

SPLINE APPROXIMATION FOR SYSTEM OF n-th ORDER NONLINEAR ORDINARY DIFFERENTIAL EQUATIONS III.

M.N. Abou El-Ela (Ismailia, Egypt)

Abstract. A spline method for approximating solution of system of nonlinear ordinary differential equations

$$x_j^{(n)} = f_j(t, x_1, x_1', x_1'', \dots, x_m, x_m', x_m''), \quad x_j^{(i)}(t_0) = x_{j,0}^{(i)},$$

where $f_j \in C([0, 1] \times R^{3m})$, $j = 1, 2, \dots, m$ and $i = 0(1)n - 1$ is presented. Errors of the method are estimated.

1. Introduction

Consider the following system of first order differential equations

$$(1.1) \quad \begin{aligned} y' &= f_1(x, y, z), & y(0) &= y_0 \\ z' &= f_2(x, y, z), & z(0) &= z_0 \end{aligned}$$

where $f_1, f_2 \in C([0, B] \times R^2)$ and satisfy the Lipschitz condition:

$$\begin{aligned} |f_i(x, y_1, z_1) - f_i(x, y_2, z_2)| &\leq A\{|y_1 - y_2| + |z_1 - z_2|\}, \quad \forall(x, y_1, z_1), \\ & \quad (x, y_2, z_2) \in [0, B] \times R^2 \quad \text{and} \quad i = 1, 2. \end{aligned}$$

Micula [2,3] has found approximate solutions for the Cauchy problem (1.1) which are polynomial splines of degree m . He also discussed the existence and uniqueness of the approximate solution and the convergence of spline approximation to the exact solution as $h \rightarrow 0$.

J. Györfvári [1] has used modified Lacunary spline functions of type $(0, 2, 3)$ for finding an approximate solution for the following Cauchy problem:

$$(1.2) \quad y''(x) = f(x, y(x), y'(x)), \quad y(0) = y_0, \quad y'(0) = y'_0, \quad x \in [0, 1],$$

where $f(x, y, y') \in C^3([0, 1] \times R^2)$, that is $y(x) \in C^5([0, 1])$, and

$$|f^{(q)}(x, y_1, y'_1) - f^{(q)}(x, y_2, y'_2)| \leq L\{|y_2 - y_1| + |y'_2 - y'_1|\}, \quad q = 0, 1, 2, 3.$$

Th. Fawzy and Samia Soliman [4-6] have discussed a spline method for finding the approximate solution of the two systems of n -th order nonlinear ordinary differential equations:

$$(1.3) \quad \begin{aligned} y^{(n)} &= f_1(x, y, z), & y^{(i)}(x_0) &= y_0^{(i)}, \\ z^{(n)} &= f_2(x, y, z), & z^{(i)}(x_0) &= z_0^{(i)}, \end{aligned}$$

where $f_1, f_2 \in C([0, 1] \times R^2)$, $f_1, f_2 \in C^r([0, 1] \times R^2)$, $i = 0(1)n - 1$;

$$(1.4) \quad \begin{aligned} y^{(n)} &= f_1(x, y, y', z, z'), & y^{(i)}(x_0) &= y_0^{(i)}, \\ z^{(n)} &= f_2(x, y, y', z, z'), & z^{(i)}(x_0) &= z_0^{(i)}, \end{aligned}$$

where $f_1, f_2 \in C([0, 1] \times R^4)$, $f_1, f_2 \in C^r([0, 1] \times R^4)$, $i = 0(1)n - 1$. They also proved the stability of the method.

In this paper, we introduce a method for approximating the solution of the system of nonlinear ordinary differential equations of n -th order:

$$(1.5) \quad x_j^{(n)} = f_j(t, x_1, x_1', x_1'', \dots, x_m, x_m', x_m''), \quad x_j^{(i)}(t_0) = x_{j,0}^{(i)},$$

where $f_j \in C([0, 1] \times R^{3m})$, $j = 1, 2, \dots, m$ and $i = 0(1)n - 1$.

The method introduced here is a one-step method $O(h^{n+\alpha})$ in $x_1^{(i)}(t), x_2^{(i)}(t), \dots, x_m^{(i)}(t)$, where $i = 0(1)n$ and $0 < \alpha \leq 1$.

2. Description of the method

Our method is to use spline functions, which are not necessarily polynomial splines, for approximating the solution $\{x_j(t) : j = 1, 2, \dots, m\}$ of the system of differential equations in consideration on the interval $[0, 1]$. These spline functions will be denoted by $S_{j,\Delta}(t)$, where $j = 1, 2, \dots, m$ and Δ is the mesh point

$$\Delta : 0 = t_0 < t_1 < \dots < t_k < t_{k+1} < \dots < t_N = 1$$

and

$$t_{k+1} - t_k = h < 1, \quad \forall k = 0(1)N - 1.$$

Let L_j be the Lipschitz constants satisfied by the functions f_j , i.e.,

$$|f_j(t, x_1, x_1', x_1'', \dots, x_m, x_m', x_m'') - f_j(t, X_1, X_1', X_1'', \dots, X_m, X_m', X_m'')| <$$

$$(2.1) \quad < L_j \sum_{p=0}^2 \left(\sum_{j=1}^m |x_j^{(p)} - X_j^{(p)}| \right)$$

$$\forall (t, x_1, x'_1, x''_1, \dots, x_m, x'_m, x''_m), (t, X_1, X'_1, X''_1, \dots, X_m, X'_m, X''_m) \in$$

$$(2.2) \quad \in [0, 1] \times R^{3m}$$

and $j = 1, 2, \dots, m$.

Then, we define the spline functions approximating the solution $\{x_j(t) : j = 1, 2, \dots, m\}$ by $S_{j,\Delta}(t)$, where

$$(2.3) \quad S_{j,\Delta}(t) = \begin{cases} S_{j,0}(t), & t_0 \leq t \leq t_1, \\ S_{j,k}(t), & t_k \leq t \leq t_{k+1}, k = 1(1)N - 1. \end{cases}$$

$S_{j,\Delta}(t)$ is given by the following expression

$$S_{j,0}(t) = \sum_{\ell=0}^{n-1} \frac{1}{\ell!} x_{j,0}^{(\ell)}(t-t_0)^\ell + \int_{t_0}^t \dots \int_{t_0}^{\theta_{n-1}} f_j \left(\theta_n, x_{1,0}^*(\theta_n), \right.$$

$$(2.4) \quad \text{n-times}$$

$$\left. x_{1,0}^{***}(\theta_n), x_{1,0}^{****}(\theta_n), \dots, x_{m,0}^*(\theta_n), x_{m,0}^{**}(\theta_n), x_{m,0}^{***}(\theta_n) \right) d\theta_n \dots d\theta_1,$$

$$S_{j,k}(t) = \sum_{\ell=0}^{n-1} \frac{1}{\ell!} S_{j,k-1}^{(\ell)}(t_k)(t-t_k)^\ell + \int_{t_k}^t \dots \int_{t_k}^{\theta_{n-1}} f_j \left(\theta_n, S_{1,k-1}^*(\theta_n), \right.$$

$$(2.5) \quad \text{n-times}$$

$$\left. S_{1,k-1}^{**}(\theta_n), S_{1,k-1}^{***}(\theta_n), \dots, S_{m,k-1}^*(\theta_n), S_{m,k-1}^{**}(\theta_n), S_{m,k-1}^{***}(\theta_n) \right) d\theta_n \dots d\theta_1,$$

where

$$(2.6) \quad x_{j,0}^*(\theta) = \sum_{\ell=0}^n \frac{1}{\ell!} x_{j,0}^{(\ell)}(\theta-t_0)^\ell,$$

$$x'_{j,0}{}^{**}(\theta) = \sum_{\ell=0}^{n-2} \frac{1}{\ell!} x_{j,0}^{(\ell+1)}(\theta - t_0)^\ell + \int_{t_0}^{\theta} \dots \int_{t_0}^{\theta_{n-2}} f_j \left(\theta_{n-1}, x_{1,0}^*(\theta_{n-1}), \right.$$

n-1 times

(2.7)

$$\left. x_{1,0}^*(\theta_{n-1}), x_{1,0}''^*(\theta_{n-1}), \dots, x_{m,0}^*(\theta_{n-1}), x_{m,0}'^*(\theta_{n-1}), x_{m,0}''^*(\theta_{n-1}) \right) d\theta_{n-1} \dots d\theta_1,$$

$$x''_{j,0}{}^{***}(\theta) = \sum_{\ell=0}^{n-3} \frac{1}{\ell!} x_{j,0}^{(\ell+2)}(\theta - t_0)^\ell + \int_{t_0}^{\theta} \dots \int_{t_0}^{\theta_{n-3}} f_j \left(\theta_{n-2}, x_{1,0}''^*(\theta_{n-2}), \right.$$

n-2 times

(2.8)

$$\left. x_{1,0}''^*(\theta_{n-2}), x_{1,0}''''^*(\theta_{n-2}), \dots, x_{m,0}''^*(\theta_{n-2}), x_{m,0}''''^*(\theta_{n-2}), x_{m,0}''''^*(\theta_{n-2}) \right) d\theta_{n-2} \dots d\theta_1.$$

Also,

$$S'_{j,k-1}(\theta) = \sum_{\ell=0}^{n-1} \frac{1}{\ell!} S_{j,k-1}^{(\ell)}(t_k)(\theta - t_k)^\ell + \frac{1}{n!} f_j \left(t_k, S_{1,k-1}(t_k), \right.$$

$$(2.9) \quad \left. S'_{1,k-1}(t_k), S''_{1,k-1}(t_k), \dots, S_{m,k-1}(t_k), S'_{m,k-1}(t_k), S''_{m,k-1}(t_k) \right) (\theta - t_k)^n,$$

$$S'_{j,k-1}{}^{**}(\theta) = \sum_{\ell=0}^{n-2} \frac{1}{\ell!} S_{j,k-1}^{(\ell+1)}(t_k)(\theta - t_k)^\ell + \int_{t_k}^{\theta} \dots \int_{t_k}^{\theta_{n-2}} f_j \left(\theta_{n-1}, S_{1,k-1}^*(\theta_{n-1}), \right.$$

(2.10)

n-1 times

$$\left. S_{1,k-1}^*(\theta_{n-1}), S_{1,k-1}''^*(\theta_{n-1}), \dots, S_{m,k-1}^*(\theta_{n-1}), S_{m,k-1}'^*(\theta_{n-1}), S_{m,k-1}''^*(\theta_{n-1}) \right) \cdot d\theta_{n-1} \dots d\theta_1,$$

$$S''_{j,k-1}{}^{***}(\theta) = \sum_{\ell=0}^{n-3} \frac{1}{\ell!} S_{j,k-1}^{(\ell+2)}(t_k)(\theta - t_k)^\ell + \int_{t_k}^{\theta} \dots \int_{t_k}^{\theta_{n-3}} f_j \left(\theta_{n-2}, S_{1,k-1}''^*(\theta_{n-2}), \right.$$

(2.11)

n-2 times

$$\left. S_{1,k-1}''^*(\theta_{n-2}), S_{1,k-1}''''^*(\theta_{n-2}), \dots, S_{m,k-1}''^*(\theta_{n-2}), S_{m,k-1}''''^*(\theta_{n-2}), \right.$$

$$S''_{m,k-1}(\theta_{n-2}) \Big) d\theta_{n-2} \dots d\theta_1.$$

By this construction it is clear that $S_{j,\Delta}(t) \in C^{n-1}[0, 1]$, $j = 1, 2, \dots, m$. Also, we need the following definitions:

Definition 2.1. Let $A = [a_{ij}]$ and $B = [b_{ij}]$ be two matrices of the same order; then we say that $A \leq B$ iff:

- (i) a_{ij} and b_{ij} are non-negative numbers;
- (ii) $a_{ij} \leq b_{ij}$ for all i, j .

Definition 2.2. Let $A = [a_{ij}]$ be an $m \times n$ matrix, then the norm of A is denoted by $\|A\|$ and is defined by

$$\|A\| = \max_i \sum_{j=1}^n |a_{ij}|.$$

Definition 2.3. Let $f(t)$ be defined on an interval I and suppose we can find two positive constants M_0 and α such that

$$|f(t_1) - f(t_2)| \leq M_0 |t_1 - t_2|^\alpha \quad \text{for all } t_1, t_2 \in I.$$

Then f is said to satisfy a Lipschitz condition of order α . The class of such functions will be designated by $\text{Lip } M_0^\alpha$.

3. Error estimates

For this purpose it is convenient to write the exact solution $\{x_j : j = 1, 2, \dots, m\}$ in special forms as follows:

For all $t \in [t_0, t_1]$ the exact solution of (1.5) can be written by means of Taylor's expansion in the forms:

$$x_j(t) = \sum_{\ell=0}^{n-1} \frac{1}{\ell!} x_{j,0}^{(\ell)}(t-t_0)^\ell + \int_{t_0}^t \dots \int_{t_0}^{\theta_{n-1}} f_j(\theta_n, \bar{x}_{1,0}(\theta_n), \bar{x}_{1,0}'(\theta_n),$$

(3.1) n times

$$\bar{x}_{1,0}''(\theta_n), \dots, \bar{x}_{m,0}(\theta_n), \bar{x}_{m,0}'(\theta_n), \bar{x}_{m,0}''(\theta_n)) d\theta_n \dots d\theta_1,$$

where

$$\bar{x}_{j,0}(\theta) = \sum_{\ell=0}^{n-1} \frac{1}{\ell!} x_{j,0}^{(\ell)}(\theta - t_0)^\ell + \frac{1}{n!} x_j^{(n)}(\xi_{j,0})(\theta - t_0)^n,$$

$$(3.2) \quad \xi_{j,0} \in (t_0, t_1),$$

$$\bar{x}'_{j,0}(\theta) = \sum_{\ell=0}^{n-2} \frac{1}{\ell!} x_{j,0}^{(\ell+1)}(\theta - t_0)^\ell + \int_{t_0}^{\theta} \dots \int_{t_0}^{\theta_{n-2}} f_j \left(\theta_{n-1}, \bar{x}_{1,0}(\theta_{n-1}), \right.$$

n-1 times

$$(3.3)$$

$$\left. \bar{x}'_{1,0}(\theta_{n-1}), \bar{x}''_{1,0}(\theta_{n-1}), \dots, \bar{x}_{m,0}(\theta_{n-1}), \bar{x}'_{m,0}(\theta_{n-1}), \bar{x}''_{m,0}(\theta_{n-1}) \right) d\theta_{n-1} \dots d\theta_1,$$

$$\bar{x}''_{j,0}(\theta) = \sum_{\ell=0}^{n-3} \frac{1}{\ell!} x_{j,0}^{(\ell+2)}(\theta - t_0)^\ell + \int_{t_0}^{\theta} \dots \int_{t_0}^{\theta_{n-3}} f_j \left(\theta_{n-2}, \bar{x}_{1,0}(\theta_{n-2}), \right.$$

n-2 times

$$(3.4)$$

$$\left. \bar{x}''_{1,0}(\theta_{n-2}), \bar{x}'''_{1,0}(\theta_{n-2}), \dots, \bar{x}_{m,0}(\theta_{n-2}), \bar{x}'_{m,0}(\theta_{n-2}), \bar{x}''_{m,0}(\theta_{n-2}) \right) d\theta_{n-2} \dots d\theta_1.$$

3.1 Estimation of $|X_j^{(i)}(t) - S_{j,0}^{(i)}(t)|$

From equations (2.4), (3.1) and applying Lipschitz condition (2.1) we have

$$|X_j^{(i)}(t) - S_{j,0}^{(i)}(t)| \leq L_j \int_{t_0}^t \dots \int_{t_0}^{\theta_{n-i-1}} \left\{ \sum_{j=1}^m |\bar{X}_{j,0}(\theta_{n-i}) - X_{j,0}^*(\theta_{n-i})| + \right.$$

$$(3.1.1)$$

n-i times

$$\left. + \sum_{j=1}^m |\bar{X}'_{j,0}(\theta_{n-i}) - X'_{j,0}(\theta_{n-i})| + \sum_{j=1}^m |\bar{X}''_{j,0}(\theta_{n-i}) - X''_{j,0}(\theta_{n-i})| \right\} d\theta_{n-i} \dots d\theta_1.$$

Equations (2.6) and (3.2) at $\theta = \theta_{n-i}$ show that

$$(3.1.2) \quad \left| \bar{X}_{j,0}^{(p)}(\theta_{n-i}) - X_{j,0}^{(p)*}(\theta_{n-i}) \right| \leq \frac{1}{(n-p)!} w(X_j^{(n)}, h) |\theta_{n-i} - t_0|^{n-p},$$

$$p = 0, 1, 2$$

where $w(X_j^{(n)}, h)$ is the modulus of continuity of the function $X_j^{(n)}(t)$. By using equations (2.7), (3.3) and (3.1.2) we get:

$$\left| \bar{X}_{j,0}^{(p)*}(\theta_{n-2}) - X_{j,0}^{(p)**}(\theta_{n-2}) \right| \leq mL_j w(h) \left[\sum_{r=0}^2 \frac{|\theta_{n-2} - t_0|^{2n-p-r}}{(2n-p-r)!} \right],$$

$$(3.1.3) \quad p = 0, 1, 2$$

where $w(h) = \max\{w(x_j^{(n)}, h) : j = 1, 2, \dots, m\}$. Using equations (2.8), (3.4) and (3.1.3) we get:

$$(3.1.4) \quad \left| \bar{X}_{j,0}^{(p)**}(\theta_{n-i}) - X_{j,0}^{(p)***}(\theta_{n-i}) \right| \leq mw(h) L_j \sum_{j=1}^m L_j \left[\frac{|\theta_{n-i} - t_0|^{3n-2}}{(3n-2)!} + \right. \\ \left. 2 \frac{|\theta_{n-i} - t_0|^{3n-3}}{(3n-3)!} + 3 \frac{|\theta_{n-i} - t_0|^{3n-4}}{(3n-4)!} + 2 \frac{|\theta_{n-i} - t_0|^{3n-5}}{(3n-5)!} + \right. \\ \left. + \frac{|\theta_{n-i} - t_0|^{3n-6}}{(3n-6)!} \right].$$

Substituting from equations (3.1.2), (3.1.3) and (3.1.4) in equation (3.1.1) and calculating the integrals we get:

$$(3.1.5) \quad \left| X_j^{(i)}(t) - S_{j,0}^{(i)}(t) \right| \leq mL_j \left\{ \frac{1}{(2n-i)!} + \left(\sum_{j=1}^m L_j \right) \left[\sum_{r=1}^3 \frac{h^{n-r}}{(3n-r-i)!} \right] + \right. \\ \left. + \left(\sum_{j=1}^m L_j \right)^2 \left[\frac{h^{2n-2}}{(4n-2-i)!} + \frac{2h^{2n-3}}{(4n-3-i)!} + \frac{3h^{2n-4}}{(4n-4-i)!} + \frac{2h^{2n-5}}{(4n-5-i)!} + \right. \right. \\ \left. \left. + \frac{h^{2n-6}}{(4n-6-i)!} \right] \right\} w(h) h^{2n-i} \leq \mu_{j,0} w(h) h^{2n-i} = O(h^{2n+\alpha-i}),$$

where $i = 0(1)n, 0 < \alpha \leq 1$, and

$$\begin{aligned} \mu_{j,0} = m L_j \left\{ \frac{1}{(2n-i)!} + \sum_{j=1}^m L_j \sum_{r=1}^3 \frac{1}{(3n-i-r)!} + \right. \\ \left. + \left(\sum_{j=1}^m L_j \right)^2 \left[\frac{1}{(4n-i-2)!} + \frac{2}{(4n-i-3)!} + \frac{3}{(4n-i-4)!} + \right. \right. \\ \left. \left. + \frac{2}{(4n-i-5)!} + \frac{1}{(4n-i-6)!} \right] \right\} \end{aligned}$$

is a constant independent of h .

3.2. Estimation of $|X_j^{(i)}(t) - S_{j,k}^{(i)}(t)|$

Finally we study the approximations on the general subinterval $[t_k, t_{k+1}]$, $k = 1, \dots, N-1$. For this purpose, the exact solution of (1.5) can be written in the following form:

$$X_j(t) = \sum_{\ell=0}^{n-1} \frac{1}{\ell!} X_{j,k}^{(\ell)}(t-t_k)^\ell + \int_{t_k}^t \dots \int_{t_k}^{\theta_{n-1}} f_j(\theta_n, \bar{X}_{1,k}(\theta_n),$$

n times

$$(3.2.1) \quad \bar{X}'_{1,k}(\theta_n), \bar{X}''_{1,k}(\theta_n), \dots, \bar{X}_{m,k}(\theta_n), \bar{X}'_{m,k}(\theta_n), \bar{X}''_{m,k}(\theta_n)) d\theta_n \dots d\theta_1,$$

where

$$\bar{X}_{j,k}(\theta) = \sum_{\ell=0}^{n-1} \frac{1}{\ell!} X_{j,k}^{(\ell)}(\theta-t_k)^\ell + \frac{1}{n!} X_j^{(n)}(\xi_{j,k})(\theta-t_k)^n,$$

$$(3.2.2) \quad \xi_{j,k} \in (t_k, t_{k+1})$$

$$\bar{X}'_{j,k}(\theta) = \sum_{\ell=0}^{n-2} \frac{1}{\ell!} X_{j,k}^{(\ell+1)}(\theta-t_k)^\ell + \int_{t_k}^{\theta} \dots \int_{t_k}^{\theta_{n-2}} f_j(\theta_{n-1}, \bar{X}_{1,k}(\theta_{n-1}),$$

$$(3.2.3) \quad \text{n-1 times}$$

$$\bar{X}'_{1,k}(\theta_{n-1}), \bar{X}''_{1,k}(\theta_{n-1}), \dots, \bar{X}_{m,k}(\theta_{n-1}), \bar{X}'_{m,k}(\theta_{n-1}), \bar{X}''_{m,k}(\theta_{n-1})) d\theta_{n-1} \dots d\theta_1,$$

$$\bar{X}_{j,k}''''(\theta) = \sum_{\ell=0}^{n-3} \frac{1}{\ell!} X_{j,k}^{(\ell+2)}(\theta - t_k)^\ell + \int_{t_k}^{\theta} \dots \int_{t_k}^{\theta_{n-3}} f_j(\theta_{n-2}, \bar{X}_{1,k}^*(\theta_{n-2}),$$

(3.2.4) n-2 times

$$\bar{X}_{1,k}^*(\theta_{n-2}), \bar{X}_{1,k}''(\theta_{n-2}), \dots, \bar{X}_{m,k}^*(\theta_{n-2}), \bar{X}_{m,k}'(\theta_{n-2}), \bar{X}_{m,k}''(\theta_{n-2})) d\theta_{n-2} \dots d\theta_1.$$

Next, we proceed to prove the convergence. Using equations (2.5), (3.2.1) we get

$$\begin{aligned} |X_j^{(i)}(t) - S_{j,k}^{(i)}(t)| &\leq \sum_{\ell=0}^{n-i-1} \frac{1}{\ell!} |X_{j,k}^{(i+\ell)} - S_{j,k-1}^{(i+\ell)}(t_k)| |t - t_k|^\ell + \\ &+ L_j \int_{t_k}^t \dots \int_{t_k}^{\theta_{n-i-1}} \left\{ \sum_{j=1}^m |\bar{X}_{j,k}(\theta_{n-i}) - S_{j,k-1}^*(\theta_{n-i})| + \right. \\ &\quad \left. + \sum_{j=1}^m |\bar{X}_{j,k}'(\theta_{n-i}) - S_{j,k-1}'(\theta_{n-i})| + \sum_{j=1}^m |\bar{X}_{j,k}''(\theta_{n-i}) - S_{j,k-1}''(\theta_{n-i})| \right\} \cdot \\ &\quad \cdot d\theta_{n-i} \dots d\theta_1. \end{aligned}$$

(3.2.5) n-i times

Using (2.9) and (3.2.2) we get

$$\begin{aligned} &|\bar{X}_{j,k}(\theta_{n-i}) - S_{j,k-1}^*(\theta_{n-i})| \leq \\ (3.2.6) \quad &\leq \sum_{\ell=0}^{n-1} \frac{1}{\ell!} e_{j,k}^{(\ell)} |\theta_{n-i} - t_k|^\ell + \frac{1}{n!} N_{j,1} |\theta_{n-i} - t_k|^n, \end{aligned}$$

where

$$\begin{aligned} N_{j,1} \leq &|X_j^{(n)}(\xi_{j,k}) - X_{j,k}^{(n)}| + |f_j(t_k, X_{1,k}, X'_{1,k}, X''_{1,k}, \dots, X_{m,k}, X'_{m,k}, X''_{m,k}) - \\ &- f_j(t_k, S_{1,k-1}(t_k), S'_{1,k-1}(t_k), S''_{1,k-1}(t_k), \dots, S_{m,k-1}(t_k), \\ &S'_{m,k-1}(t_k), S''_{m,k-1}(t_k))|. \end{aligned}$$

Applying Lipschitz condition (2.1) for the second term

$$(3.2.7) \quad N_{j,1} \leq w(X_j^{(n)}, h) + L_j \sum_{k=1}^m (e_{j,k} + \dot{e}_{j,k} + \ddot{e}_{j,k}),$$

where $e_{j,k}^{(i)} = |X_{j,k}^{(i)} - S_{j,k-1}^{(i)}|$ is the estimated error of $X_{j,k}^{(i)}$ at any point $t_k \in [0, 1]$, $i = 0, 1, 2$.

Using (3.2.7) in (3.2.6)

$$\begin{aligned} & \left| \bar{X}_{j,k}^{(p)}(\theta_{n-i} - S_{j,k-1}^{(p)*}(\theta_{n-i})) \right| \leq \sum_{\ell=0}^{n-p-1} \frac{1}{\ell!} e_{j,k}^{(\ell+p)} |\theta_{n-i} - t_k|^\ell + \\ & + \frac{1}{(n-p)!} \left[w(X_j^{(n)}, h) + L_j \sum_{k=1}^m (e_{j,k} + \dot{e}_{j,k} + \ddot{e}_{j,k}) \right] |\theta_{n-i} - t_k|^{n-p}, \end{aligned}$$

$$(3.2.8) \quad p = 0, 1, 2.$$

From (2.10), (3.2.3) and (3.2.8) we have

$$\begin{aligned} & \left| \bar{X}_{j,k}^{(p)*}(\theta_{n-i}) - S_{j,k-1}^{(p)**}(\theta_{n-i}) \right| \leq \sum_{\ell=0}^{n-p-1} \frac{1}{\ell!} e_{j,k}^{(\ell+p)} |\theta_{n-i} - t_k|^\ell + \\ (3.2.9) \quad & + L_j \left[\sum_{q=0}^2 \left(\sum_{\ell=0}^{n-q-1} \frac{1}{(\ell+n-p)!} \sum_{j=1}^m e_{j,k}^{(\ell+q)} \right) \right] |\theta_{n-i} - t_k|^{\ell+n-p} + \\ & L_j [mw(h) + \sum_{j=1}^m L_j \sum_{k=1}^m (e_{j,k} + \dot{e}_{j,k} + \ddot{e}_{j,k})] \sum_{q=0}^2 \frac{|\theta_{n-i} - t_k|^{2n-p-q}}{(2n-p-q)!}, \\ & p = 0, 1, 2. \end{aligned}$$

From (2.11), (3.2.4) and (3.2.9) we have:

$$\begin{aligned} & \left| \bar{X}_{j,k}^{(p)**}(\theta_{n-i}) - S_{j,k-1}^{(p)***}(\theta_{n-i}) \right| \leq \sum_{\ell=0}^{n-3} \frac{1}{\ell!} e_{j,k}^{(\ell+2)} |\theta_{n-i} - t_k|^\ell + \\ & + L_j \sum_{q=0}^2 \left\{ \sum_{\ell=0}^{n-q-1} \frac{1}{(\ell+n-2)!} \sum_{j=1}^m e_{j,k}^{(\ell+q)} \right\} |\theta_{n-i} - t_k|^{\ell+n-2} + \end{aligned}$$

$$\begin{aligned}
& + L_j \sum_{j=1}^m L_j \sum_{r=2}^4 \left\{ \sum_{q=0}^2 \sum_{\ell=0}^{n-q-1} \frac{1}{(\ell + 2n - r)!} \sum_{j=1}^m e_{j,k}^{(\ell+q)} \right\} \cdot |\theta_{n-i} - t_k|^{\ell+2n-r} + \\
(3.2.10) \quad & + L_j \sum_{j=1}^m L_j \{mw(h) + \sum_{j=1}^m L_j \sum_{j=1}^m (e_{j,k} + \dot{e}_{j,k} + \ddot{e}_{j,k})\} \times \\
& \times \left\{ \frac{|\theta_{n-i} - t_k|^{3n-2}}{(3n-2)!} + \frac{2|\theta_{n-i} - t_k|^{3n-3}}{(3n-3)!} + \frac{3|\theta_{n-i} - t_k|^{3n-4}}{(3n-4)!} + \right. \\
& \left. + \frac{2|\theta_{n-i} - t_k|^{3n-5}}{(3n-5)!} + \frac{|\theta_{n-i} - t_k|^{3n-6}}{(3n-6)!} \right\}.
\end{aligned}$$

Substituting from (3.2.8), (3.2.9) and (3.2.10) in (3.2.5)

$$\begin{aligned}
e_j^{(i)}(t) & \leq \sum_{\ell=0}^{n-i-1} \frac{1}{\ell!} e_{j,k}^{(i+\ell)} h^\ell + L_j \sum_{q=0}^2 \sum_{\ell=0}^{n-q-1} \left[\frac{h^{\ell+n-i}}{(\ell+n-i)!} + \right. \\
& + \sum_{j=1}^m L_j \sum_{r=1}^2 \frac{h^{\ell+2n-i-r}}{(\ell+2n-i-r)!} + \left. \left(\sum_{j=1}^m L_j \right)^2 \sum_{r=2}^4 \frac{h^{\ell+3n-i-r}}{(\ell+3n-i-r)!} \right] \sum_{j=1}^m e_{j,k}^{(\ell+q)} + \\
(3.2.11) \quad & + L_j \sum_{j=1}^m L_j \sum_{j=1}^m (e_{j,k} + \dot{e}_{j,k} + \ddot{e}_{j,k}) \left[\frac{h^{2n-i}}{(2n-i)!} + \right. \\
& + \sum_{j=1}^m L_j \sum_{r=1}^3 \frac{h^{3n-i-r}}{(3n-i-r)!} + \left. \left(\sum_{j=1}^m L_j \right)^2 \left(\frac{h^{4n-i-2}}{(4n-i-2)!} + \frac{2h^{4n-i-3}}{(4n-i-3)!} + \right. \right. \\
& \left. \left. + \frac{3h^{4n-i-4}}{(4n-i-4)!} + \frac{2h^{4n-i-5}}{(4n-i-5)!} + \frac{h^{4n-i-6}}{(4n-i-6)!} \right) \right] + C_{i,j} w(h) h^{2n-i},
\end{aligned}$$

where

$$\begin{aligned}
C_{i,j} & = mL_j \left[\frac{1}{(2n-i)!} + \sum_{r=1}^3 \frac{1}{(3n-i-r)!} \sum_{j=1}^m L_j + \left(\frac{1}{(4n-i-2)!} + \right. \right. \\
& \left. \left. + \frac{2}{(4n-i-3)!} + \frac{3}{(4n-i-4)!} + \frac{2}{(4n-i-5)!} + \frac{1}{(4n-i-6)!} \right) \left(\sum_{j=1}^m L_j \right)^2 \right]
\end{aligned}$$

is a constant independent of h and $i = 0(1)n-1$, $j = 1, 2, \dots, m$.

Hence, the inequality (3.2.11) can be written in general form as

$$(3.2.12) \quad e_j^{(i)}(t) \leq \sum_{\bar{j}=1}^m \left(\sum_{\ell=0}^{n-1} Q_{j,\bar{j},i,\ell} e_{j,\bar{j},k}^{(\ell)} \right) + C_{i,j} w(h) h^{2n-i},$$

where $i = 0(1)n - 1$, and $j = 1, 2, \dots, m$.

$$Q_{j,\bar{j},i,\ell} = \begin{cases} (1 + q_{j,\bar{j},i,\ell} h) & \text{if } j = \bar{j} \text{ and } i = \ell \\ q_{j,\bar{j},i,\ell} h^{\ell-i} & \text{if } j = \bar{j} \text{ and } i < \ell \\ q_{j,\bar{j},i,\ell} h^{n-i} & \text{if } j = \bar{j}, i > \ell \text{ and } \ell = 0, 1, 2 \\ q_{j,\bar{j},i,\ell} h^{n+\ell-i-2} & \text{if } j = \bar{j}, i > \ell \text{ and } \ell \geq 3 \\ q_{j,\bar{j},i,\ell} h^{n-i} & \text{if } j \neq \bar{j} \text{ and } \ell = 0, 1, 2 \\ q_{j,\bar{j},i,\ell} h^{n+\ell-i-2} & \text{if } j \neq \bar{j} \text{ and } \ell \geq 3 \end{cases}$$

If we use the definition 2.1 and since $h < 1$, from the inequality (3.2.12),

$$\begin{bmatrix} \bar{e}_1(t) \\ \bar{e}_2(t) \\ \vdots \\ \bar{e}_m(t) \end{bmatrix} \leq \left(\begin{bmatrix} M_{1,1} M_{1,2} \dots M_{1,m} \\ M_{2,1} M_{2,2} \dots M_{2,m} \\ \vdots \\ M_{m,1} M_{m,2} \dots M_{m,m} \end{bmatrix} + h \begin{bmatrix} \bar{Q}_{1,1}^* \bar{Q}_{1,2}^* \dots \bar{Q}_{1,m}^* \\ \bar{Q}_{2,1}^* \bar{Q}_{2,2}^* \dots \bar{Q}_{2,m}^* \\ \vdots \\ \bar{Q}_{m,1}^* \bar{Q}_{m,2}^* \dots \bar{Q}_{m,m}^* \end{bmatrix} \right) \begin{bmatrix} \bar{e}_{1,k} \\ \bar{e}_{2,k} \\ \vdots \\ \bar{e}_{m,k} \end{bmatrix} + w(h) h^{n+1} \begin{bmatrix} \bar{C}_1^* \\ \bar{C}_2^* \\ \vdots \\ \bar{C}_m^* \end{bmatrix}$$

where

$$\bar{e}_j(t) = \begin{bmatrix} e_j(t) \\ \dot{e}_j(t) \\ \vdots \\ e_j^{(n-1)}(t) \end{bmatrix}, \quad \bar{e}_{j,k} = \begin{bmatrix} e_{j,k} \\ \dot{e}_{j,k} \\ \vdots \\ e_{j,k}^{(n-1)} \end{bmatrix}, \quad \bar{C}_j^* = \begin{bmatrix} C_{0,j} \\ C_{1,j} \\ \vdots \\ C_{n-1,j} \end{bmatrix}$$

$$\bar{Q}_{j,\bar{j}}^* = \begin{bmatrix} q_{j,\bar{j},0,0} & q_{j,\bar{j},0,1} & q_{j,\bar{j},0,n-1} \\ q_{j,\bar{j},1,0} & q_{j,\bar{j},1,1} & q_{j,\bar{j},1,n-1} \\ \vdots & \vdots & \vdots \\ q_{j,\bar{j},n-1,0} & q_{j,\bar{j},n-1,1} & q_{j,\bar{j},n-1,n-1} \end{bmatrix} \quad \forall j, \bar{j} = 1, 2, \dots, m$$

and

$$M_{j,\bar{j}} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \vdots & \vdots & \vdots \\ 0 & 0 & 1 \end{bmatrix} \quad \text{if } j = \bar{j}, \quad M_{j,\bar{j}} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \vdots & \vdots & \vdots \\ 0 & 0 & 0 \end{bmatrix} \quad \text{if } j \neq \bar{j}.$$

Then

$$(3.2.13) \quad \bar{E}(t) \leq (I + h\bar{Q})\bar{E}_k + w(h)h^{n+1}\bar{C},$$

where

$$\begin{aligned} \bar{E}(t) &= (\bar{e}_1(t) \ \bar{e}_2(t) \ \dots \ \bar{e}_m(t))^T, \\ E_k &= (\bar{e}_{1,k} \ \bar{e}_{2,k} \ \dots \ \bar{e}_{m,k})^T, \\ \bar{C} &= (\bar{C}_1^* \ \bar{C}_2^* \ \dots \ \bar{C}_m^*)^T \end{aligned}$$

Q is the $(mn \times mn)$ matrix whose elements are constants independent of h , I is the mn -th order unit matrix and \bar{C} is the $(mn \times 1)$ matrix whose elements are constants independent of h .

By using definition 2.2 of the matrix norm

$$(3.2.14) \quad \|\bar{E}(t)\| \leq (1 + h \|\bar{Q}\|) \|\bar{E}_k\| + w(h)h^{n+1} + \|\bar{C}\|.$$

Since (3.2.13) is valid for all $t \in [t_k, t_{k+1}]$, then the following inequalities hold true:

$$\begin{aligned} \|\bar{E}(t)\| &\leq (1 + h \|\bar{Q}\|) \|\bar{E}_k\| + w(h)h^{n+1} \|\bar{C}\|, \\ (1 + h \|\bar{Q}\|) \|\bar{E}_k\| &\leq (1 + h \|\bar{Q}\|)^2 \|\bar{E}_{k-1}\| + (1 + h \|\bar{Q}\|)w(h)h^{n+1} \|\bar{C}\|, \end{aligned}$$

$$\begin{aligned} &(1 + h \|\bar{Q}\|)^2 \|\bar{E}_{k-1}\| \leq \\ &\leq (1 + h \|\bar{Q}\|)^3 \|\bar{E}_{k-2}\| + (1 + h \|\bar{Q}\|)^2 w(h)h^{n+1} \|\bar{C}\|, \end{aligned}$$

$$\begin{aligned} &(1 + h \|\bar{Q}\|)^k \|\bar{E}_1\| \leq \\ &\leq (1 + h \|\bar{Q}\|)^{k+1} \|\bar{E}_0\| + (1 + h \|\bar{Q}\|)^k w(h)h^{n+1} \|\bar{C}\|. \end{aligned}$$

Then, from these inequalities, and noting that $\|\bar{E}_0\| = 0$, we get

$$\begin{aligned} \|\bar{E}(t)\| &\leq w(h)h^{n+1} \|\bar{C}\| \sum_{\ell=0}^k (1 + h \|\bar{Q}\|)^\ell = \\ &= w(h)h^n \frac{\|\bar{C}\|}{\|\bar{Q}\|} [(1 + h \|\bar{Q}\|)^{k+1} - 1] \leq w(h)h^n \frac{\|\bar{C}\|}{\|\bar{Q}\|} \left[\left(1 + \frac{1}{N} \|\bar{Q}\|\right)^N - 1 \right] \end{aligned}$$

$$(3.2.15) \quad \leq w(h)h^n \frac{\|\bar{C}\|}{\|\bar{Q}\|} (e^{\|\bar{Q}\|} - 1).$$

$$\| \bar{E}(t) \| \leq \mu_1 w(h) h^n,$$

where $\mu_1 = \frac{\|\bar{C}\|}{\|Q\|} (e^{\|Q\|} - 1)$ is a constant independent of h . Thus, by using the definition 2.3, we get

$$(3.2.16) \quad e_j^{(i)}(t) \leq \mu_1 w(h) h^n = O(h^{n+\alpha}),$$

where $i = 0(1)n - 1$, $j = 1, 2, \dots, m$ and $0 < \alpha \leq 1$.

3. Estimation of $|X^{(n)}(t) - S_{j,k}^{(n)}(t)|$

From (2.5), (3.2.1), (3.2.8), (3.2.9) and (3.2.10), we get

$$(3.3.1) \quad \begin{aligned} e_j^{(n)}(t) = |X_j^{(n)}(t) - S_{j,k}^{(n)}(t)| &\leq L_j \sum_{q=0}^2 \sum_{\ell=0}^{n-q-1} \left[\frac{h^\ell}{\ell!} + \sum_{j=1}^m L_j \sum_{r=1}^2 \frac{h^{\ell+n-r}}{(\ell+n-r)!} + \right. \\ &+ \left. \left(\sum_{j=1}^m L_j \right)^2 \sum_{r=2}^4 \frac{h^{\ell+2n-r}}{(\ell+2n-r)!} \right] \sum_{j=1}^m e_{j,k}^{(\ell+q)} + L_j \sum_{j=1}^m L_j \sum_{j=1}^m (e_{j,k} + \\ &+ \dot{e}_{j,k} + \ddot{e}_{j,k}) \left[\frac{h^n}{n!} + \sum_{j=1}^m L_j \sum_{r=1}^3 \frac{h^{2n-r}}{(2n-r)!} + \left(\sum_{j=1}^m L_j \right)^2 \left(\frac{h^{3n-2}}{(3n-2)!} + \right. \right. \\ &+ \left. \left. \frac{2h^{3n-3}}{(3n-3)!} + \frac{3h^{3n-4}}{(3n-4)!} + \frac{2h^{3n-5}}{(3n-5)!} + \frac{h^{3n-6}}{(3n-6)!} \right) \right] + C_{n,j} w(h) h^n, \end{aligned}$$

where

$$\begin{aligned} C_{n,j} = m L_j \left[\frac{1}{n!} + \sum_{j=1}^m L_j \sum_{r=1}^3 \frac{1}{(2n-r)!} + \left(\sum_{j=1}^m L_j \right)^2 \frac{1}{(3n-2)!} + \right. \\ \left. + \frac{2}{(3n-3)!} + \frac{3}{(3n-4)!} + \frac{2}{(3n-5)!} + \frac{1}{(3n-6)!} \right] \end{aligned}$$

is a constant independent of h .

By using (3.2.16), the inequality (3.3.1) becomes

$$(3.3.2) \quad e_j^{(n)}(t) \leq \mu_{2,j} w(h) h^n = O(h^{n+\alpha}),$$

where $j = 1, 2, \dots, n$ and $\mu_{2,j}$ is a constant independent of h .

Thus, we have proved the following

Theorem 3.1 Let $S_{j,\Delta}(t)$ be the approximate solutions to the exact ones $X_j(t)$, $j = 1, 2, \dots, m$ of the problem (1.5) given by equations (2.3)-(2.5) and let $f_j \in C([t_0, t_N] \times \mathbb{R}^{3m})$, $j = 1, 2, \dots, m$. Then, for all $t \in [t_0, t_1]$, we have

$$\begin{aligned} |X_j^{(i)}(t) - S_{j,0}^{(i)}| &\leq \mu_{j,0} w(h) h^{2n-i}, \\ i &= 0(1)n \quad \text{and} \quad j = 1, 2, \dots, m \quad \text{and} \\ \mu_{j,0} &= mL_j \left\{ \frac{1}{(2n-i)!} + \sum_{j=1}^m L_j \left(\sum_{r=1}^3 \frac{1}{(3n-i-r)!} \right) + \right. \\ &+ \left. \left(\sum_{j=1}^m L_j \right)^2 \left[\frac{1}{(4n-i-2)!} + \frac{2}{(4n-i-3)!} + \frac{3}{(4n-i-4)!} + \right. \right. \\ &\left. \left. + \frac{2}{(4n-i-5)!} + \frac{1}{(4n-i-6)!} \right] \right\} \end{aligned}$$

is a constant independent of h , and for all $t \in [t_k, t_{k+1}]$, $k = 1(1)N - 1$ we have

$$|X_j^{(i)}(t) - S_{j,k}^{(i)}(t)| \leq \mu w(h) h^n,$$

where $i = 0(1)n$, $j = 1, 2, \dots, m$ and $\mu = \max\{\mu_1, \mu_{2,j} : j = 1(1)m\}$ is a constant independent of h .

References

- [1] Györfvári J., Cauchy problem and modified lacunary spline functions, *Constructive Theory of Functions' 84*, eds. B. Sendov, P. Petrushev, R. Mallev, and S. Tashev, Bulgarian Academy of Sciences, Sofia, 1984, 392-396.
- [2] Micula Gh., Approximate integration of system of differential equations by spline functions, *Studia Univ.Babes-Bolyai Ser.Math.-Mech.*, **16** (1971), Fasc. 2, 27-39.
- [3] Micula Gh., Functii spline de grad superior de aproximare a solutiilor sistemelor de ecuatii diferentiale, *Studia Univ.Babes-Bolyai Ser. Math.-Mech.*, **17** (1972), Fasc.1, 21-32.
- [4] Fawzy Th. and Samia Soliman, Spline approximation for system on n-th order ordinary differential equations I., *Studia Univ.Babes-Bolyai Ser.Math.-Mech.* (submitted)

- [5] **Fawzy Th. and Ramadan Z.**, Spline approximation for system of ordinary differential equations II., *Annales Univ.Sci.Budapest,Sect.Comp.*, **7** (1987), 53-61.
- [6] **Fawzy Th. and Samia Soliman**, Stability of the spline approximation method for solving system of n-th order ordinary differential equations (submitted)

(Received December 1, 1990)

M.N. Abou El-Ela

Dept. of Math., Faculty of Science,

Suez Canal University

Ismailia, Egypt