

A CHARACTERIZATION OF SOME ADDITIVE ARITHMETICAL FUNCTIONS II.

J.-L. Mauclaire (Paris, France)

Dedicated to the memory of Béla Kovács

I. Introduction

Let G be an abelian group. A function f defined on the set of the positive integers \mathbb{N}^* is a G -valued additive arithmetical function if $f(mn) = f(m) + f(n)$ when $(m, n) = 1$. In 1946 P.Erdős [1] proved that if a real-valued additive arithmetical function f satisfies the condition $(f(n+1) - f(n)) \rightarrow 0, n \rightarrow +\infty$, then there exists a constant C such that the equality $f(n) = C \log n$ holds for all n in \mathbb{N}^* .

In his article [4] I.Z.Ruzsa has suggested to consider the problem of the distribution of group-valued additive arithmetical functions, and in this context I have extended the result of P.Erdős to the case of arithmetical additive functions with values in a locally compact abelian group: an additive arithmetical function with values in G satisfies the condition $(f(n+1) - f(n)) \rightarrow 0, n \rightarrow +\infty$, if and only if there exists a continuous homomorphism $\varphi : \mathbb{R} \rightarrow G$ such that for any n in \mathbb{N}^* , $f(n) = \varphi(\log n)$ [2]. And as proved by I.Z.Ruzsa and R.Tijdeman [5] this cannot be generalized to all groups.

Answering a question of P.Erdős asked for in the abovementioned article [1], E.Wirsing [6] provided a characterization of a real-valued additive arithmetical function satisfying the condition $(f(n+1) - f(n)) = O(1)$: a real-valued additive arithmetical function satisfies the condition $(f(n+1) - f(n)) = O(1)$ if and only if there exists a constant C such that the sequence $(f(n) - C \log n)$ is bounded.

In this article I shall consider the same question for arithmetical additive functions with values in a locally compact abelian group G , and shall provide a characterization of G -valued arithmetical additive functions satisfying the

condition: there exists a compact neighborhood V of zero such that for all n in \mathbb{N}^* $(f(n+1) - f(n))$ belongs to V .

II. The results

We have the following result:

Theorem. *Let G be a locally compact abelian group with group law denoted additively and f a G -valued arithmetical additive function. The following assertions are equivalent:*

- i) *there exists a compact neighborhood V of zero such that for all n in \mathbb{N}^* $(f(n+1) - f(n))$ belongs to V ;*
- ii) *there exists a continuous homomorphism $\varphi : \mathbb{R} \rightarrow G$ and a compact neighborhood of zero W such that for all n in \mathbb{N}^* $(f(n) - \varphi(\log n))$ belongs to W .*

Remark. To obtain the Theorem we shall use the following result.

Proposition. *If G is an abelian group and f is a G -valued additive arithmetical function such that the sequence $(f(n+1) - f(n))$ takes only a finite number of values, then the sequence $(f(n))$ takes only a finite number of values.*

N.B. This proposition is an answer to my naive question IV.2.1 in [3].

III. Proofs of the results

III.1. Proof of the proposition

If G is an abelian group and f is a G -valued additive arithmetical function such that the sequence $(f(n+1) - f(n))$ takes only a finite number of values, then clearly the sequence $(f(n))$ takes its values in a finitely generated \mathbb{Z} -module G' . Now as a finitely generated \mathbb{Z} -module G' is isomorphic to a product $\mathbb{Z}^r \times (\mathbb{Z}/n_1\mathbb{Z}) \times \dots \times (\mathbb{Z}/n_s\mathbb{Z})$, where r is a nonnegative integer, and the finite sequence (n_i) , $1 \leq i \leq s$, of positive integers greater than 2 is such that n_i divides n_{i-1} for $i = 2$ to s . To this isomorphism we can associate $(f)_{1 \leq j \leq r+s}$, a decomposition of the function f , where f_j is a \mathbb{Z} -valued additive arithmetical function for $1 \leq j \leq r$ and f_j is a $\mathbb{Z}/n_{j-r}\mathbb{Z}$ -valued additive arithmetical

function for $r + 1 \leq j \leq r + s$. Now, for $r + 1 \leq j \leq r + s$, $f_j(\mathbb{N}^*)$ is contained in the finite set $\mathbb{Z}/n_{j-r}\mathbb{Z}$, and so to obtain the proposition it will be sufficient to prove that a \mathbb{Z} -valued additive arithmetical function f such that the sequence $(f(n + 1) - f(n))$ takes only a finite number of values, takes also only a finite number of values. To do that we recall the following recent result of mine [3]:

Let f be a real-valued additive arithmetical function satisfying the condition (H): There exists a finite set Ω such that

$$\lim_{n \rightarrow +\infty} \min_{\omega \in \Omega} |f(n + 1) - f(n) - \omega| = 0.$$

Then there exists a constant C such that the sequence $f'(n) = f(n) - C \text{Log} n$ takes a finite number of values on \mathbb{N}^ .*

Since a \mathbb{Z} -valued additive arithmetical function can be viewed as a real-valued additive arithmetical function, this result gives us that there exists a constant C such that the sequence $f'(n) = f(n) - C \text{Log} n$ takes a finite number of values on \mathbb{N}^* . Since $f(n + 1) - f(n)$ takes a finite number of values on \mathbb{N}^* , by difference, we get that the number of values of $C(\text{Log}(n + 1) - \text{Log} n)$ is finite, too, which implies that the value of C is equal to zero, and so, that $f'(n) = f(n)$ and this gives that the number of the values of the sequences $f(n)$ is finite. This ends the proof of the proposition.

III.2. Proof of the theorem

III.2.1. Proof of $ii) \Rightarrow i)$

We assume that there exists a continuous homomorphism $\varphi : \mathbb{R} \rightarrow G$ and a compact neighborhood of zero W such that for all n in \mathbb{N}^* $(f(n) - \varphi(\log(n)))$ belongs to W . Then we have $((f(n + 1) - \varphi(\log(n + 1))) - (f(n) - \varphi(\log(n))))$ belongs to $W' = W - W$, which is still a compact neighborhood of zero. This gives that $((f(n + 1) - f(n)) - \varphi(\log(1 + 1/n)))$ is in W' , and since φ is a continuous group homomorphism, $\varphi(\log(1 + 1/n))$ tends to zero, and so there exists a compact neighborhood W'' of zero such that for all n in \mathbb{N}^* $\varphi(\log(1 + 1/n))$ is in W'' . Hence we get that for all n in \mathbb{N}^* $(f(n + 1) - f(n))$ belongs to the compact neighborhood of zero V defined by $V = W' - W''$.

III.2.2. Proof of $i) \Rightarrow ii)$

We assume that there exists a compact neighborhood V of zero such that for all n in \mathbb{N}^* $(f(n + 1) - f(n))$ belongs to V and shall prove that there exists a continuous homomorphism $\varphi : \mathbb{R} \rightarrow G$ and a compact neighborhood of zero W such that for all n in \mathbb{N}^* $(f(n) - \varphi(\log(n)))$ belongs to W .

Since G is a locally compact abelian group, the structure theorem gives that G can be written as $\mathbb{R}^m \times G'$, where G' contains an open compact subgroup H . Now, as above in the proof of the proposition, we associate to f a decomposition $f = (f_1, \dots, f_m, g)$, where f_j , $1 \leq j \leq m$, are real valued additive functions and g is a G' -valued additive function. Since there exists a compact neighborhood V of zero in G such that for all n in \mathbb{N}^* $(f(n+1) - f(n))$ belongs to V , we get that there exist m compact neighborhoods V_j of zero in \mathbb{R} , $1 \leq j \leq m$, and W , a compact neighborhood of zero in G' such that for all n in \mathbb{N}^* $(f_j(n+1) - f_j(n))$ belongs to V_j and $(g(n+1) - g(n))$ belongs to W . Now, by the original result of Wirsing, we get that for all j , $1 \leq j \leq m$, there exists a real number C_j such that the sequence $(f_j(n) - C_j \log n)$ is bounded.

It is clear that to end the proof of the theorem it will be sufficient to prove it for the special case of a function f taking its values in a group G' , where G' contains an open compact subgroup H .

Let K be the subgroup of G' generated by W . K is open and closed in G' and H' , the intersection of H and K is also open and closed since H is so, and compact as a closed subgroup of H . As a consequence, the quotient group K/H' is discrete. Let T be the canonical homomorphism $K \rightarrow K/H'$. The sequence $T(f(n))$ is a K/H' -valued additive arithmetical function, and for all n $T(f(n+1)) - T(f(n))$ belongs to $T(W)$. But T is continuous and so $T(W)$ is compact, hence finite since K/H' is discrete. This gives us that $T(f(n+1)) - T(f(n))$ takes only a finite number of values, and by the Proposition the sequence $T(f(n))$ takes also a finite number of values, say $\overline{\alpha}_u$, $u \in U$, a finite set. This implies that $f(n)$ belongs to the union of a finite number of cosets $\alpha_u + H'$, where α_u is in $\overline{\alpha}_u$. But $f(1)$ is equal to zero. This implies that one of the α_u is equal to zero. Now, since each of the cosets $\alpha_u + H'$ is compact, their union W is also compact and is a compact neighborhood of zero since it contains H' . Hence we have proved that for all n $f(n)$ is in a compact neighborhood of zero, and this ends the proof of the theorem.

References

- [1] Erdős P., On the distribution function of additive functions, *Annals of Math.*, **47** (1946), 1-20.
- [2] Mauclaire J.-L., On the regularity of group valued additive arithmetical functions, *Publ. Math.*, **44** (1994), 285-290.
- [3] Mauclaire J.-L., A characterization of some additive arithmetical functions (to appear)

- [4] **Ruzsa I.Z.**, General multiplicative functions, *Acta Arith.*, **32** (1977), 313-347.
- [5] **Ruzsa I.Z. and Tijdeman R.**, On the difference of integer-valued additive functions, *Publ. Math. Debrecen*, **39** (1991), 353-358.
- [6] **Wirsing E.**, A characterization of the logarithm as an additive arithmetic function, *Symposia Mathematica*, **4** (1970), 45-57.

J.-L. Mauclaire

U.M.R. 9994 du C.N.R.S., Institut Mathématique de Jussieu

Université Paris VII

2, Place Jussieu

75251 Paris Cedex 05, France