

**A NOTE ON THE PRODUCT
OF CONSECUTIVE ELEMENTS
OF AN ARITHMETIC PROGRESSION**

Yuan Jin (Xi'an, China)

1. Introduction

For an integer $x > 1$ we denote by $P(x)$ the greatest prime factor of x and by $\pi(x)$ the number of primes $\leq x$. We consider the equation

$$(1.1) \quad (n + d)(n + 2d) \dots (n + kd) = y^\ell$$

in positive integers d, k, ℓ, n, y subject to $\gcd(n, d) = 1, k > 2, \ell \geq 2$.

P.Erdős and J.L.Selfridge confirm in [1] an old conjecture that equation (1.1) has no solution if $d = 1$. Furthermore, Erdős conjectured that equation (1.1) implies that k is bounded by an absolute constant.

R.Marszalek [2] considered equation (1.1) with $d \geq 2$. He showed that k is bounded if d is fixed. More precisely, he proved that for any solution of (1.1) with $d \geq 2$ we have

$$\begin{aligned} k &< 2 \exp[d(d + 1)^{1/2}] && \text{if } \ell = 2, \\ k &< \max\{30000, (3/2) \exp[1/2d(d + 2)(d + 1)^{1/3}]\} && \text{if } \ell = 3, \\ k &< \max\{30000, (1/4)d(d + 2)(d + 1)^{1/2}\} && \text{if } \ell = 4, \\ k &< \max\{30000, (3/2)(d + 1)\} && \text{if } \ell \geq 5. \end{aligned}$$

The results in this paper considerably improve the results of Marszalek. We will prove the following result

Theorem. *For every integer $d \geq 2$ and $\ell \geq 2$ there exists a constant $k_0(d, \ell)$ such that for $k \geq k_0(d, \ell)$ the equation (1.1) has no solution. For $k_0(d, \ell)$ we can take the following values:*

$$\begin{aligned} k_0(d, 2) &= \max[64, 2 \exp(d)], \\ k_0(d, 3) &= \max[30000, (3/2) \exp(d^{1/3})], \\ k_0(d, \ell) &= \max[30000, d] \quad \text{for } \ell \geq 4. \end{aligned}$$

2. Lemmas

For the proof we need the following results.

Lemma 1. (T.N.Shorey and R.Tijdeman [3]) *If $d > 1$ and $(n+d, d, k) \neq (2, 7, 3)$, then $P(\Delta) > k$, where $\Delta = (n+d)(n+2d)\dots(n+kd)$.*

Lemma 2. (R.Marszalek [2]) *Let d be a positive integer and let f be a real function for which there exists a positive integer k_0 , such that f is positive and nondecreasing on the interval $[k_0, \infty)$. If the positive integers n and k satisfy*

$$\begin{aligned} \gcd(n, d) &= 1, & n + d &> kf(k), \\ k &> \max\{k_0, 2\pi[1 + d/f(k_0)]\}, \end{aligned}$$

then

$$\pi[P(\Delta)] > k\{\log[f(k) + d]/[\log(f(k) + d) + \log k]\}.$$

Lemma 3. *The equation (1.1) with $d \geq 2$ has no solution if $k \geq \max(d, n)$.*

Proof. If the equation (1.1) has solution, by Lemma 1 there exists a prime $P > k$ dividing exactly one factor of Δ . Thus

$$(2.1) \quad n + kd \geq (k+1)^\ell \geq (k+1)^2.$$

On the other hand, if $k \geq \max(n, d)$ we have

$$(2.2) \quad n + kd \leq k + k^2 < (k+1)^2.$$

However (2.1) contradicts to (2.2). This completes the proof of Lemma 3.

Lemma 3 implies that we may confine ourselves to the case

$$(2.3) \quad d \leq k < n$$

to complete the proof of our theorem.

We assume that d, k, n, ℓ and y are positive integers satisfying the equation (1.1). Thus, for $1 \leq i \leq k$ we can write

$$(2.4) \quad n + id = a_i x_i^\ell,$$

where a_i is ℓ -th power-free and its prime factors are less than k .

Lemma 4. *The products $a_i a_j$ are all distinct provided*

- (1) $k \geq d$ for $\ell > 3$,
- (2) $k \geq (3/2) \exp(d^{4/3})$ for $\ell = 3$.

Proof. By Lemma 1 and (2.3) we have

$$(k + 1)^\ell \leq n + kd \leq n + k^2.$$

Therefore

$$(2.5) \quad k^\ell < n \quad \text{if } \ell \geq 3.$$

For $1 \leq i, j, r, s \leq k$ and $\langle i, j \rangle \neq \langle r, s \rangle$ we have $\gcd(n + id, n + rd) < k$, $\gcd(n + id, n + sd) < k$ and by (2.5) $n + id > k^2$. If $n + id$ divides $(n + rd)(n + sd)$, then $\gcd[n + id, (n + rd)(n + sd)] = n + id > k^2$. However, this is not possible. So, it follows that $n + id$ cannot divide $(n + rd)(n + sd)$. Hence the products $(n + id)(n + jd)$ and $(n + rd)(n + sd)$ are distinct.

Suppose that for some $1 \leq i, j, r, s \leq k$ and $\langle i, j \rangle \neq \langle r, s \rangle$ one has $a_i a_j = a_r a_s$. Putting $T = (n + id)(n + jd) - (n + rd)(n + sd)$ (which we may assume to be positive) and $A = a_i a_j$, we get

$$\begin{aligned} (n + id)(n + jd) &= a_i a_j x^\ell = Ax^\ell, \\ (n + rd)(n + sd) &= a_r a_s y^\ell = Ay^\ell. \end{aligned}$$

Hence $Ax^\ell > Ay^\ell$, and therefore $x \geq y + 1$. Thus $T \geq A[(y + 1)^\ell - y^\ell] > Aly^{\ell-1}$. Since $Ay^\ell \geq (n + d)^2$ and A is an integer, so we obtain

$$(2.6) \quad T > \ell(n + d)^{2(\ell-1)/\ell}.$$

On the other hand

$$T \leq (n + kd)^2 - (n + d)^2 = 2kdn + k^2d^2 - 2nd - d^2.$$

Using (2.5) we get

$$2nd > 2k^\ell d \geq 2k^3 d > k^2 d^2.$$

So

$$(2.7) \quad T < 2kdn.$$

By (2.6) and (2.7) it follows

$$\ell(n+d)^{2(\ell-1)/\ell} < 2kdn < 2kd(n+d).$$

Then

$$(2.8) \quad \ell^\ell(n+d)^{\ell-2} < 2^\ell k^\ell d^\ell.$$

Now we have to consider separately the cases $\ell > 3$ and $\ell = 3$. If $\ell > 3$ and $k \geq d$, then

$$3^\ell(n+d)^2 \leq \ell^\ell(n+d)^{\ell-2} < 2^\ell k^\ell d^\ell \leq 2^\ell k^{2\ell}.$$

However, this contradicts to (2.5).

In the case $\ell = 3$ by (2.5) we see that $n+d > k(k^2-d)$. This enables us to utilize Lemma 2 for $f(k) = k^2 - d$. Therefore there exists a prime P dividing Δ such that $\pi(P) > 2/3k$. By $x > \pi(x) \log \pi(x)$, this gives

$$P > (2/3)k \log(2k/3) \geq (2/3)kd^{4/3} \quad \text{for } k \text{ satisfying (2).}$$

From (2.4) and the fact that P divides only one factor of Δ , we get

$$(2.9) \quad n + kd > [(2/3)kd^{4/3}]^3.$$

Since

$$n + kd = n + d + (k-1)d,$$

then from (2.8) and (2.9)

$$[(2/3)kd^{4/3}]^3 < n + d + k^2 < [(2/3)kd]^3 + k^2.$$

This implies $d < 2$, and Lemma 4 is proved.

Let G be the set of primes p dividing Δ with $p \leq k-1$. For every $p \in G$ we choose a $u(p) \in \{1, 2, \dots, k\}$ such that

$$(2.10) \quad \text{ord}_p[n + u(p)d] = \max\{\text{ord}_p(n + jd)\},$$

where $1 \leq j \leq k$. We denote by H the set of all elements from $\{1, 2, \dots, k\}$ which do not appear in the range of u . Then we have

Lemma 5.

$$(2.11) \quad \prod_{j \in H} a_j \mid (k-1)!$$

Proof. For each prime $p \in G$, if $1 \leq j \leq k$ and $j \neq u(p)$, we have

$$(2.12) \quad \text{ord}_p(n + jd) \leq \text{ord}_p[u(p) - j],$$

since if $p^m \mid n + jd$, then (2.10) and $\text{gcd}(n, d) = 1$ imply $p^m \mid u(p) - j$. Hence

$$\begin{aligned} \text{ord}_p \left[\prod_{1 \leq j \leq k, j \neq u(p)} (n + jd) \right] &\leq \text{ord}_p \left[\prod_{1 \leq j \leq k, j \neq u(p)} (u(p) - j) \right] = \\ &= \text{ord}_p[(u(p) - 1)!(k - u(p)!)] \leq \text{ord}_p[(k - 1)!]. \end{aligned}$$

Thus, (2.11) follows from

$$\text{ord}_p \left(\prod_{j \in H} a_j \right) \leq \text{ord}_p \left[\prod_{j \in H} (n + jd) \right] \leq \text{ord}_p \left[\prod_{1 \leq j \leq k, j \neq u(p)} (n + jd) \right].$$

Note that

$$(2.13) \quad |H| \geq k - \pi(k - 1),$$

where $|A|$ denotes the cardinality of set A .

Lemma 6. (P.Erdős and J.L.Selfridge [1]) *Let $b_1 < b_2 < \dots < b_k$ be positive integers such that the products $b_i b_j$ are all distinct. Then for $k \geq 30000$*

$$(2.14) \quad \prod_{i \in D} b_i > k!,$$

where D is any subset of $\{1, 2, \dots, k\}$ satisfying $|D| \geq k - \pi(k)$.

Lemma 7. *If $k \geq 2 \exp(q)$ and $q \geq 5$, then*

$$(2.15) \quad 3^{(k-6)/4} q^{(k-1)/(q-1)} > 2^{(k+6)/3} k^4,$$

where k and q are positive integers.

Proof. First we prove that

$$(2.16) \quad 3^{(k-6)/4} > 2^{(k+6)/3}, \quad \text{if } k \geq 2 \exp(5).$$

If (2.16) is false, then

$$3^{(k-6)/4} \leq 2^{(k+6)/3}.$$

So

$$4(k+6)\log 2 \geq 3(k-6)\log 3.$$

This implies

$$(2.17) \quad k(3\log 3 - 4\log 2) \leq 24\log 2 + 18\log 3.$$

However it is impossible for $k \geq 2\exp(5)$. Thus we have (2.16).

Next we prove that if $k \geq 2\exp(q)$ and $q \geq 5$, then

$$(2.18) \quad q^{(k-1)/(q-1)} > k^4.$$

If $k \geq 2\exp(q)$ and $q \geq 5$, then

$$q^{(k-1)^{1/2}} > k.$$

Thus

$$(k-1)^{1/2} > (\log k)/(\log q).$$

Since

$$(k-1)^{1/2} \leq (k-1)/4(q-1),$$

we have

$$(\log k)/(\log q) < (k-1)/4(q-1).$$

Consequently, (2.18) is true.

3. Proof of the Theorem

a) The case $\ell \geq 3$. Lemma 4 enables us to apply Lemma 6 to the set H given by Lemma 5. Thus in the case $\ell \geq 3$, since (2.11) and (2.14) are in contradiction for k satisfying (1), (2) and $k \geq 30000$, we have proved: if

$$\begin{aligned} k &\geq \max\{30000, 3/2\exp(d^{4/3})\} && \text{for } \ell = 3, \\ k &\geq \max\{30000, d\} && \text{for } \ell > 3, \end{aligned}$$

then the equation (1.1) has no solution.

b) The case $\ell = 2$. Now suppose that the theorem is false for $\ell = 2$. We shall first prove that if $k \geq 2\exp(d)$ and $i \neq j$, then $a_i \neq a_j$. Suppose that $a_i = a_j$ for some $i \neq j$. Assuming that $x_i \geq x_j + 1$, we have

$$d(k-1) = (n+kd) - (n+d) \geq (n+id) - (n+jd) = a_j(x_i^2 - x_j^2) > 2x_j a_j \geq$$

$$\geq 2(n + d)^{1/2}.$$

Hence

$$(3.1) \quad (n + d) < [d^2(k - 1)^2]/4.$$

On the other hand, by Lemma 1, we have $n + d > k(k - d)$. Thus we may utilize Lemma 2 for $f(k) = k - d$. Therefore there exists a prime P dividing Δ , such that $\pi(P) > 1/2k$, which by $x > \pi(x) \log \pi(x)$ gives $P > (kd)/2$ for $k \geq 2 \exp(d)$.

Since P divides only one factor of Δ which is a square, we get $n + kd > > P^2 > (k^2d^2)/4$. Thus

$$(3.2) \quad (n + d) > [(k^2d^2)/4] - (k - 1)d.$$

By (3.1) and (3.2) we have

$$(3.3) \quad [k^2d^2/4] - (k - 1)d < [d^2(k - 1)^2]/4.$$

Thus (3.3) gives

$$2k(d - 2) < d - 4.$$

However, this is not possible for $d \geq 2$. Thus for $k \geq 2 \exp(d)$ the a 's are distinct and square-free. So by Lemma 5

$$(3.4) \quad \prod_{1 \leq j \leq k} a_j \mid (k - 1)! \prod_{p < k} p.$$

Let us for a prime q put $g_q = \text{ord}_q \left(\prod_{1 \leq j \leq k} a_j \right)$ and $h_q = \text{ord}_q[(k - 1)!]$. Then by (3.4) if $g_2 \geq h_2$, then we have

$$\prod_{1 \leq j \leq k} a_j \mid (k - 1)! 2^{g_2 - h_2} \prod_{p < k} p,$$

and if $g_2 < h_2$, then there exists an integer w which satisfies $\text{ord}_2(w) > h_2 - g_2$ and

$$w \prod_{1 \leq j \leq k} a_j = (k - 1)! \prod_{p < k} p.$$

So we get

$$(3.5) \quad \prod_{1 \leq j \leq k} a_j \mid (k - 1)! 2^{g_2 - h_2} \prod_{p < k} p.$$

Similarly, we have

$$(3.6) \quad \prod_{1 \leq j \leq k} a_j \mid (k-1)! 2^{g_2 - h_2} 3^{g_3 - h_3} \prod_{p < k} p.$$

If 2 cannot divide d and 3 also cannot divide d , then there is a prime $q \geq 5$ such that $q \mid d$. Therefore q cannot divide a_i . Thus

$$(3.7) \quad \prod_{1 \leq j \leq k} a_j \mid (k-1)! 2^{g_2 - h_2} 3^{g_3 - h_3} q^{-h_q} \prod_{p < k} p.$$

On the other hand, for a prime q we have

$$g_q \leq [k/(q+1)] + \log_q k + 1 \quad (\text{cf. [2] p.221})$$

and also

$$h_q \geq [(k-1)/(q-1)] - \log_q k \quad (\text{cf. [2] p.221}).$$

Therefore

$$(3.8) \quad g_2 - h_2 \leq -(2/3)k + 2 \log_2 k + 2, \quad g_3 - h_3 \leq -(1/4)k + 2 \log_3 k + (3/2).$$

Further, using the above inequality,

$$(3.9) \quad \prod_{p < k} p < 3^k, \quad \text{for } k = 1, 2, \dots$$

(see for example [4]) and the fact that the product of k consecutive square-free integers is greater than $k!(3/2)^k$ for $k \geq 64$ (see [1]), we obtain

$$(3.10) \quad 3^{(k-6)/4} q^{(k-1)/(q-1)} < 2^{(k+6)/3} k^4,$$

which is in contradiction with Lemma 7. If $2 \mid d$ or $3 \mid d$, then by (3.6) and (3.8) we get

$$3^{(k-6)/4} < 2k^2 \quad \text{or} \quad 3^{(k-1)/2} < 2^{(k+6)/3} k^2$$

which also give a contradiction for $k \geq \max[64, 2 \exp(d)]$. If $2 \mid d$ and $3 \mid d$, then we get

$$3^{(k-1)/2} < 2k.$$

This leads also to a contradiction. So we complete the proof of the theorem in the case $\ell = 2$.

References

- [1] **Erdős P. and Selfridge J.L.**, The product of consecutive integers is never a power, *Illinois J. Math.*, **19** (1975), 292-301.
- [2] **Marszalek R.**, On the product of consecutive elements of an arithmetic progression, *Monatsh. Math.*, **100** (1985), 215-222.
- [3] **Shorey T.N. and Tijdeman R.**, On the greatest prime factors of an arithmetical progression III., *Diophantine approximations and transcendental numbers*, de Gruyter, 1992.
- [4] **Hanson D.**, On a theorem of Sylvester and Schur, *Canad. Math. Bull.*, **16** (1973), 195-199.

(Received November 29, 1994)

Yuan Jin
Department of Mathematics
Northwest University
Xi'an, China