

## PERMUTATIONS AVOIDING CONSECUTIVE PATTERNS

**R. Warlimont** (Johannesburg, South Africa)

*Karl-Heinz Indlekofer zum 60. Geburtstag*

**Abstract.** Given some permutation  $\sigma \in S_m$  ( $m \geq 3$ ) denote by  $B(n, m; \sigma)$  the set of all permutations  $\pi \in S_n$  with the property that

$$\pi(k + \sigma(1)) < \dots < \pi(k + \sigma(m)) \text{ is false for } 0 \leq k \leq n - m.$$

We conjecture that to any  $\sigma \in S_m$  there are constants  $c(\sigma) > 0$ ,  $\xi(\sigma) > 1$  such that

$$|B(n, m; \sigma)|/n! \sim c(\sigma)\xi(\sigma)^{-n} \text{ as } n \rightarrow \infty.$$

We prove this for  $m = 3$  and for the particular  $\sigma \in S_m$  being the identity:  $\sigma(j) = j$  ( $1 \leq j \leq m$ ).

### 1. Introduction

Let  $S_n$  denote the set of all permutations  $\pi$  of  $1, \dots, n$ . Let  $m \geq 3$  and some permutation  $\sigma \in S_m$  be fixed,  $\epsilon$  the identity ( $\epsilon(j) = j$  for  $1 \leq j \leq m$ ).

1. Let  $A(n, m; \sigma)$  denote the set of all  $\pi \in S_n$  with the property that

$$\pi(j_{\sigma(1)}) < \dots < \pi(j_{\sigma(m)}) \text{ is false for all } 1 \leq j_1 < \dots < j_m \leq n.$$

In particular  $A(n, m; \epsilon)$  is the set of all those  $\pi \in S_n$  with the property that

$$\pi(j_1) < \dots < \pi(j_m) \text{ is false for all } 1 \leq j_1 < \dots < j_m \leq n.$$

The Stanley-Wilf conjecture states that to any  $m \geq 3$  there is some constant  $K(m) > 1$  such that

$$|A(n, m; \sigma)| \leq K(m)^n \quad \text{for all } \sigma \in S_m.$$

The following facts are known:

$$\begin{aligned} |A(n, 3; \sigma)| &= \frac{1}{n+1} \binom{2n}{n} \quad \text{for all } \sigma \in S_3, \\ (*) \quad |A(n, m; \epsilon)| &\leq (m-1)^{2n}, \\ |A(n, m; \epsilon)| &\sim c(m) \frac{(m-1)^{2n}}{n^{m(m-2)/2}} \quad \text{as } n \rightarrow \infty. \end{aligned}$$

$|A(n, m; \sigma)|$  varies considerably with  $\sigma$  already for  $m = 4$ .

References for this can be found in Richard Arratia's paper "On the Stanley-Wilf conjecture for the number of permutations avoiding a given pattern", *Electronic J. Combin.* 6 (1999), #N1. The right guess for  $K(m)$  should be  $(m-1)^2$ .

We mention that we could sharpen (\*):

$$|A(n, m; \epsilon)| \leq \frac{(m-1)^{2n}}{\prod_{k=1}^{m-2} \left(1 + \frac{1}{k}\right)^k}.$$

2. Let  $B(n, m; \sigma)$  denote the set of all  $\pi \in S_n$  with the property that

$$\pi(k + \sigma(1)) < \dots < \pi(k + \sigma(m)) \quad \text{is false for } 0 \leq k \leq n - m.$$

In particular  $B(n, m; \epsilon)$  is the set of all  $\pi \in S_n$  with the property that

$$\pi(k) < \dots < \pi(k + m - 1) \quad \text{is false for } 1 \leq k \leq n - m + 1.$$

Since the conditions imposed on  $\pi$  there concern only  $m$  consecutive values of the argument of  $\pi$  they are less restrictive than those met before; hence  $A(n, m; \sigma) \subset B(n, m; \sigma)$ . Our results for  $B$  are at this stage as incomplete as those for  $A$ . Put

$$\begin{aligned} b(n, m; \sigma) &:= \frac{|B(n, m; \sigma)|}{n!}, \quad b(0, m; \sigma) := 1, \\ b(n, m) &:= b(n, m; \epsilon), \quad b(n) := b(n, 3), \\ \tilde{b}(n) &:= b(n, 3; \sigma_1), \quad \text{where } \sigma_1 := \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}. \end{aligned}$$

We obtained:

- (1)  $b(n, m; \sigma) \leq \left(1 - \frac{1}{m!}\right)^{\lfloor \frac{n}{m} \rfloor}$  for all  $\sigma \in S_m$ ;
- (2)  $\lim_{n \rightarrow \infty} b(n, m; \sigma)^{\frac{1}{n}}$  exists and is equal to  $\inf_n b(n, m; \sigma)^{\frac{1}{n}}$  for all  $\sigma \in S_m$ ;
- (3)  $\sum_{n=0}^{\infty} b(n, m)z^n = P_m(z)^{-1}$ , where

$$P_m(z) := \sum_{\substack{k=0 \\ k \equiv 0(m)}}^{\infty} \frac{z^k}{k!} - \sum_{\substack{k=0 \\ k \equiv 1(m)}}^{\infty} \frac{z^k}{k!};$$

- (4)  $b(n, m) \sim c(m)\xi(m)^{-n}$  as  $n \rightarrow \infty$ , where

$$c(m) > 0 \text{ and } \xi(m) = 1 + \frac{1}{m!}(1 + o(1)) \text{ as } m \rightarrow \infty;$$

in particular

- (5)  $b(n) \sim \exp\left(\frac{\pi}{\sqrt{3}}\right) \left(\frac{3\sqrt{3}}{2\pi}\right)^{n+1}$  as  $n \rightarrow \infty$ ;
- (6)  $\sum_{n=0}^{\infty} \tilde{b}(n)z^n = \tilde{P}(z)^{-1}$ , where

$$\tilde{P}(z) := 1 - \int_0^z \exp\left(-\frac{1}{2}w^2\right) dw;$$

- (7)  $\tilde{b}(n) \sim \frac{\exp\left(\frac{1}{2}\xi^2\right)}{\xi^{n+1}}$  as  $n \rightarrow \infty$ , where  $\xi$  is given by

$$\int_0^{\xi} \exp\left(-\frac{1}{2}t^2\right) dt = 1 \quad (1 < \xi < \infty);$$

- (8)  $b(n, 3; \sigma) = b(n)$  for  $\sigma = \epsilon, \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$ ,

$b(n, 3; \sigma) = \tilde{b}(n)$  for the 4 remaining  $\sigma \in S_3$ .

We conjecture that to any  $\sigma \in S_m$  ( $m \geq 3$ ) there are constants  $c(\sigma) > 0$ ,  $\xi(\sigma) > 1$  such that

$$b(n, m; \sigma) \sim c(\sigma)\xi(\sigma)^{-n} \text{ as } n \rightarrow \infty.$$

In the last section of our paper we report on  $m = 4$ , where we could achieve only partial results.

**3. Acknowledgement.** The author thanks Arnold Knopfmacher and Helmut Prodinger for bringing to his attention the PhD problem of their student Albert Tshifhumulo on permutations avoiding consecutive patterns.

Tshifhumulo in his thesis will study these problems mainly using an approach based on geometrically distributed random variables. (3) of our list of results will appear in the thesis. Here I will study Tshifhumulo's problem from a different (elementary) perspective based on a direct treatment of permutations.

Finally I am obliged to Arnold Knopfmacher for localizing and bringing to my attention Arratia's paper.

## 2. The proofs of (1.1), (1.2)

Since  $B(m, m; \sigma) = S_m - \{\sigma^{-1}\}$ , we get

$$(1) \quad b(m, m; \sigma) = 1 - \frac{1}{m!}.$$

Put  $B(k) := |B(k, m; \sigma)|$  and  $b(k) := \frac{B(k)}{k!}$ ,  $b(0) = 1$ . Then we have

$$(2) \quad b(n) \leq b(p)b(n-p) \quad \text{for } 0 \leq p \leq n.$$

**Proof.** This is true for  $p = 0, n$ . Assume  $0 < p < n$ . From

$$B(n, m; \sigma) = \bigcup_{\substack{P \subseteq [1, n] \\ |P|=p}} \{\pi \in B(n, m; \sigma) \mid \pi([1, p]) = P\}$$

and

$$\#\{\pi \in B(n, m; \sigma) \mid \pi([1, p]) = P\} \leq B(p)B(n-p)$$

we obtain

$$B(n) \leq \binom{n}{p} B(p)B(n-p)$$

which gives (2).

From (2), (1) we infer

$$b(n) \leq b(m)^{\lfloor \frac{n}{m} \rfloor} b\left(n - \left\lfloor \frac{n}{m} \right\rfloor m\right) \leq b(m)^{\lfloor \frac{n}{m} \rfloor} = \left(1 - \frac{1}{m!}\right)^{\lfloor \frac{n}{m} \rfloor}$$

which is (1.1).

From (2) we get

$$b(p + q) \leq b(p)b(q) \text{ for all } p, q \geq 0,$$

which yields (1.2).

### 3. A formula for $b(n, m)$ and the proof of (1.3)

We shall need two formulae whose proofs are left to the reader.

$$(1) \quad \sum_{\substack{\delta_s \geq 0 \\ \sum_{s=1}^t \delta_s \leq N}} \binom{N - \sum_{s=1}^t \delta_s}{1 \leq s \leq t} = \binom{t + N}{t + 1}.$$

Put

$$S(h, q, b) := \sum_{\substack{0 \leq \delta_i \leq q \\ \sum_{i=1}^h \delta_i = b}} 1.$$

Put  $m = q + 2$ . Then

$$(2) \quad \begin{aligned} f_m(a) &:= \sum_{h=0}^a (-1)^{h+1} S(h, q, a - h) = \\ &= \begin{cases} -1 & \text{if } a \equiv 0(m), \\ 1 & \text{if } a \equiv 1(m), \\ 0 & \text{else.} \end{cases} \end{aligned}$$

Put  $p = m - 1$ . The sieve formula yields

$$b(n, m) = 1 + \sum_{j=1}^{n-p} (-1)^j \sum_{1 \leq k_1 < \dots < k_j \leq n-p} \frac{D(k_1, \dots, k_j)}{n!},$$

where

$$D(k_1, \dots, k_j) = \# \{ \pi \in S_n \mid \pi(k_i) < \dots < \pi(k_i + p) \text{ for } 1 \leq i \leq j \}.$$

We find

$$D(k_1, \dots, k_j) = \frac{n!}{M(k_1, \dots, k_j)},$$

where  $M(k_1, \dots, k_j)$  is defined in the following way:

$$\begin{aligned} j = 1: & \quad M(k) = m! \\ j \geq 2: & \quad \text{Put} \end{aligned}$$

$$L(k_1, \dots, k_j) := \{ i \mid 1 \leq i \leq j, k_{i+1} - k_i \geq m \}.$$

If  $L(k_1, \dots, k_j) = \emptyset$  then

$$M(k_1, \dots, k_j) = (k_j - k_1 + m)!$$

If  $L(k_1, \dots, k_j) \neq \emptyset$  write

$$L(k_1, \dots, k_j) = \{ \ell_1, \dots, \ell_t \} \quad \text{with } 1 \leq \ell_1 < \dots < \ell_t < j.$$

Furthermore, put  $\ell_0 = 0$ ,  $\ell_{t+1} = j$ . Then

$$M(k_1, \dots, k_j) = \prod_{s=1}^{t+1} (k_{\ell_s} - k_{\ell_{s-1}+1} + m)!$$

This, too, should be verified by the reader.

We have

$$mt \leq \sum_{i=1}^{j-1} (k_{i+1} - k_i) = k_j - k_1 \leq n - m.$$

Therefore  $0 \leq t \leq T = \left\lceil \frac{n}{m} \right\rceil - 1$ . We get

$$b(n, m) = 1 + \sum_{t=0}^T X(t),$$

where

$$X(0) = -\frac{n-p}{m!} + \sum_{j=2}^{n-p} (-1)^j \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n-p \\ L(k_1, \dots, k_j) = 0}} M(k_1, \dots, k_j)^{-1}$$

and ( $t \geq 1$ )

$$X(t) = \sum_{j=t+1}^{n-p} (-1)^j \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n-p \\ |L(k_1, \dots, k_j)| = t}} M(k_1, \dots, k_j)^{-1}.$$

We have to evaluate ( $j \geq 2, t \geq 0$ ) the inner sums

$$X_j(t) := \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n-p \\ |L(k_1, \dots, k_j)| = t}} M(k_1, \dots, k_j)^{-1}.$$

We begin with  $t = 0$ .

$$\begin{aligned} X_j(0) &= \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n-p \\ k_{i+1} - k_i \leq p \ (1 \leq i < j)}} \frac{1}{(k_j - k_1 + m)!} = \\ &= \sum_{j-1 \leq d \leq n-m} \frac{1}{(d+m)!} \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n-p \\ k_{i+1} - k_i \leq p \ (1 \leq i < j) \\ k_j - k_1 = d}} 1, \\ X_j(0) &= \sum_{j-1 \leq d \leq n-m} \frac{n-p-d}{(d+m)!} S(j-1, q, d-(j-1)). \end{aligned}$$

Now let  $t \geq 1$ .

$$X_j(t) = \sum_{1 \leq \ell_1 < \dots < \ell_t < j} \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n-p \\ L(k_1, \dots, k_j) = \{\ell_1, \dots, \ell_t\}}} M(k_1, \dots, k_j)^{-1}.$$

Denote the inner sum by  $S_j(\ell_1, \dots, \ell_t)$  and write  $L(t) := \{\ell_1, \dots, \ell_t\}$ . Then

$$S_j(\ell_1, \dots, \ell_t) = \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n-p \\ L(k_1, \dots, k_j) = L(t)}} \left( \sum_{s=1}^{t+1} (k_{\ell_s} - k_{\ell_{s-1}+1} + m)! \right)^{-1}.$$

We introduce the differences  $d_i = k_{i+1} - k_i$  ( $1 \leq i \leq j$ ). They satisfy

$$d_i \leq p \quad (i \notin L(t)) \quad \text{and} \quad d_{\ell_s} \geq m \quad (1 \leq s \leq t).$$

We have

$$k_{\ell_s} - k_{\ell_{s-1}+1} = \sum_{i=\ell_{s-1}+1}^{\ell_s-1} d_i$$

and

$$\sum_{\substack{i=1 \\ i \notin L(t)}}^{j-1} d_i = k_j - k_1 - \sum_{\ell \in L(t)} d_\ell \leq n - m(t+1).$$

Therefore

$$S_j(\ell_1, \dots, \ell_t) = \sum_{\substack{1 \leq d_i \leq p \quad (i \notin L(t)) \\ D := \sum_{i \notin L(t)} d_i \leq n - m(t+1)}} \left( \prod_{s=1}^{t+1} \left( \sum_{i=\ell_{s-1}+1}^{\ell_s-1} d_i + m \right)! \right)^{-1} \cdot X,$$

where

$$\begin{aligned} X &= \sum_{\substack{d_{\ell_s} \geq m \quad (1 \leq s \leq t) \\ \sum_{s=1}^t d_{\ell_s} \leq n - m - D}} \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n - p \\ k_{i+1} - k_i = d_i \quad (1 \leq i < j)}} 1 = \\ &= \sum_{\substack{d_{\ell_s} \geq m \quad (1 \leq s \leq t) \\ \sum_{s=1}^t d_{\ell_s} \leq n - m - D}} \binom{n - p - D - \sum_{s=1}^t d_{\ell_s}}{t+1} = \\ &= \binom{n - p(t+1) - D}{t+1} \quad \text{by (1)}. \end{aligned}$$

Therefore

$$\begin{aligned} S_j(\ell_1, \dots, \ell_t) &= \sum_{\substack{1 \leq d_i \leq p \quad (i \notin L(t)) \\ \sum_{i \notin L(t)} d_i \leq n - m(t+1)}} \binom{n - p(t+1) - \sum_{i \notin L(t)} d_i}{t+1} \times \\ &\quad \times \left( \prod_{s=1}^{t+1} \left( \sum_{i=\ell_{s-1}+1}^{\ell_s-1} d_i + m \right)! \right)^{-1} = \\ &= \sum_{\substack{a_1 + \dots + a_{t+1} \leq n - m(t+1) \\ a_s \geq \ell_s - \ell_{s-1} - 1 \quad (1 \leq s \leq t+1)}} \binom{n - p(t+1) - \sum_{s=1}^{t+1} a_s}{t+1} \left( \prod_{s=1}^{t+1} (a_s + m)! \right)! \cdot Y, \end{aligned}$$



where

$$\begin{aligned}
 Y &= \sum_{\substack{1 \leq d_i \leq p \ (i \notin L(t)) \\ \sum_{i=\ell_{s-1}+1}^{\ell_s-1} d_i = a_s \ (1 \leq s \leq t+1)}} 1 = \prod_{s=1}^{t+1} \sum_{\substack{1 \leq d_i \leq p \ (\ell_{s-1} < i < \ell_s) \\ \sum_{i=\ell_{s-1}+1}^{\ell_s-1} d_i = a_s}} 1 = \\
 &= \prod_{s=1}^{t+1} S(\ell_s - \ell_{s-1} - 1, q, a_s - (\ell_s - \ell_{s-1} - 1)).
 \end{aligned}$$

Therefore we obtain

$$\begin{aligned}
 &S_j(\ell_1, \dots, \ell_t) = \\
 &= \sum_{\substack{a_1 + \dots + a_{t+1} \leq n - m(t+1) \\ a_s \geq \ell_s \ell_{s-1} - 1 \ (1 \leq s \leq t+1)}} \binom{n - p(t+1) - \sum_{s=1}^{t+1} a_s}{t+1} \prod_{s=1}^{t+1} \frac{\sigma(a_s, \ell_s - \ell_{s-1})}{(a_s + m)!},
 \end{aligned}$$

where

$$\sigma(a_s, \ell_s - \ell_{s-1}) := S(\ell_s - \ell_{s-1} - 1, q, a_s - (\ell_s - \ell_{s-1} - 1)).$$

Now we have

$$\begin{aligned}
 X(t) &= \sum_{j=t+1}^{n-p} (-1)^j X_j(t) = \sum_{j=t+1}^{n-p} (-1)^j \sum_{1 \leq \ell_1 < \dots < \ell_t < j} S_j(\ell_1, \dots, \ell_t) = \\
 &= \sum_{\substack{a_1 + \dots + a_{t+1} \leq n - m(t+1) \\ a_s \geq 0 \ (1 \leq s \leq t+1)}} \binom{n - p(t+1) - \sum_{s=1}^t a_s}{t+1} \cdot \prod_{s=1}^{t+1} \frac{1}{(a_s + m)!} \cdot F(a_1, \dots, a_{t+1}),
 \end{aligned}$$

where

$$F(a_1, \dots, a_{t+1}) = \sum_{j=t+1}^{n-p} (-1)^j \sum_{\substack{1 \leq \ell_1 < \dots < \ell_t < j \\ \ell_s - \ell_{s-1} \leq a_s + 1 \ (1 \leq s \leq t+1)}} \prod_{s=1}^{t+1} \sigma(a_s, \ell_s - \ell_{s-1}).$$

Since

$$j = \ell_{t+1} - \ell_0 = \sum_{s=1}^{t+1} (\ell_s - \ell_{s-1}),$$

we get

$$\begin{aligned}
 F(a_1, \dots, a_{t+1}) &= \sum_{j=t+1}^{n-p} \sum_{\substack{1 \leq \ell_1 < \dots < \ell_t < j \\ \ell_s - \ell_{s-1} \leq a_s + 1 \quad (1 \leq s \leq t+1)}} \prod_{s=1}^{t+1} (-1)^{\ell_s - \ell_{s-1}} \sigma(a_s, \ell_s - \ell_{s-1}) = \\
 &= \prod_{s=1}^{t+1} \sum_{h=0}^{a_s} (-1)^{h+1} S(h, q, a_s - h) = \prod_{s=1}^{t+1} f_m(a_s) \quad \text{by (2)}.
 \end{aligned}$$

We summarize. Writing  $t + 1 = k$  we arrive at the formula

$$b(n, m) = 1 + \sum_{k=1}^{\lfloor \frac{n}{m} \rfloor} S_k(n, m),$$

where

$$S_k(n, m) = \sum_{\substack{a_j \geq 0 \quad (1 \leq j \leq k) \\ \sum_{j=1}^k a_j \leq n - mk}} \binom{n - (m - 1)k - \sum_{j=1}^k a_j}{k} \prod_{j=1}^k \frac{f_m(a_j)}{(a_j + m)!}$$

with

$$f_m(a) = \begin{cases} -1 & \text{if } a \equiv 0 \pmod{m}, \\ 1 & \text{if } a \equiv 1 \pmod{m}, \\ 0 & \text{else.} \end{cases}$$

**Proof of (1.3).** Put  $S_0(n, m) = 1$  and introduce

$$F_m(z) := \sum_{a=0}^{\infty} \frac{f_m(a)}{(a + m)!} z^a.$$

Then we have

$$\begin{aligned}
 \sum_{n=0}^{\infty} b(n, m) z^n &= \sum_{n=0}^{\infty} \left( \sum_{0 \leq k \leq \frac{n}{m}} S_k(n, m) \right) z^n = \\
 &= \sum_{k=0}^{\infty} \sum_{n \geq mk} S_k(n, m) z^n = \\
 &= \sum_{k=0}^{\infty} z^{mk} \sum_{\ell=0}^{\infty} S_k(\ell + mk, m) z^\ell.
 \end{aligned}$$

The inner sum is equal to

$$\sum_{\ell=0}^{\infty} \left( \sum_{\substack{a_j \geq 0 \\ \sum_{j=1}^k a_j \leq \ell}} \binom{\ell + k - \sum_{j=1}^k a_j}{k} \prod_{j=1}^k \frac{f_m(a_j)}{(a_j + m!)} \right) z^\ell = (1 - z)^{-(k+1)} F_m(z)^k.$$

Therefore

$$\sum_{n=0}^{\infty} b(n, m) z^n = \frac{1}{1 - z} \sum_{k=0}^{\infty} \left( \frac{z^m F_m(z)}{1 - z} \right)^k = \frac{1}{1 - z - z^m F_m(z)} = P_m(z)^{-1}.$$

#### 4. The proof of (1.5)

We have

$$\sum_{n=0}^{\infty} b(n) z^n = P_3(z)^{-1} \quad \text{with} \quad P_3(z) = \sum_{\substack{k=0 \\ k \equiv 0 \pmod{3}}}^{\infty} \frac{z^k}{k!} - \sum_{\substack{k=0 \\ k \equiv 1 \pmod{3}}}^{\infty} \frac{z^k}{k!}.$$

Putting  $\xi := \exp\left(\frac{2\pi i}{3}\right)$  we get

$$P_3(z) = \frac{1}{3} (\xi - 1) \exp(\xi z) (\xi^{-1} - \exp(\xi(\xi - 1)z)).$$

Therefore the zeros of  $P_3(z)$  are

$$w_\ell = \frac{2\pi}{\sqrt{3}} \left( \frac{1}{3} - \ell \right) \quad (\ell = 0, \pm 1, \dots).$$

We find

$$P'_3(w_\ell) = (-1)^{\ell+1} \exp\left(\frac{\pi}{\sqrt{3}}(3\ell - 1)\right).$$

The residue theorem yields

$$b(n) \sim \sum_{\ell=-\infty}^{\infty} \frac{1}{-P_3'(w_\ell)} \frac{1}{w_\ell^{n+1}}.$$

In particular we get (1.5).

## 5. The proof of (1.4)

Since  $m = 3$  has been settled we may assume  $m \geq 4$ . We show first that  $P_m(z)$  has a zero on the open interval

$$I(m) = \left] 1 + \frac{1}{m!}(1 - \alpha(m)), 1 + \frac{1}{m!}(1 + \beta(m)) \right[ ,$$

where

$$\alpha(m) = \frac{m! + 1}{(m+1)! + 1}, \quad \beta(m) = \frac{2(m+1)}{m! - 2(m+1)}.$$

We have

$$P_m(x) = 1 - x + \frac{x^m}{m!} \left( 1 - \frac{x}{m+1} \right) + S_m(x),$$

$$S_m(x) = \sum_{\ell=2}^{\infty} \frac{x^{m\ell}}{(m\ell)!} \left( 1 - \frac{x}{m\ell+1} \right).$$

Let  $1 \leq x \leq 2$ . Then  $S_m(x) > 0$  and

$$S_m(x) < \sum_{\ell=2}^{\infty} \left( \frac{x^m}{m!} \right)^\ell = \left( \frac{x^m}{m!} \right)^2 \frac{1}{1 - \frac{x^m}{m!}}.$$

From this we see that for  $1 \leq x \leq 2$  we have

$$(1) \quad P_m(x) > 1 - x + \frac{1}{m!} \left( 1 - \frac{x}{m+1} \right),$$

$$(2) \quad P_m < \frac{1}{1 - \frac{x^m}{m!}} - x.$$

In (1) take  $x = 1 + \frac{1}{m!}(1 + \delta)$ ,  $0 < \delta < 1$ . Then

$$P_m(x) > \frac{1}{m!} \left( \delta \left( 1 + \frac{1}{(m+1)!} \right) - \frac{1}{m+1} \left( 1 + \frac{1}{m!} \right) \right) = 0 \text{ for } \delta = \alpha(m).$$

In (2) take  $x = 1 + \frac{1}{m!}(1 + \delta)$ ,  $0 < \delta < 1$ , and put  $\epsilon = \frac{1 + \delta}{m!}$ . We get

$$P_m(x) < \frac{y}{m! - x^m},$$

where

$$\begin{aligned} y &= (1 + \epsilon)^{m+1} - (1 + \delta) \leq \\ &\leq \exp(\epsilon(m+1)) - (1 + \delta) \leq \\ &\leq (1 + 2\epsilon(m+1)) - (1 + \delta) = \\ &= \frac{2}{m!}(m+1) - \delta \left( 1 - \frac{2}{m!}(m+1) \right) = \\ &= 0 \text{ for } \delta = \beta(m). \end{aligned}$$

Next we show that  $P_m(z)$  has on the disk  $|z| \leq 1 + \frac{1}{m}$  just one zero and that this zero is simple.

This will follow from Rouché’s theorem if

$$|P_m(z) - (1 - z)| < |1 - z| \text{ for } |z| = 1 + \frac{1}{m}.$$

Since

$$\begin{aligned} |P_m(z) - (1 - z)| &= \left| \sum_{\ell=1}^{\infty} \frac{z^{m\ell}}{(m\ell)!} \left( 1 - \frac{z}{m\ell+1} \right) \right| \leq \\ &\leq \left( 1 + \frac{|z|}{m+1} \right) \sum_{\ell=1}^{\infty} \left( \frac{|z|}{m!} \right)^\ell \end{aligned}$$

and since  $|1 - z| \geq |z| - 1$ , it is enough to show that

$$\left( 1 + \frac{|z|}{m+1} \right) \sum_{\ell=1}^{\infty} \left( \frac{|z|^m}{m!} \right)^\ell < |z| - 1 \text{ for } |z| = 1 + \frac{1}{m}.$$

Since  $\left(1 + \frac{1}{m}\right)^m < 3$  the left side is

$$< \left(1 + \frac{1}{m}\right) \frac{3}{m!} \sum_{\ell=0}^{\infty} \left(\frac{3}{4!}\right)^{\ell} \leq \frac{5}{7} \frac{1}{m} < \frac{1}{m}.$$

Denote the unique zero of  $P_m(z)$  on the disk  $|z| \leq 1 + \frac{1}{m}$  by  $\xi(m)$ . Since it is simple we have  $P'_m(\xi(m)) \neq 0$ . Since

$$1 + \frac{1}{m!}(1 + \beta(m)) < 1 + \frac{1}{m} \quad (m \geq 4)$$

we have  $\xi(m) \in I(m)$ .

The residue theorem yields

$$b(n, m) = \frac{c(m)}{\xi(m)^n} + O\left(\left(\frac{m}{m+1}\right)^n\right) \quad \text{as } n \rightarrow \infty,$$

where

$$c(m) = -(\xi(m)P'_m(\xi(m)))^{-1}.$$

## 6. An invariance property of $b(n, m; \sigma)$

Let  $\tau \in S_n$  be given by  $\tau(k) = n+1-k$  ( $1 \leq k \leq n$ ). Let  $\rho \in S_m$  be given by  $\rho(\ell) = m+1-\ell$  ( $1 \leq \ell \leq m$ ). One finds

$$\pi \in \mathcal{B}(n, m; \sigma) \iff \pi \circ \tau \in \mathcal{B}(n, m; \rho \circ \sigma),$$

$$\pi \in \mathcal{B}(n, m; \sigma) \iff \tau \circ \pi \in \mathcal{B}(n, m; \sigma \circ \rho).$$

From this we deduce

$$b(n, m; \sigma) = b(n, m; \rho \circ \sigma) = b(n, m; \sigma \circ \rho) = b(n, m; \rho \circ \sigma \circ \rho).$$

We apply this on

$$\begin{aligned} S_3 &= \left\{ \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}, \right. \\ &\quad \left. \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \right\} = \\ &= \{\epsilon, \rho, \sigma_1, \sigma_2, \sigma_3, \sigma_4\} \end{aligned}$$

and see that  $b(n, 3; \sigma)$  is the same for  $\sigma = \epsilon, \rho (= b(n))$  and for  $\sigma = \sigma_1, \sigma_2, \sigma_3, \sigma_4$ . We select  $\sigma_1$  as a sample for our analysis.

$B(n, 3; \sigma_1)$  is the set of all  $\pi \in S_n$  with the property that

$$\pi(k + 1) < \pi(k) < \pi(k + 2) \text{ is false for } 1 \leq k \leq n - 2.$$

The fraction

$$\tilde{b}(n) := \frac{|B(n, 3; \sigma_1)|}{n!}$$

will be studied in the next section.

### 7. A formula for $\tilde{b}(n)$ and proof of (1.6), (1.7)

The sieve formula yields

$$\tilde{b}(n) = 1 + \sum_{j=1}^{n-2} (-1)^j \sum_{1 \leq k_1 < \dots < k_j \leq n-2} \frac{\tilde{D}(k_1, \dots, k_j)}{n!},$$

where

$$\tilde{D}(k_1, \dots, k_j) = \# \{ \pi \in S_n \mid \pi(k_i + 1) < \pi(k_i) < \pi(k_i + 2) \text{ for } 1 \leq i \leq j \}.$$

We find

$$\tilde{D}(k_1, \dots, k_j) = \frac{n!}{\tilde{M}(k_1, \dots, k_j)},$$

where  $\tilde{M}(k_1, \dots, k_j)$  is defined in the following way.

$$j = 1 : \tilde{M}(k) = 3!$$

$j \geq 2 : \tilde{D}(k_1, \dots, k_j) = 0$  if there is some  $i_0$  ( $1 \leq i_0 < j$ ) such that  $k_{i_0+1} - k_{i_0} = 1$ .

Now suppose that  $k_{i+1} - k_i \geq 2$  ( $1 \leq i < j$ ). Put

$$L(k_1, \dots, k_j) := \{ i \mid 1 \leq i < j, k_{i+1} - k_i \geq 3 \}.$$

If  $L(k_1, \dots, k_j) = \emptyset$  then

$$\tilde{M}(k_1, \dots, k_j) = (2j + 1)2^j j!.$$

If  $L(k_1, \dots, k_j) \neq \emptyset$  write

$$L(k_1, \dots, k_j) = \{\ell_1, \dots, \ell_t\} \quad \text{with } 1 \leq \ell_1 < \dots < \ell_t < j.$$

Further put  $\ell_0 = 0$ ,  $\ell_{t+1} = j$  and define  $d_s := \ell_{s+1} - \ell_s$  ( $0 \leq s \leq t$ ). Then

$$\tilde{M}(k_1, \dots, k_j) = \prod_{s=0}^t (2d_s + 1)2^{d_s} d_s!.$$

We must leave the proof of this to the reader. We now have

$$\tilde{b}(n) = 1 - \frac{n-2}{3!} + \sum_{2 \leq j \leq \frac{n-1}{2}} (-1)^j S(j),$$

where ( $j \geq 2$ )

$$S(j) = \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n-2 \\ k_{i+1} - k_i \geq 2 \quad (1 \leq i < j)}} \tilde{M}(k_1, \dots, k_j)^{-1}.$$

For  $L \subset \{1, \dots, j-1\}$  put

$$S(j; L) := \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n-2 \\ k_{i+1} - k_i \geq 2 \quad (1 \leq i < j) \\ L(k_1, \dots, k_j) = L}} \tilde{M}(k_1, \dots, k_j)^{-1}.$$

Then we have ( $j \geq 2$ )

$$\begin{aligned} S(j) &= \sum_{L \subset \{1, \dots, j-1\}} S(j; L) = \\ &= S(j; \emptyset) + \sum_{t=1}^{j-1} \sum_{1 \leq \ell_1 < \dots < \ell_t < j} S(j; \{\ell_1, \dots, \ell_t\}). \end{aligned}$$

We have

$$S(j; \emptyset) = \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n-2 \\ k_{i+1} - k_i = 2 \quad (1 \leq i < j)}} \frac{1}{(2j+1)2^j j!} = \frac{n-2j}{(2j+1)2^j j!}.$$

We have ( $t \geq 1$ )

$$S(j; \{\ell_1, \dots, \ell_t\}) = \left( \prod_{s=0}^t (2d_s + 1)2^{d_s} d_s! \right)^{-1} \cdot X,$$



where

$$\begin{aligned}
 X &= \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n-2 \\ k_{i+1} - k_i \geq 2 \ (1 \leq i < j) \\ L(k_1, \dots, k_j) = \{\ell_1, \dots, \ell_t\} =: L}} 1 = \\
 &= \sum_{\vartheta_\ell \geq 3 \ (\ell \in L)} \sum_{\substack{1 \leq k_1 < \dots < k_j \leq n-2 \\ k_{i+1} - k_i = 2 \ (i \notin L) \\ k_{\ell+1} - k_\ell = \vartheta_\ell \ (\ell \in L)}} 1 = \\
 &= \sum_{\substack{\delta_\ell \geq 0 \ (\ell \in L) \\ \sum_{\ell \in L} \delta_\ell \leq n-2j-t}} \binom{n-2j-t - \sum_{\ell \in L} \delta_\ell}{t+1} = \\
 &= \binom{n-2j}{t+1} \text{ by (3.1).}
 \end{aligned}$$

Altogether we get  $\left(2 \leq j \leq \frac{n-1}{2}\right)$

$$S(j) = \sum_{0 \leq t \leq \min\{j-1, n-2j-1\}} \binom{n-2j}{t+1} \sigma(j, t)$$

with

$$\sigma(j, 0) = \frac{1}{(2j+1)2^j j!}$$

and  $(t \geq 1)$

$$\begin{aligned}
 \sigma(j, t) &= \sum_{1 \leq \ell_1 < \dots < \ell_t < j} \left( \prod_{s=0}^t (2d_s + 1) 2^{d_s} d_s! \right)^{-1} = \\
 &= \sum_{\substack{d_s \geq 1 \ (0 \leq s \leq t) \\ \sum_{s=0}^t d_s = j}} \left( \prod_{s=0}^t (2d_s + 1) 2^{d_s} d_s! \right)^{-1}.
 \end{aligned}$$

We arrive at the formula

$$\tilde{b}(n) = 1 + \sum_{0 \leq k \leq \frac{n-3}{2}} (-1)^{k+1} \sum_{t=0}^k \binom{n-2k-2}{t+1} S(k, t),$$

where

$$S(k, t) = \sum_{\substack{a_s \geq 0 \\ (0 \leq s \leq t)}} \left( \prod_{s=0}^t (2a_s + 3) 2^{a_s+1} (a_s + 1)! \right)^{-1} \cdot \sum_{s=0}^t a_s = k - t$$

**Proof of (1.6).** The formula above gives

$$\sum_{n=0}^{\infty} \tilde{b}(n) z^n = \frac{1}{1-z} + S(z),$$

where

$$\begin{aligned} S(z) &= \sum_{n=3}^{\infty} \left( \sum_{0 \leq k \leq \frac{n-3}{2}} (-1)^{k+1} \sum_{t=0}^k \binom{n-2k-2}{t+1} S(k, t) \right) z^n = \\ &= \sum_{k=0}^{\infty} (-1)^{k+1} \sum_{t=0}^k S(k, t) \sum_{n=2k+3}^{\infty} \binom{n-2k-2}{t+1} z^n = \\ &= z^3 \sum_{k=0}^{\infty} (-1)^{k+1} z^{2k} \sum_{t=0}^k S(k, t) \frac{z^t}{(1-z)^{t+2}} = \\ &= -\frac{z^3}{(1-z)^2} \sum_{t=0}^{\infty} \left( \frac{-z}{1-z} \right)^t \sum_{\ell=0}^{\infty} (-1)^{\ell} S(\ell+t, t) z^{2\ell}. \end{aligned}$$

Since

$$S(\ell+t, t) = \sum_{\substack{a_s \geq 0 \\ (0 \leq s \leq t)}} \left( \prod_{s=0}^t (2a_s + 3) 2^{a_s+1} (a_s + 1)! \right)^{-1},$$

the inner sum above is equal to  $F(z)^{t+1}$ , where

$$F(z) := \sum_{a=0}^{\infty} \frac{(-1)^a}{(2a+3)2^{a+1}(a+1)!} z^{2a}.$$

Therefore

$$\begin{aligned} S(z) &= -\frac{z^3}{(1-z)^2} \sum_{t=0}^{\infty} \left( -\frac{z^3}{1-z} F(z) \right)^t = \\ &= -\frac{z^3}{1-z} \frac{F(z)}{1-z+z^3F(z)}. \end{aligned}$$

We conclude

$$\sum_{n=0}^{\infty} \tilde{b}(n)z^n = \frac{1}{1-(z-z^3F(z))}.$$

But

$$z - z^3F(z) = \int_0^z \exp\left(-\frac{1}{2}w^2\right) dw.$$

**Proof of (1.7).** Put

$$E(z) := \int_0^z \exp\left(-\frac{1}{2}w^2\right) dw.$$

Since  $E(1) < 1$ ,  $E(\infty) = \sqrt{\frac{\pi}{2}} > 1$  and  $E(x)$  is strictly increasing for  $x \geq 0$ , there is a unique  $\xi > 1$  with  $E(\xi) = 1$ . (1.7) follows with the theorem of residues if

$$(1) \quad E(z) \neq 1 \text{ for all } z, |z| \leq \xi \text{ and } x \neq \xi.$$

We first show that

$$(2) \quad z = x + iy, y \neq 0, |z| \leq \sqrt{\pi} \Rightarrow \text{Im}E(z) \neq 0.$$

Namely from

$$\text{Im}E(z) = \exp\left(-\frac{1}{2}x^2\right) \int_0^y \exp\left(\frac{1}{2}t^2\right) \cos(xt) dt$$

we see that

$$\text{Im}E(z) \neq 0 \text{ if } y \neq 0 \text{ and } |xy| \leq \frac{\pi}{2}.$$

But  $|xy| \leq \frac{\pi}{2}$  if  $|z| \leq \sqrt{\pi}$ .

From (2) we infer

(3) Let  $0 < r \leq \sqrt{\pi}$ . Then

$$E(z) \neq E(r) \text{ for all } z, |z| \leq r \text{ and } z \neq r.$$

**Proof of (1).** According to (3) it is enough to establish  $\xi \leq \sqrt{\pi}$ . But this holds true since  $E(\sqrt{\pi}) > 1$ .

**Concluding remark.** Comparing (1.5), (1.7) one should know if  $\frac{3\sqrt{3}}{2\pi} > \xi^{-1}$ ,  $\frac{2\pi}{3\sqrt{3}} < \xi$ . This is true if  $E\left(\frac{2\pi}{3\sqrt{3}}\right) < 1$ .

One has (courtesy Dr. Harald Schmidt, Universität Regensburg)

$$E\left(\frac{2\pi}{3\sqrt{3}}\right) = 0.9693304740 \dots$$

## 8. $m = 4$ , outlook

We identified 7 classes of  $S_4$  on each of which  $b(n, 4; \sigma)$  is the same:

$$C_1 = \{e, \rho\}$$

$$C_2 = \{(1423), (4132), (2314), (3241)\} \cup \{(4123), (1432), (2341), (3214)\},$$

$$C_3 = \{(1243), (4312), (2134), (3421)\},$$

$$C_4 = \{(1342), (3124), (4213), (2431)\},$$

$$C_5 = \{(1324), (4231)\},$$

$$C_6 = \{(3142), (2413)\},$$

$$C_7 = \{(3412), (2143)\}.$$

We obtained

$$\sum_{n=0}^{\infty} b(n, 4; C_2) z^n = \left(1 - \int_0^z \exp\left(-\frac{1}{6}w^3\right) dw\right)^{-1}$$

from which we derive

$$b(n, 4; C_1) \sim \frac{\exp\left(\frac{1}{6}\xi_2^3\right)}{\xi_2^{n+1}} \text{ as } n \rightarrow \infty,$$

where  $\xi_2 > 1$  is given by  $\int_0^{\xi_2} \exp\left(-\frac{1}{6}t^3\right) dt = 1$ .

Further we got

$$\sum_{n=0}^{\infty} b(n, 4; C_3) z^n = F(z)^{-1} = (1 - z - R(z))^{-1},$$

where

$$R(z) = \sum_{m=1}^{\infty} (-1)^m \frac{\prod_{k=0}^{m-1} (3k+1)}{(3m+1)!} z^{3m+1}.$$

We have  $F(1) > 0$ ,  $F\left(\frac{3}{2}\right) > 0$  and  $F(z)$  has on the disk  $|z| \leq \frac{3}{2}$  precisely one zero  $\xi_3$  and it is simple. Therefore

$$b(n, 4; C_3) \sim \frac{1}{(-F'(\xi_3))\xi_3^{n+1}} \quad \text{as } n \rightarrow \infty.$$

For the remaining classes we have no results.

### **R. Warlimont**

The John Knopfmacher Centre for Applicable Analysis and Number Theory  
University of the Witwatersrand  
WITS, 2050  
Johannesburg, South Africa  
dikeledi@mweb.co.za