

**ON THE AVERAGE OF
 $d(n)\omega(n)$ AND SIMILAR FUNCTIONS
ON SHORT INTERVALS**

J.-M. DeKoninck (Québec, Canada)

I. Kátai (Budapest, Hungary)

1. Introduction

For each integer $n \geq 2$, let $\omega(n)$, $\Omega(n)$ and $\tau(n)$ stand for the number of distinct prime divisors of n , the number of prime divisors of n counting their multiplicity and the number of positive divisors of n , respectively, with $\omega(1) = \Omega(1) = 0$ and $\tau(1) = 1$. Given an integer $n \geq 2$, let $\beta(n)$ be the sum of the distinct prime divisors of n . Moreover, given any positive integer k and any complex number z , let

$$\tau_k(n) = \#\{(d_1, d_2, \dots, d_k) : d_1 d_2 \dots d_k = n, d_i \in \mathbb{N}\}.$$

Finally, let $x_1 = \log x$, $x_{j+1} = \log x_j$ for each integer $j \geq 1$.

It was shown by De Koninck and Ivić [1], using analytic methods, that as $x \rightarrow \infty$,

$$(1) \quad \sum_{n \leq x} \tau(n)\omega(n) = 2xx_1x_2 + Axx_1 + O(x),$$

where

$$A = 2 \left(\sum_p \left(\log \left(1 - \frac{1}{p} \right) + \left(\frac{1}{p} + \frac{3}{2p^2} + \frac{4}{2p^3} + \dots \right) \left(1 - \frac{1}{p} \right)^2 \right) - \Gamma'(2) \right),$$

The first author supported in part by a grant from NSERC.

The second author supported by the Hungarian National Foundation for Scientific Research under grant OTKA T046993 and the fund of the Applied Number Theory Research Group of the Hungarian Academy of Sciences and a grant from NSERC.

where Γ stands for the Gamma function.

This result was later improved by Sitaramachandrarao [10] who showed that

$$\sum_{n \leq x} \tau(n)\omega(n) = 2xx_1x_2 + Axx_1 + Bxx_2 + Cx + O\left(\frac{x}{x_1}\right),$$

with explicit constants $B \neq 0$ and C . Observe that this means that the term C_2xx_2 is missing in (1).

Later Ivić [4] investigated asymptotic formulas for sums of the type $\sum_{n \leq x} f(n)g(n)$, where f (resp. g) belong to certain classes of multiplicative (resp. additive) functions. He did so by considering the generating function

$$(2) \quad \sum_{n=1}^{\infty} \frac{f(n)z^{g(n)}}{n^s},$$

and then applying the method of A. Selberg in order to compute the asymptotic expansion of $\sum_{n \leq x} f(n)z^{g(n)}$. He could carry over this argument when

$\sum_{n=1}^{\infty} \frac{f(n)z^{g(n)}}{n^s}$ was a product of $\zeta(s)^w$ and $A(s, w)$, where $A(s, w)$ is a function which is regular in $|s - 1| < \varepsilon$. In fact, Ivić proved the following two results.

Theorem A. *Let $k \geq 2$ be fixed and $N > k$ be an arbitrary but fixed integer. Then there exist computable constants $a_{k,j}, b_{k,j}, c_{k,j}$ ($a_{k,1} \neq 0$) such that*

$$\sum_{n \leq x} d_k(n)\omega(n) = x \sum_{j=1}^k (a_{k,j}x_2 + b_{k,j})x_1^{k-j} + x \sum_{j=k+1}^N c_{k,j}x_1^{k-j} + O(xx_1^{k-N-1}).$$

Theorem B. *Let $m, N \geq 1$ and $k \geq 2$ be fixed integers. Then there exist polynomials $P_{k,m,j}(t)$ ($j = 1, \dots, N$) of degree m in t with computable coefficients such that*

$$\sum_{n \leq x} d_k(n)\omega^m(n) = x \sum_{j=1}^N P_{k,m,j}(x_2)x_1^{k-j} + O(xx_1^{k-N-1}x_2^m).$$

We shall provide here short interval versions of similar theorems. In particular, we shall prove the following results.

Theorem 1. *Let $x^{7/12+\varepsilon} \leq h(x) \leq x$. Then, for each fixed integer k and suitable constants $B_0, B_1, E_1, E_2, \dots, E_k$,*

$$\begin{aligned} & \frac{1}{h(x)} \sum_{x \leq n \leq x+h(x)} \tau(n)\omega(n) = \\ & = B_0(x_1x_2 - x_1) + B_1(x_2 + 1) + \sum_{\nu=1}^k E_\nu x_1^{1-\nu} + O(x_1^{-k}x_2). \end{aligned}$$

In [2] De Koninck and Ivić proved that if $f(n) = \sum_{p|n} p^\rho L(p)$ for some $\rho > 0$, where $L(x)$ is a slowly oscillating function, and if $x^{7/12} \log^{22} x \leq h(x) \leq x$, then

$$\sum_{x \leq n \leq x+h(x)} f(n) = (\zeta(1+\rho) + o(1)) \frac{h(x)x^\rho L(x)}{\log x},$$

from which it follows, in particular, that

$$\frac{1}{h(x)} \sum_{x \leq n \leq x+h(x)} \frac{\beta(n)}{n} = (\zeta(1+\rho) + o(1)) \frac{1}{\log x}.$$

Here we show the following stronger result.

Theorem 2. *Let $x^{7/12+\varepsilon} \leq h(x) \leq x$. Then, for each fixed integer k and suitable constants D_1, D_2, \dots, D_k ,*

$$\frac{1}{h(x)} \sum_{x \leq n \leq x+h(x)} \frac{\beta(n)}{n} = \frac{D_1}{x_1} + \frac{D_2}{x_1^2} + \dots + \frac{D_k}{x_1^k} + O\left(\frac{1}{x_1^{k+1}}\right).$$

2. The Hooley-Huxley contour and Ramachandra's theorem

In 1976 K. Ramachandra [8] obtained short interval mean value theorems for those arithmetical functions such that the corresponding Dirichlet series may be written as finite products of powers of L -functions multiplied by the product of finitely many $\log L(s, \chi)$ functions and a certain function regular in

$$\Re(s) > \frac{1}{2}.$$

For our results, the main idea of the proof is to choose an appropriate line of integration in the Perron formula, namely the so-called Hooley-Huxley contour. To do so, first let S_1, S_2 and S_3 be the set of L -series, the set of their derivatives and the set of their logarithms, respectively. Observe that $\log L(s, \chi)$ is defined by analytic continuation from the halfplane $\sigma = \Re(s) > 1$; for each complex number z , we define

$$L(x, \chi)^z = \exp\{z \log L(s, \chi)\}.$$

Let $P_1(s)$ be any finite power product, with complex exponents, of functions of S_1 , and let $P_2(s)$ (resp. $P_3(s)$) be any finite power product, with non-negative integer exponents, of functions of S_2 (resp. S_3). Moreover, let c_n be a sequence of complex numbers such that $|c_n| \ll n^\varepsilon$ for every $\varepsilon > 0$ and

$$\sum_{n=1}^{\infty} \frac{|c_n|}{n^\sigma} < +\infty \quad \text{for } \sigma > \frac{1}{2}.$$

Let also $F_0(s) = \sum_{n=1}^{\infty} \frac{c_n}{n^s}$ and define the sequence g_1, g_2, \dots implicitly by

$$F_1(s) := P_1(s)P_2(s)P_3(s)F_0(s) = \sum_{n=1}^{\infty} \frac{g_n}{n^s}$$

and set

$$E(x) = \sum_{n \leq x} g_n.$$

Given a positive number $r \leq \frac{1}{2}$, we define the contour C_r by first considering the circle $\{s : |s-1| = r\}$, removing the point $1-r$, and proceeding on the remaining portion of the circle in the anticlockwise direction. Set $C_0 = C(r)$. Assume that r is small enough so that $F_1(s)$ has no singularities on the boundary and in the interior of C_0 , except possibly at the point $s = 1$.

Let $C_1 = C\left(\frac{1}{\log x}\right)$, and let L^-, L^+ be defined as the intervals on straight lines

$$\begin{aligned} L^- &= \left[\left(1 - \frac{1}{r}\right) e^{-i\pi}, \left(1 - \frac{1}{\log x}\right) e^{-i\pi} \right], \\ L^+ &= \left[\left(1 - \frac{1}{\log x}\right) e^{i\pi}, \left(1 - \frac{1}{r}\right) e^{i\pi} \right]. \end{aligned}$$

Let C^* be the contour going along L^- starting from $\left(1 - \frac{1}{r}\right) e^{-i\pi}$, then on C_1 , and finally on L^+ .

Let B be the constant appearing in the well known density result

$$N_\chi(\alpha, T) := \#\{\rho = \beta + i\gamma : L(\rho, \chi) = 0, \text{ with } \beta \geq \alpha \geq 0 \text{ and } |\gamma| \leq T\} = O\left(T^{B(1-\alpha)} \log^2 T\right),$$

which is valid for all characters χ occuring in P_1, P_2 and P_3 . Letting $\varphi = 1 - \frac{1}{B} - \varepsilon$, with an arbitrary $\varepsilon > 0$, Ramachandra [8] proved the following result.

Theorem (Ramachandra). *Let x be a large number and $1 \leq h(x) \leq x$. Set*

$$I(x, h(x)) = \frac{1}{2\pi i} \int_0^{h(x)} \left(\int_{C_0} F_1(s)(v+x)^{s-1} ds \right) dv.$$

Then

$$E(x+h(x)) - E(x) = I(x, h(x)) + O_\varepsilon\left(h(x) \cdot \exp\{-(\log x)^{1/6} \cdot x^\varphi\}\right).$$

Remark. According to Huxley's result [3], the number φ may be replaced by any constant greater than $7/12$.

Ramachandra used the Hooley-Huxley contour in order to prove his very general theorem. Later on, Kátai [5] applied Ramachandra's theorem to obtain that

$$\sum_{\substack{x \leq n \leq x+h(x) \\ \omega(n)=k}} 1 = (1 + o(1)) \frac{h(x) \cdot x_2^{k-1}}{(k-1)!x_1}$$

holds uniformly for $k \leq x_2 + c_x \sqrt{x_2}$, where c_x tends to $+\infty$ sufficiently slowly, and $h(x) \geq x^{\varphi+\varepsilon}$.

3. The proof of Theorem 1

Since

$$\sum_{m=1}^{\infty} \frac{z^{\omega(m)}}{m^s} = \zeta(s)^z G(s, z) \quad \text{and} \quad \sum_{m=1}^{\infty} \frac{z^{\Omega(m)}}{m^s} = \zeta(s)^z F(s, z),$$

where the functions $G(s, z)$ and $F(s, z)$ are regular in $\sigma > \frac{1}{2}$, it follows that the above Dirichlet series belong to the classes of functions satisfying Ramachandra's theorem. Kátai and Subbarao [6] used this to obtain asymptotic estimates of the expressions

$$(3) \quad \sum_{x \leq n \leq x+h(x)} z^{\omega(n)}, \quad \sum_{x \leq n \leq x+h(x)} z^{\omega(n)} |\mu(n)|, \quad \sum_{x \leq n \leq x+h(x)} 1/t_k(n),$$

where $h(x) = x^{7/12+\varepsilon}$. More generally, they proved that, assuming that $F(s)$ satisfies the conditions of Ramachandra's theorem, that $r > 0$ and $\varepsilon > 0$ are small numbers, that $x^{7/12+\varepsilon} \leq h(x) \leq x^{\frac{2}{3}-\frac{2r}{3}}$, and that $E(x) = \sum_{n \leq x} f(n)$ where $f(n) = z^{\omega(n)}$ or $z^{\omega(n)} |\mu(n)|$ or $1/\tau_k(n)$, then

$$\frac{E(x+h(x)) - E(x)}{h(x)} = \frac{1}{2\pi i} \int_{C^*} F(s) x^{s-1} ds + O\left(\exp\left\{-x_1^{1/6}\right\}\right),$$

where $C^* = \{s : |s-1| = 1/x_1, s \neq 1 - 1/x_1\}$.

Remark. Observe that the reason for the upper bound $x^{\frac{2}{3}-\frac{2r}{3}}$ on $h(x)$ is only due to a technical condition used in the proof; indeed one can show that the result is in fact valid for $x^{7/12+\varepsilon} \leq h(x) \leq x$.

Returning to the proof of Theorem 1, we let

$$F(s) = \sum_{n=1}^{\infty} \frac{\tau(n)\omega(n)}{n^s}.$$

One easily verifies that

$$\begin{aligned} F(s) &= \sum_p \sum_{\alpha=1}^{\infty} \frac{\tau(p^\alpha)}{p^{\alpha s}} \sum_{\substack{m=1 \\ (m,p)=1}}^{\infty} \frac{\tau(m)}{m^s} = \zeta^2(s) \sum_p \left(1 - \frac{1}{p^s}\right)^2 \sum_{\alpha=1}^{\infty} \frac{\alpha+1}{p^{\alpha s}} = \\ &= \zeta^2(s) \sum_p \left(1 - \frac{1}{p^s}\right)^2 \left\{ \frac{1}{(1-1/p^s)^2} - 1 \right\} = \zeta^2(s) \sum_p \left\{ \frac{2}{p^s} - \frac{1}{p^{2s}} \right\}. \end{aligned}$$

Since $\sum_p \frac{1}{p^s} = \log \zeta(s) - \sum_{r=2}^{\infty} \frac{1}{r} \sum_p \frac{1}{p^{rs}}$, it follows that

$$2 \sum_p \frac{1}{p^s} - \sum_p \frac{1}{p^{2s}} = 2 \log \zeta(s) - 2 \sum_p \frac{1}{p^{2s}} - \sum_{r=3}^{\infty} \frac{2}{r} \sum_p \frac{1}{p^{rs}} = 2 \log \zeta(s) - U(s),$$

say, where $U(s)$ is regular for $\Re(s) > \frac{1}{2}$, so that we may write

$$F(s) = F_1(s) - F_2(s), \quad \text{with } F_1(s) = \zeta^2(s) \cdot 2 \log \zeta(s) \text{ and } F_2(s) = \zeta^2(s)U(s).$$

Now define $\alpha_1(n)$ and $\alpha_2(n)$ implicitly by the representations

$$F_1(s) = \sum_{n=1}^{\infty} \frac{\alpha_1(n)}{n^s} \quad \text{and} \quad F_2(s) = \sum_{n=1}^{\infty} \frac{\alpha_2(n)}{n^s}.$$

Clearly both $F_1(s)$ and $F_2(s)$ belong to Ramachandra's class of functions. It follows that

$$\frac{1}{h(x)} \sum_{x \leq n \leq x+h(x)} \alpha_1(n) = \frac{1}{2\pi i} \int_{C^*} F_1(s) x^{s-1} ds + O\left(\exp\left\{-x_1^{1/6}\right\}\right).$$

Now we can write

$$\begin{aligned} F_1(s) &= (s-1)^2 \zeta^2(s) \cdot \frac{1}{(s-1)^2} \cdot 2 \left\{ \log((s-1)\zeta(s-1)) + \log \frac{1}{s-1} \right\} = \\ &= \frac{2 \log \frac{1}{s-1}}{(s-1)^2} ((s-1)^2 \zeta(s-1)) + \frac{2}{(s-1)^2} ((s-1)^2 \zeta(s)) \log((s-1)\zeta(s)). \end{aligned}$$

But since $(s-1)\zeta(s) \rightarrow 1$ as $s \rightarrow 1$, it follows that $(s-1)\zeta(s)$ and $\log((s-1)\zeta(s))$ are regular in the neighbourhood of 1.

Now define the constants B_0, B_1, \dots, B_k , C_0, C_1, \dots, C_k and the functions $U_k(s)$ and $V_k(s)$ implicitly by the relations

$$2(\zeta(s)(s-1))^2 = B_0 + B_1(s-1) + \dots + B_k(s-1)^k + U_k(s)(s-1)^{k+1},$$

$$2(\zeta(s)(s-1))^2 \log(\zeta(s)(s-1)) =$$

$$= C_0 + C_1(s-1) + \dots + C_k(s-1)^k + V_k(s)(s-1)^{k+1},$$

so that $U_k(s)$ and $V_k(s)$ are regular and bounded for $|s-1| \leq 1/x_1$. Thus

$$\frac{1}{2\pi i} \int_{C^*} F_1(s) x^{s-1} ds = \sum_{\nu=0}^k B_\nu \cdot \mathcal{I}_\nu + L_1 + \sum_{\nu=0}^k C_\nu \mathcal{J}_\nu + L_2,$$

where

$$\mathcal{I}_\nu = \frac{1}{2\pi i} \int_{C^*} \frac{x^{s-1}}{(s-1)^{2-\nu}} \log \frac{1}{s-1} ds \quad (0 \leq \nu \leq k),$$

$$\mathcal{J}_\nu = \frac{1}{2\pi i} \int_{C^*} \frac{x^{s-1}}{(s-1)^{2-\nu}} ds \quad (0 \leq \nu \leq k),$$

$$L_1 = \frac{1}{2\pi i} \int_{C^*} \frac{x^{s-1}}{(s-1)^{2-(k+1)}} U_k(s) \log(s-1) ds,$$

$$L_2 = \frac{1}{2\pi i} \int_{C^*} \frac{x^{s-1}}{(s-1)^{2-(k+1)}} V_k(s) \log(s-1) ds.$$

Since $s = 1 + \frac{e^{i\theta}}{x_1}$, $ds = \frac{1}{x_1} i e^{i\theta}$, $\log(s-1) = \log(1/x_1) + i\theta$ and $\log \frac{1}{s-1} = x_2 - i\theta$, it follows that, taking into account that $x^{e^{i\theta} \cdot x_1^{-1}} = e^{e^{i\theta}}$, we have

$$\begin{aligned} \mathcal{I}_\nu &= \frac{x_1^{2-\nu}}{2\pi x_1} \int_{-\pi}^{\pi} e^{e^{i\theta}} e^{(\nu-2)i\theta} (x_2 - i\theta) e^{i\theta} d\theta = \\ &= \frac{x_1^{1-\nu} \cdot x_2}{2\pi} \int_{-\pi}^{\pi} e^{e^{i\theta}} \cdot e^{i(\nu-1)\theta} d\theta - \frac{i x_1^{1-\nu}}{2\pi} \int_{-\pi}^{\pi} e^{e^{i\theta}} \cdot e^{i(\nu-1)\theta} \theta d\theta. \end{aligned}$$

Define $\eta_0 = 0$ and observe that, for $h \neq 0$, we have

$$\begin{aligned} \eta_h &:= \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ih\theta} \theta d\theta = \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(\frac{e^{ih\theta}}{ih} \right)' \theta d\theta = \\ &= \frac{1}{2\pi} \left[\frac{e^{ih\theta}}{ih} \theta \right]_{-\pi}^{\pi} - \frac{1}{2\pi ih} \int_{-\pi}^{\pi} e^{ih\theta} d\theta = \\ &= \frac{1}{2\pi ih} [\pi e^{ih\theta} + \pi e^{-ih\theta}] = \\ &= \frac{(-1)^h}{ih}. \end{aligned}$$

Therefore, using the representation $e^{e^{i\theta}} = \sum_{t=0}^{\infty} \frac{1}{t!} e^{it\theta}$, it follows from this that

$$(4) \quad \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{e^{i\theta}} e^{i(\nu-1)\theta} d\theta = \begin{cases} 0 & \text{if } \nu \geq 2, \\ 1 & \text{if } \nu = 0, 1, \end{cases}$$

and that

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{e^{i\theta}} e^{i(\nu-1)\theta} \theta d\theta = \sum_{t=0}^{\infty} \frac{1}{t!} \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(t+\nu-1)\theta} \theta d\theta = \frac{1}{i} \sum_{\substack{t=0 \\ t \neq 1-\nu}}^{\infty} \frac{1}{t!} \frac{(-1)^{t+\nu-1}}{t+\nu-1}.$$

Gathering these estimates, we obtain that

$$\mathcal{I}_0 = x_1 x_2 - x_1, \quad \mathcal{I}_1 = x_2 + 1, \quad \mathcal{I}_\nu = -x_1^{1-\nu} \mathcal{D}_\nu \quad \text{for } \nu \geq 2,$$

where each \mathcal{D}_ν is a computable constant.

On the other hand, it follows from (4) that

$$\mathcal{I}_\nu = \frac{x_1^{1-\nu}}{2\pi} \int_{-\pi}^{\pi} e^{e^{i\theta}} e^{i(\nu-1)\theta} d\theta = \begin{cases} x_1 & \text{if } \nu = 0, \\ 1 & \text{if } \nu = 1, \\ 0 & \text{if } \nu \geq 2. \end{cases}$$

Moreover, we have

$$|L_1| \leq x_1^{-k} \int_{-\pi}^{\pi} \left| U_k \left(1 + \frac{1}{x_1} e^{i\theta} \right) \right| \cdot (x_2 + |\theta|) d\theta \ll x_2 \cdot x_1^{-k}$$

and one can also easily establish that

$$|L_2| \ll x_1^{-k}.$$

Combining these estimates, the proof of Theorem 1 is thus complete.

Remark 1. From the Ramachandra's theorem, it follows that under the assumption

$$N(\sigma, T) \ll T^{(2+\varepsilon)(1-\sigma)},$$

which is somewhat weaker than the Riemann hypothesis, Theorem 1 holds for the shorter interval $x^{\frac{1}{2}+\varepsilon} \leq h(x) \leq x$.

Remark 2. It is clear from the proof of Theorem 1 that similar estimates can be obtained for the following sums:

$$\sum_{x \leq n \leq x+h(x)} \tau_k(n) \omega(n), \quad \sum_{x \leq n \leq x+h(x)} \tau_k(n) \Omega(n), \quad \sum_{x \leq n \leq x+h(x)} r(n) \omega(n),$$

where $r(n) = \#\{(u, v) : n = u^2 + v^2\}$.

4. The proof of Theorem 2

We begin by writing

$$\begin{aligned} F(s) &:= \sum_{n=1}^{\infty} \frac{\beta(n)}{n} = \left[\sum_p \left\{ \sum_{\alpha=1}^{\infty} \frac{p}{p^{\alpha s}} \right\} \left(1 - \frac{1}{p^s} \right) \right] \zeta(s) = \\ &= \zeta(s) \left\{ \sum_p \frac{1}{p^{s-1}} \right\} = \{\log \zeta(s-1) + u(s)\} \zeta(s), \end{aligned}$$

say, with $u(s)$ bounded and regular for $\Re(s) > 2$. Thus

$$\begin{aligned} F(s) &= \left(\log \frac{1}{s-2} \right) \zeta(s) + \zeta(s) [\log((s-1)\zeta(s-1)) + u(s)] = \\ &= \left(\log \frac{1}{s-2} \right) \zeta(s) + \zeta(s)v(s), \end{aligned}$$

say, with $v(s)$ regular for $\Re(s) > 2$. It follows that

$$F(s+1) = \sum_{n=1}^{\infty} \frac{\beta(n)/n}{n^s} = \zeta(s+1) \log \frac{1}{s-1} + \zeta(s+1)v(s+1).$$

Now observe that

$$\begin{aligned} &\frac{1}{h(x)} \sum_{x \leq n \leq x+h(x)} \frac{\beta(n)}{n} = \\ &= \frac{1}{x} \cdot \frac{1}{h(x)} \sum_{x \leq n \leq x+h(x)} \beta(n) - \frac{1}{h(x)} \sum_{x \leq n \leq x+h(x)} \beta(n) \left(\frac{1}{n} - \frac{1}{x} \right) = \\ &= \frac{1}{x} \cdot \frac{1}{h(x)} \sum_{x \leq n \leq x+h(x)} \beta(n) + O \left(\frac{1}{x} \sum_{x \leq n \leq x+h(x)} \frac{\beta(n)}{n} \right). \end{aligned}$$

Hence, proceeding as in the proof of Theorem 1, we get that

$$\begin{aligned} \frac{1}{h(x)} \sum_{x \leq n \leq x+h(x)} \frac{\beta(n)}{n} &= \frac{1}{2\pi i} \int_{C^*} F(s+1)x^{s-1} ds + O \left(\exp \left\{ -x_1^{1/6} \right\} \right) = \\ &= \frac{1}{2\pi i} \int_{C^*} x^{s-1} \zeta(s+1) \log \frac{1}{s-1} ds = \\ &= \sum_{\nu=0}^{\infty} a_{\nu} T_{\nu}, \end{aligned}$$

where the a_ν 's are defined implicitly by

$$\zeta(s+1) = \zeta(2) + \zeta'(2)(s-1) + \dots = a_0 + a_1(s-1) + \dots$$

and where the T_ν 's can be written as

$$\begin{aligned} T_\nu &= \frac{1}{2\pi i} \int_{C^*} (s-1)^\nu x^{s-1} \log \frac{1}{s-1} ds = \\ &= \frac{1}{2\pi} \frac{1}{x_1^{\nu+1}} \int_{-\pi}^{\pi} e^{i(\nu+1)\theta} e^{e^{i\theta}} (x_2 - i\theta) d\theta. \end{aligned}$$

Collecting the above estimates completes the proof of Theorem 2.

References

- [1] **DeKoninck J.-M. and Ivić A.**, *Topics in arithmetical functions*, North-Holland, Amsterdam, 1980.
- [2] **DeKoninck J.-M. and Ivić A.**, The average prime divisor of an integer in short intervals, *Arch. Math.*, **52** (1989), 440-448.
- [3] **Huxley M.N.**, On the difference between consecutive primes, *Invent. Math.*, **15** (1972), 164-170.
- [4] **Ivić A.**, Sums of products of certain arithmetical functions, *Publ. Inst. Math. (Beograd) N.S.*, **41** (55) (1987), 31-41.
- [5] **Kátai I.**, A remark on a paper of K. Ramachandra, *Number theory. Proc. Ootacamund*, ed. K. Alladi, Lecture Notes in Math. **1122**, Springer, 1984, 147-152.
- [6] **Kátai I. and Subbarao M.V.**, Some remarks on a paper of Ramachandra, *Lietuvos mat. rink.*, **43** (4) (2003), 497-506.
- [7] **Левин Б.В. и Тимофеев Н.М.**, Распределение арифметических функций в среднем по прогрессиям (теоремы типа Виноградова-Бомбьери), *Мат. сборник*, **125 (167)** (1984) (4) (12), 558-572. (**Levin B.V. and Timofeev N.M.**, Distribution of arithmetical functions in the mean over progressions (theorems of A.I. Vinogradov - E. Bombieri type), *Math. USSR Sbornik*, **53** (1986), 563-579. (English version))
- [8] **Ramachandra K.**, Some problems of analytic number theory, *Acta Arith.*, **31** (1976), 313-324.

- [9] **Selberg A.**, Note on a paper by L.G. Sathe, *J. Indian Math. Soc.*, **18** (1954), 83-87.
- [10] **Sitaramachandrarao R.**, *Letter to Jean-Marie De Koninck dated March 3, 1986*.
- [11] **Wolke D.**, Über die mittlere Verteilung der Werte Zahlentheoretischer Funktionen auf Restklassen I., *Math. Ann.*, **202** (1973), 1-25.

(Received November 30, 2004)

J.-M. DeKoninck

Département de mathématiques
Université Laval
Québec G1K 7P4
Canada
jmdk@mat.ulaval.ca

I. Kátai

Department of Computer Algebra
Eötvös Loránd University
Pázmány Péter s. 1/C.
H-1117 Budapest, Hungary
katali@compalg.inf.elte.hu