

ON THE PAIRS OF MULTIPLICATIVE FUNCTIONS SATISFYING A CONGRUENCE PROPERTY

Bui Minh Phong (Budapest, Hungary)

Dedicated to Professor Ferenc Schipp on his 70th anniversary

Dedicated to Professor Péter Simon on his 60th anniversary

Abstract. All solutions of the congruence

$$g(An + B) \equiv Cf(n) + D \pmod{n}$$

are given for integer-valued completely multiplicative functions f and g with some integers $A > 0$, $B > 0$, C and $D \neq 0$. We prove that, except for some special cases, there exist a non-negative integer α and a real-valued Dirichlet character $\chi_A \pmod{A}$ such that $n|f(n)$ and $g(m) = m^\alpha \chi_A(m)$ are satisfied for all $n, m \in \mathbb{N}$, $(m, A) = 1$. In the case when $C = 0$, we also determine all multiplicative functions g of the above congruence.

1. Introduction

Let k be a positive integer and let \mathbb{N}_k denote the set of the natural numbers coprime to k . An arithmetical function $g(n) \not\equiv 0$ is said to be multiplicative function on the set \mathbb{N}_k if $n, m \in \mathbb{N}_k$, $(n, m) = 1$ implies

$$g(nm) = g(n)g(m)$$

The European Union and the European Social Fund have provided financial support to the project under the grant agreement TÁMOP-4.2.1/B-09/1/KMR-2010-0003.

and it is called completely multiplicative on the set \mathbb{N}_k if this equation holds for all pairs of positive integers $n \in \mathbb{N}_k$ and $m \in \mathbb{N}_k$. In the following let \mathcal{M}_k (\mathcal{M}_k^*) be the set of integer-valued multiplicative (completely multiplicative) functions on the set \mathbb{N}_k . In the case $k = 1$, we use the following notations:

$$\mathbb{N} := \mathbb{N}_1, \mathcal{M} := \mathcal{M}_1 \quad \text{and} \quad \mathcal{M}^* := \mathcal{M}_1^*.$$

In 1966 M.V. Subbarao [8] proved the following assertion: If $g \in \mathcal{M}$ satisfies

$$(1) \quad g(n+m) \equiv g(m) \pmod{n} \quad \text{for all } n, m \in \mathbb{N},$$

then there is a non-negative integer α such that

$$(2) \quad g(n) = n^\alpha \quad \text{for all } n \in \mathbb{N}.$$

A. Iványi [2] extended this result proving that if $g \in \mathcal{M}^*$ and (1) holds for a fixed $m \in \mathbb{N}$ and for all $n \in \mathbb{N}$, then $g(n)$ has also the same form (2). In the joint paper with J. Fehér, we improved in [6] the results of Subbarao and Iványi mentioned above by proving that if $M \in \mathbb{N}$, $g \in \mathcal{M}$ satisfy the conditions $g(M) \neq 0$ and

$$g(n+M) \equiv g(M) \pmod{n} \quad \text{for all } n \in \mathbb{N},$$

then (2) holds.

Another characterization of n^α by using congruence property was found by A. Iványi [3], namely he proved that if $g \in \mathcal{M}$ satisfies the relation

$$(3) \quad g(n+m) \equiv g(n) + g(m) \pmod{n} \quad \text{for all } n, m \in \mathbb{N},$$

then $g(n)$ is a power of n with positive integer exponent. In [6] we determined all solutions $g \in \mathcal{M}^*$ of (3) under the condition that the congruence (3) holds for a fixed $m \in \mathbb{N}$ and for all $n \in \mathbb{N}$. Later, in the papers [3, 4, 5, 7] we obtained some generalizations of this result, namely we proved the following theorems:

Theorem A. ([3]) *If the integers $A > 0$, $B > 0$, $C \neq 0$, $N > 0$ with $(A, B) = 1$ and $g \in \mathcal{M}$ satisfy the relation*

$$g(An+B) \equiv C \pmod{n} \quad \text{for all } n \geq N,$$

then $g(B) = C$ and there are a non-negative integer α , a real-valued Dirichlet character $\chi_A \pmod{A}$ such that

$$g(n) = \chi_A(n)n^\alpha \quad \text{for all } n \in \mathbb{N}_A.$$

Theorem B. ([5]) *Let $A > 0$, B , $a > 0$, b , $N > 0$ and $D \neq 0$ be fixed positive integers. If a function $g \in \mathcal{M}^*$ satisfies the congruence*

$$g\left[A(an + b) + B\right] \equiv D \pmod{an + b} \quad \text{for all } n \in \mathbb{N}, n > N,$$

then there are a non-negative integer α , a real-valued Dirichlet character $\chi_{aA} \pmod{aA}$ such that

$$g(n) = \chi_{aA}(n)n^\alpha$$

holds for all $n \in \mathbb{N}_{aA}$.

Theorem C. [7] *Assume that $A > 0$, $B > 0$, C , $D \neq 0$ are fixed integers with $(A, B) = 1$ and a function $g \in \mathcal{M}^*$ satisfies the congruence*

$$g(An + B) \equiv Cg(n) + D \pmod{n} \quad \text{for all } n \in \mathbb{N}.$$

Then the following assertions hold:

(A) *If $g(p) = 0$ for some prime p with $(p, A) = 1$, then*

$$p = 2, \quad -C = D = 1, \quad (2, AB) = 1 \quad \text{and } g(n) = \chi_2(n) \quad \text{for all } n \in \mathbb{N}_2,$$

(B) *If $g(n) \neq 0$ for all $n \in \mathbb{N}_A$, then either*

$$C + D = 1 \quad \text{and } g(n) = 1 \quad \text{for all } n \in \mathbb{N},$$

or there are a non-negative integer α , a real-valued Dirichlet character $\chi_{aA} \pmod{aA}$ such that

$$g(n) = \chi_{aA}(n)n^\alpha$$

holds for all $n \in \mathbb{N}_{aA}$.

In [4] we completely solved the equation

$$g(An + B) \equiv g(An) + D \pmod{n}$$

for a multiplicative function g under conditions $A, B \in \mathbb{N}$, $D \in \mathbb{Z} \setminus \{0\}$ and $(A, B) = 1, (A, 2) = 1$.

The main purpose of this paper is to extend the result of Theorem A and prove a result similar to Theorem C for two completely multiplicative functions. We prove

Theorem 1. *Assume that the integers $A > 0, B > 0, N > 0, D \neq 0, E \neq 0$ and $g \in \mathcal{M}$ satisfy the relation*

$$Eg(An + B) \equiv D \pmod{n} \quad \text{for all } n \in \mathbb{N}, n > N.$$

Let $(A, B) = d$ and $A = da$. Then there are a non-negative integer α and a real-valued Dirichlet character $\chi_a \pmod{a}$ such that

$$g(dn) = g(d)\chi_a(n)n^\alpha \quad \text{for all } n \in \mathbb{N}_a.$$

We note that Theorem A is a special case of Theorem 1 when $E = 1$ and $(A, B) = 1$.

Theorem 2. *Assume that the integers $A > 0, B > 0, (A, B) = 1, C \neq 0, D \neq 0, E \neq 0$ and $f, g \in \mathcal{M}^*$ satisfy the relation*

$$Eg(An + B) \equiv Cf(n) + D \pmod{n} \quad \text{for all } n \in \mathbb{N}.$$

Then the following assertions hold:

(I) *If $f(\pi) = 0$ for some prime π , then*

(I. a) *there are a non-negative integer α and a real-valued Dirichlet character $\chi_A \pmod{A}$ such that*

$$n|f(n) \quad \text{and} \quad g(m) = \chi_A(m)m^\alpha \quad (n \in \mathbb{N}, m \in \mathbb{N}_A).$$

(I. b) *If $\pi|A$ and $(\pi, B) = 1$, then $C = -2D$ and all further solutions (f, g) have the form*

$$\pi = 2, f(n) = \chi_2(n) \quad \text{and} \quad g(m) = \chi_{2A}(m)m^\alpha \quad (n \in \mathbb{N}, m \in \mathbb{N}_A),$$

where α is a non-negative integer and χ_{2A} is a real-valued Dirichlet character $\pmod{2A}$ with the condition $\chi_{2A}(A + B) = -\chi_{2A}(B)$.

(I. c) *If $(\pi, AB) = 1$, then $C = -D$ and all further solutions (f, g) have the form*

$$\pi = 2, f(n) = \chi_2(n), g(2) = 0 \quad \text{and} \quad g(m) = \chi_{2A}(m)m^\alpha \quad (n \in \mathbb{N}, m \in \mathbb{N}_A),$$

where α is a non-negative integer and χ_{2A} is a real-valued Dirichlet character $\pmod{2A}$.

(II) If $g(An + B) = 0$ for some $N \in \mathbb{N}$ and $f(n) \neq 0$ for all $n \in \mathbb{N}$, then

$$f(n) = 1 \quad \text{for all } n \in \mathbb{N},$$

and either

$$g(An + B) = 0 \quad \text{for all } n \in \mathbb{N} \quad \text{if } C + D = 0,$$

or there are a non-negative integer α and a real-valued Dirichlet character χ_A such that

$$g(n) = \chi_A(n)n^\alpha \quad \text{for all } n \in \mathbb{N}_A \quad \text{if } C + D \neq 0.$$

(III) If $g(An + B)f(n) \neq 0$ for all $n \in \mathbb{N}$, then there are a non-negative α and a real-valued Dirichlet character $\chi_A \pmod{A}$ such that

$$g(n) = \chi_A(n)n^\alpha \quad \text{for all } n \in \mathbb{N}_A,$$

furthermore either

$$C + D = Eg(B) \quad \text{and } f(n) = 1 \quad \text{for all } n \in \mathbb{N}$$

or $n \mid f(n)$ for all $n \in \mathbb{N}$.

2. Proof of Theorem 1

Assume that the conditions of Theorem 1 are satisfied, i.e.

$$Eg(An + B) \equiv D \pmod{n} \quad \text{for all } n \in \mathbb{N}, n > N,$$

where $g \in \mathcal{M}$, $A, B, N \in \mathbb{N}$ and $D \neq 0$ is a nonzero integer.

Let $d = (A, B)$, $A = da$, $B = db$, $(a, b) = 1$. Then we have

$$(4) \quad Eg[d(an + b)] \equiv D \pmod{n} \quad \text{for all } n \in \mathbb{N}, n > N.$$

First we prove that

$$g(d) \neq 0, \quad G \in \mathcal{M}_a^* \quad \text{and} \quad Eg(B) = D,$$

where

$$G(n) := \frac{g(dn)}{g(d)} \quad \text{for all } n \in \mathbb{N}.$$

Since $(a, b) = 1$, there are infinitely many $n \in \mathbb{N}$ such that $n > N$ and $(an + b, d) = 1$. For these n , the relation (4) gives

$$Eg(d)g(an + b) \equiv D \pmod{n},$$

which with $D \neq 0$ shows that $g(d) \neq 0$.

Assume that k and l are fixed positive integers, for which $(kl, a) = 1$. Let q be a prime for which

$$q > \max\{k, l, N, |B|, |D|, |E|, |Eg(kB)g(dl) - Dg(dkl)|\}.$$

Since $(kl, qa) = 1$, $(a, b) = 1$ and $q > |B|$, one can deduce from the Chinese Remainder Theorem that there are positive integers x , u , y and v such that

$$kx = aqy + 1, \quad (x, klbd) = 1$$

and

$$lu = aqv + b, \quad (u, kldx) = 1.$$

Then by (4), we have

$$klxu = aqT + b,$$

where $T := by + v + aqyv$. These with (4) and the multiplicativity of g imply that

$$Eg(kB)g(x) = Eg(kBx) = Eg(d(aqby + b)) \equiv D \pmod{q},$$

$$Eg(dl)g(u) = Eg(dlu) = Eg(d(aqv + b)) \equiv D \pmod{q}$$

and

$$Eg(dkl)g(x)g(u) = Eg(d(klxu)) = Eg(d(aqT + b)) \equiv D \pmod{q}.$$

Hence $q > |D|$ implies that $g(x)g(u) \not\equiv 0 \pmod{q}$, consequently

$$Eg(kB)g(dl) \equiv Dg(dkl) \pmod{q}.$$

This relation and the fact $q > |Dg(dkl) - Eg(kB)g(dl)|$ imply that

$$Dg(dkl) = Eg(kB)g(dl)$$

holds for all $k, l \in \mathbb{N}$, $(kl, a) = 1$.

By applying this relation with $l = 1$, we have $Eg(kB)g(d) = Dg(dk)$ for all $k \in \mathbb{N}$, $(k, a) = 1$, therefore we obtain

$$Eg(B) = D \quad \text{and} \quad g(dkl) = \frac{Eg(kB)g(dl)}{D} = \frac{Dg(dk)g(dl)}{Dg(d)} = \frac{g(dk)g(dl)}{g(d)}.$$

Consequently

$$G(kl) = \frac{g(dkl)}{g(d)} = \frac{g(dk)g(dl)}{g(d)g(d)} = G(k)G(l), \quad G \in \mathcal{M}_a^*.$$

Hence we infer from (4) and the fact $(a, b) = 1$ that

$$Eg[d(an + b)] = Eg(d)G(an + b) \equiv D \pmod{n} \quad \text{for all } n \in \mathbb{N}, n > N$$

which, using the fact $G \in \mathcal{M}_a^*$, $Eg(d)G(b) = Eg(db) = Eg(B) = D$, gives

$$(5) \quad Eg(B)G(an + 1) = DG(An + 1) \equiv D \pmod{n} \quad \text{for all } n \in \mathbb{N}, n > N.$$

If $G(aM + 1) = 0$ for some $M \in \mathbb{N}$, then we get from (5) that

$$0 = DG((aM + 1)^t) \equiv D \pmod{\frac{(aM + 1)^t - 1}{a}}$$

for all $t \in \mathbb{N}$. This is impossible because

$$\frac{(aM + 1)^t - 1}{a} \rightarrow \infty \quad \text{as } t \rightarrow \infty.$$

Consequently

$$G(an + 1) \neq 0 \quad \text{for all } n \in \mathbb{N}.$$

On the other hand, if $G(a\ell + 1) = -1$ for some $\ell \in \mathbb{N}$, then from (5) we have

$$-D = DG[(a\ell + 1)^{2t+1}] \equiv D \pmod{\frac{(a\ell + 1)^{2t+1} - 1}{a}},$$

which is impossible. If $G(an + 1) = 1$ for all $n \in \mathbb{N}$, then $G(n) = \chi_a(n)$ for all $n \in \mathbb{N}_a$ and so $g(dn) = g(d)G(n) = g(d)\chi_a(n)$. Theorem 1 is proved in this case with $\alpha = 0$.

In the following we assume that $G(an + 1) \notin \{0, -1, 1\}$ for all $n \in \mathbb{N}$. Let $M = am + 1 \in \mathbb{N}$, $N = an + 1 \in \mathbb{N}$. Then we infer from (5) that

$$DG(N)G(M)^{2t} = DG(NM^{2t}) \equiv D \pmod{\frac{NM^{2t} - 1}{a}},$$

and so

$$(NM^{2t} - 1) \mid aD(G(N)G(M)^{2t} - 1)$$

hold for all $t \in \mathbb{N}$.

To complete the proof of Theorem 1, we shall use the following result of [5] (see Lemma in [5])

Lemma 1. *Let $U \geq 1$, V , $u \geq 1$, v , $\beta > 1$, $\gamma > 1$, $k \geq 1$, l and $F \neq 0$ be fixed integers. If*

$$(U\gamma^{kn+l} + V) \mid F(u\beta^{kn+l} + v)$$

for all $n \in \mathbb{N}$, then there is a positive integer e such that

$$\beta = \gamma^e \quad \text{and} \quad u(-V)^e + vU^e = 0.$$

Since $G(M)^2 > 1$, therefore Lemma 1 shows that there is a non-negative integer α such that

$$G(M)^2 = M^{2\alpha}$$

and

$$G(N) - N^\alpha = G(an + 1) - (an + 1)^\alpha = 0.$$

Let

$$\mathcal{G}(n) := \frac{G(n)}{n^\alpha}, \quad G(n) = n^\alpha \mathcal{G}(n) \quad (n \in \mathbb{N}).$$

Then we have

$$\mathcal{G}(an + 1) = 1 \quad \text{for all } n \in \mathbb{N},$$

which gives that $\mathcal{G}(n) = \chi_a(n)$ for all $n \in \mathbb{N}_a$, where χ_a is a real-valued Dirichlet character (mod a). Therefore

$$g(dn) = g(d)G(n) = n^\alpha g(d)\mathcal{G}(n) = n^\alpha g(d)\chi_a(n) \quad \text{for all } n \in \mathbb{N}_a.$$

The proof of Theorem 1 is finished.

3. Proof of (I) of Theorem 2

First we prove the following

Lemma 2. *Assume that $a, b \in \mathbb{N}$, $c, d \in \mathbb{Z}$, $c \neq 0$ and $H \in \mathcal{M}^*$ satisfy the condition*

$$(6) \quad cH(an + b) + d \equiv 0 \pmod{an + b} \quad \text{for all } n \in \mathbb{N}.$$

Then we have:

If $d \neq 0$, then $H(n) = \chi_a(n)$ for all $n \in \mathbb{N}_a$.

If $d = 0$, then either $H(an + b) = 0$ for all $n \in \mathbb{N}$ or $H(n) \equiv 0 \pmod{n}$ for all $n \in \mathbb{N}_a$.

Proof. Assume that (6) holds for all $n \in \mathbb{N}$.

First we consider the case when $d \neq 0$. In this case, by (6) we have

$$H(b) \neq 0 \quad \text{and} \quad cH(b)H(an + 1) \equiv -d \pmod{an + 1}.$$

Then, for each $N \in \mathbb{N}$ we have

$$\begin{aligned} -dH(aN + 1) &= cH(b)H(aN + 1)H(an + 1) = \\ &= cH(b)H\left[(aN + 1)(an + 1)\right] \equiv -d \pmod{an + 1}, \end{aligned}$$

consequently

$$-dH(aN + 1) = -d, \quad H(aN + 1) = 1$$

hold for each $N \in \mathbb{N}$. This shows that $H(n) = \chi_a(n)$ for all $n \in \mathbb{N}_a$.

Assume now that $d = 0$. In this case, by (6) we have

$$cH(an + b) \equiv 0 \pmod{an + b} \quad \text{for all } n \in \mathbb{N}.$$

We have two possibilities: either $H(an + b) = 0$ for all $n \in \mathbb{N}$ or $H(aN + b) \neq 0$ for some $N \in \mathbb{N}$. Assume that $H(aN + b) \neq 0$. Then

$$cH(aN + b)H(an + 1) = cH\left[(aN + b)(an + 1)\right] \equiv 0 \pmod{an + 1}$$

for all $n \in \mathbb{N}$. Thus, for each prime p , $(p, a) = 1$, we have

$$cH(aN + b)H(p)^{\varphi(a)t} = cH(aN + b)H(p^{\varphi(a)t}) \equiv 0 \pmod{p^{\varphi(a)t}} \quad \text{for all } t \in \mathbb{N}.$$

This with $cH(aN + b) \neq 0$ shows that $p \mid H(p)$. Lemma 2 is proved.

Now assume that the integers $A > 0$, $B > 0$, $(A, B) = 1$, $C \neq 0$, $D \neq 0$, $E \neq 0$ and $f, g \in \mathcal{M}^*$ satisfy the relation

$$(7) \quad Eg(An + B) \equiv Cf(n) + D \pmod{n} \quad \text{for all } n \in \mathbb{N}.$$

We shall prove the following lemma.

Lemma 3. *Assume that there is a prime π such that $f(\pi) = 0$. Then $Eg(B) = D$ and the following assertions hold:*

(a) *There are a non-negative integer α and a character $\chi_A \pmod{A}$ such that*

$$n|f(n) \quad \text{and} \quad g(m) = \chi_A(m)m^\alpha \quad (n \in \mathbb{N}, m \in \mathbb{N}_A).$$

(b) *If $\pi|A$ and $(\pi, B) = 1$, then $C = -2D$ and all further solutions (f, g) of (7) have the form*

$$\pi = 2, \quad f(n) = \chi_2(n) \quad \text{and} \quad g(m) = \chi_{2A}(m)m^\alpha \quad (n \in \mathbb{N}, m \in \mathbb{N}_A),$$

where α is a non-negative integer and χ_{2A} is a character $\pmod{2A}$ with the condition $\chi_{2A}(A+B) = -\chi_{2A}(B)$.

(c) *If $(\pi, AB) = 1$, then $C = -D$ and all further solutions (f, g) of (7) have the form*

$$\pi = 2, \quad f(n) = \chi_2(n), \quad g(2) = 0 \quad \text{and} \quad g(m) = \chi_{2A}(m)m^\alpha, \quad (n \in \mathbb{N}, m \in \mathbb{N}_A),$$

where α is a non-negative integer and χ_{2A} is a character $\pmod{2A}$.

Proof. Assume that there is a prime π such that $f(\pi) = 0$. Then, by writing $n\pi$ in the place of n in (7), we have

$$Eg(A\pi n + B) \equiv Cf(\pi n) + D = Cf(\pi)f(n) + D = D \pmod{n}$$

for all $n \in \mathbb{N}$. This, by using Theorem 1, implies that there are a non-negative integer α and a real-valued Dirichlet character $\chi_{\pi A} \pmod{\pi A}$ such that

$$g(n) = \chi_{\pi A}(n)n^\alpha \quad \text{holds for all } n \in \mathbb{N}_{\pi A}.$$

In the following let

$$G(n) := \frac{g(n)}{n^\alpha} \quad \text{for all } n \in \mathbb{N}.$$

Then

$$(8) \quad g(n) = n^\alpha G(n) \quad \text{and} \quad G(m) = \chi_{\pi A}(m) \quad \text{for all } n \in \mathbb{N}, m \in \mathbb{N}_{\pi A},$$

and we infer from (7) that

$$\begin{aligned} EB^\alpha G(An + B) &\equiv E(An + B)^\alpha G(An + B) = Eg(An + B) \equiv \\ &\equiv Cf(n) + D \pmod{n}. \end{aligned}$$

This gives

$$(9) \quad EB^\alpha G(An + B) \equiv Cf(n) + D \pmod{n} \text{ for all } n \in \mathbb{N}.$$

Next we prove that

$$(10) \quad Eg(B) = D.$$

Indeed, by applying (9) with $n = B\pi m$, we obtain from (8)

$$\begin{aligned} Eg(B) &\equiv Eg(B)G(A\pi m + 1) = EB^\alpha G(B)G(A\pi m + 1) = \\ &= EB^\alpha G(AB\pi m + B) \equiv Cf(B\pi m) + D = D \pmod{m}, \end{aligned}$$

which proves (10).

We shall use the notation $\varphi_\alpha(n) := n^\alpha$, $\alpha \in \mathbb{N}$. Furthermore let $\mathcal{D} \in \mathcal{M}^*$ such that $n|\mathcal{D}(n)$ for all $n \in \mathbb{N}$. It is obvious that $\mathcal{D}(n) = n\mathcal{D}_1(n)$ and $\mathcal{D}_1 \in \mathcal{M}^*$.

We shall prove that the solution (f, g) of (7) is $(\mathcal{D}, \chi_A \varphi_\alpha)$ in the following cases:

- (i) $f(p) = 0$ for some prime p , $p \neq \pi$,
- (ii) $f(B) = 0$.

Case (i): Assume that there is a prime p , $p \neq \pi$ for which $f(p) = 0$. Then, as we have seen in the proof of (8), we have

$$g(n) = n^\alpha G(n) \text{ and } G(m) = \chi_{pA}(m) \text{ for all } n \in \mathbb{N}, m \in \mathbb{N}_{pA},$$

which imply that $G(n) = \chi_A(n)$ for all $n \in \mathbb{N}_A$. In this case we infer from (7) and (10) that

$$\begin{aligned} Cf(n) &\equiv Eg(An + B) - D = E(An + B)^\alpha G(An + B) - D \equiv \\ &\equiv EB^\alpha G(B) - D = Eg(B) - D = 0 \pmod{n}, \end{aligned}$$

which by using Lemma 2 gives $n|f(n)$ for all $n \in \mathbb{N}$. Thus the solution (f, g) of (7) is $(\mathcal{D}, \chi_A \varphi_\alpha)$ in the case (i).

In the following we assume that

$$(11) \quad f(n) \neq 0 \text{ if and only if } (n, \pi) = 1.$$

Case (ii): Assume that $f(B) = 0$. Then by writing Bn in the place of n in (9), we have

$$\begin{aligned} Eg(B)G(An + 1) &= EB^\alpha G(B)G(An + 1) = \\ &= EB^\alpha G(ABn + B) \equiv Cf(B)f(n) + D \equiv D \pmod{n} \end{aligned}$$

for all $n \in \mathbb{N}$. This relation with Theorem 1 implies that $G(n) = \chi_A(n)$ for all $n \in \mathbb{N}_A$. Thus we get from (10) that $EB^\alpha G(An + B) = EB^\alpha G(B) = Eg(B) = D$, consequently from (9) that $Cf(n) \equiv 0 \pmod{n}$ for all $n \in \mathbb{N}$. Hence Lemma 2 gives $n \mid f(n)$ for all $n \in \mathbb{N}$ and so the solution of (7) is $(\mathcal{D}, \chi_A \varphi_\alpha)$ for the case (ii).

In the following we assume that

$$(12) \quad f(B) \neq 0 \quad \text{and} \quad (\pi, B) = 1.$$

For each $\ell \geq 0$ let

$$\mathcal{H}_\ell := \{ n \in \mathbb{N} \mid \pi^\ell \parallel An + B \}.$$

We note that

$$\mathbb{N} = \begin{cases} \mathcal{H}_0 & \text{if } \pi \mid A, \\ \bigcup_{\ell=0}^{\infty} \mathcal{H}_\ell & \text{if } (\pi, A) = 1. \end{cases}$$

Let $n_\ell \in \mathcal{H}_\ell$ and $An_\ell + B = \pi^\ell N_\ell$. It is clear to see from $(A\pi, B) = 1$ that $(n_\ell, A\pi) = (N_\ell, A\pi) = 1$. By writing $\pi^{\ell+1}m + n_\ell$ in place of n in (10), we infer from (8) that

$$\begin{aligned} EB^\alpha G(An_\ell + B) &= EB^\alpha G(\pi^\ell)G(N_\ell) = EB^\alpha G(\pi^\ell)G(A\pi m + N_\ell) = \\ &= EB^\alpha G\left(A(\pi^{\ell+1}m + n_\ell) + B\right) \equiv \\ &\equiv Cf(\pi^{\ell+1}m + n_\ell) + D \pmod{\pi^{\ell+1}m + n_\ell} \end{aligned}$$

and

$$(13) \quad Cf(\pi^{\ell+1}m + n_\ell) \equiv EB^\alpha G(An_\ell + B) - D := K_\ell \pmod{\pi^{\ell+1}m + n_\ell}$$

hold for all $m \in \mathbb{N}$.

We shall consider (13) in two cases, according to $K_\ell = 0$ or $K_\ell \neq 0$.

Case I. $K_\ell = 0$ for some $\ell \in \mathbb{N}$, $0 \leq \ell < \pi$.

In this case, we infer from (11) and the fact $(n_\ell, \pi) = 1$ that $f(\pi^{\ell+1}m + n_\ell) \neq 0$, consequently we obtain from Lemma 2 that $n \mid f(n)$ for all $n \in \mathbb{N}_\pi$. Hence by using the fact $f(\pi) = 0$, we get from (7) that

$$n \mid f(n) \quad \text{and} \quad Eg(An + B) \equiv Cf(n) + D \equiv D \pmod{n} \quad \text{for all } n \in \mathbb{N},$$

which gives that $g(n) = n^\alpha \chi_A(n)$ for all $(n \in \mathbb{N}_A)$, where $\alpha \in \mathbb{N}$ and χ_A is a real-valued Dirichlet character $(\bmod A)$.

Case II. $K_\ell \neq 0$ for $\ell = 0, 1, \dots, \pi - 1$.

In this case, by applying (13) for $\ell = 0$, Lemma 2 gives

$$f(n) = \chi_\pi(n) \quad \text{for all } n \in \mathbb{N}_\pi.$$

Since $f(\pi) = 0$, we have

$$(14) \quad f(n) = \chi_\pi(n) \quad \text{for all } n \in \mathbb{N}.$$

Thus, from (13) and (14) we get

$$EB^\alpha G(An_\ell + B) \equiv Cf(\pi^{\ell+1}m + n_\ell) + D = Cf(n_\ell) + D \pmod{\pi^{\ell+1}m + n_\ell}$$

and so

$$EB^\alpha G(An_\ell + B) = Cf(n_\ell) + D$$

hold for all $m \in \mathbb{N}$, $\ell \in \{0, 1, \dots, \pi - 1\}$ and $n_\ell \in \mathcal{H}_\ell$. Consequently, we have have

$$(15) \quad EB^\alpha G(An + B) = Cf(n) + D \quad \text{for all } n \in \mathbb{N}.$$

From (10) we have $EB^\alpha G(B) = Eg(B) = D$, and so we get from (15) that

$$(16) \quad DG(An + 1) = Cf(B)f(n) + D \quad \text{for all } n \in \mathbb{N}.$$

We shall deduce from (16) that

$$(17) \quad \pi = 2.$$

Assume that $\pi \geq 3$. Then there is a $\nu \in \mathbb{N}$ such that $(A\nu + 1, A\pi) = (\nu, \pi) = 1$. By (8) and (14) we have $f(\nu) = \pm 1$ and $G(A\nu + 1) = \pm 1$, consequently we infer from (16) that

$$\begin{aligned} D^2 &= [DG(A\nu + 1)]^2 = (Cf(B)f(\nu) + D)^2 = \\ &= C^2f(B)^2f(\nu)^2 + 2CDf(B)f(\nu) + D^2 = \\ &= C^2f(B)^2 + 2Cf(B)Df(\nu) + D^2, \end{aligned}$$

which implies $2Df(\nu) + Cf(B) = 0$.

Therefore from (16) we have

$$(18) \quad G(An + 1) = -2f(\nu)f(n) + 1 \quad \text{for all } n \in \mathbb{N}.$$

If $f(\nu) = -1$, then (18) gives that $G(An + 1) = 2f(n) + 1$ for all $n \in \mathbb{N}$. Hence we have $G(A+1) = 2f(1)+1 = 3$ and $9 = G(A+1)^2 = G[A(A+2)+1] = 2f(A+2) + 1$, which imply $f(A+2) = \chi_\pi(A+2) = 4$. This is impossible.

Thus, we have $f(\nu) = 1$ and so

$$G(An + 1) = -2f(n) + 1 \quad \text{for all } n \in \mathbb{N}.$$

Since

$$G(An + 1)^2 = G[An(An + 2) + 1] = -2f(n)f(An + 2) + 1$$

and

$$G(An + 1)^2 = [-2f(n) + 1]^2 = 4f(n)^2 - 4f(n) + 1,$$

we have

$$f(n)[f(An + 2) + 2f(n) - 2] = 0 \quad \text{for all } n \in \mathbb{N}.$$

This relation with the fact $f(n) = \chi_\pi(n)$ implies that

$$f(n) = 1 \quad \text{and} \quad f(An + 2) = 0 \quad \text{for all } n \in \mathbb{N}_\pi,$$

from which we obtain that

$$\pi | An + 2 \quad \text{for all } n \in \mathbb{N}_\pi.$$

Since $1 \in \mathbb{N}_\pi$ and $\pi - 1 \in \mathbb{N}_\pi$, we get from the last relation that $\pi | A + 2$, $\pi | A(\pi - 1) + 2 = A\pi - (A - 2)$. These imply $\pi = 2$.

Thus (17) is proved.

Now we prove that $G(A + B) = -G(B)$ in the case $2|A$.

Since $f(n) = \chi_\pi(n) = \chi_2(n)$ for all $n \in \mathbb{N}_2$ and $(B, 2) = 1$, we get from (16) that $DG(A + 1) = Cf(B) + D = C + D$. It follows from $2|A$ that $(A + 1, 2A) = 1$, which implies from (8) that $G(A + 1) = -1$ and $C = -2D$. Therefore (10) and (15) imply that

$$EB^\alpha G(A + B) = C + D = -2D + D = -D = -Eg(B) = -EB^\alpha G(B),$$

consequently

$$G(A + B) = -G(B).$$

The part (b) of Lemma 3 is proved.

Assume that $(2, AB) = 1$. Then for every $\nu \geq 1$ there are $n_\nu \in \mathbb{N}$ and $N_\nu \in \mathbb{N}$ such that $An_\nu + 1 = 2^\nu N_\nu$, $(N_\nu, 2) = 1$. It is obvious that

$$(n_\nu, 2) = (N_\nu, 2A) = 1, f(n_\nu) = \chi_2(n_\nu) = 1 \quad \text{and} \quad G(N_\nu) = \chi_{2A}(N_\nu) = \pm 1.$$

We infer from (16) that

$$DG(An_\nu + 1) = DG(2^\nu)G(N_\nu) = Cf(B)f(n_\nu) + D = C + D$$

holds for all $\nu \geq 1$, from which we obtain that

$$G(2)^\nu G(N_\nu) = G(2)G(N_1) = C + D \quad \text{for all } \nu \in \mathbb{N}, \nu \geq 1.$$

This shows that $G(2) \in \{0, 1, -1\}$. We shall prove that $G(2) = 0$. Assume that $G(2) = \pm 1$. Then $G(An + 1) = \pm 1$, and so we get from (16) that $DG(A + 1) = Cf(B)f(1) + D = C + D$. Consequently $G(A + 1) = -1$ and

$$D = DG(A + 1)^2 = DG[A(A + 2) + 1] = Cf(B)f(A + 2) + D = C + D,$$

which is impossible. Thus we have proved that $G(2) = 0$. Hence $g(2) = n^\alpha G(2) = 0$ and $0 = EB^\alpha G(A + B) = Cf(B) + D = C + D$, i.e. $C = -D$.

Lemma 3 and the part (I) of Theorem 2 are proved.

4. Proof of (II) of Theorem 2

We shall prove the part (II) of Theorem 2 by showing

Lemma 4. *Assume that all conditions of Theorem 2 are satisfied and $f(n) \neq 0$ for all $n \in \mathbb{N}$, $g(AN + B) = 0$ for some $N \in \mathbb{N}$. Then*

$$f(n) = 1 \quad \text{for all } n \in \mathbb{N},$$

and either

$$g(An + B) = 0 \quad \text{for all } n \in \mathbb{N} \quad \text{if } C + D = 0,$$

or there are a non-negative integer α and a real-valued character χ_A such that

$$g(n) = \chi_A(n)n^\alpha \quad \text{for all } n \in \mathbb{N}_A \quad \text{if } C + D \neq 0.$$

Proof. We infer from $g(AN + B) = 0$ that there is a prime p such that $p|AN + B$, $g(p) = 0$ and $(p, A) = 1$. By writing $n = pm + N$ in the place of n in

(7), we have $p|An+B = Apm+AN+B$, $g(An+B) = 0$ and $Cf(pm+N)+D \equiv \equiv 0 \pmod{pm+N}$. This with Lemma 2 implies that

$$(19) \quad f(n) = \chi_p(n) \text{ for all } n \in \mathbb{N}_p.$$

We shall prove that

$$(20) \quad f(n) = 1 \text{ for all } n \in \mathbb{N}$$

By writing nB in the place of n in (7), we have

$$(21) \quad Eg(B)g(An+1) \equiv Cf(B)f(n) + D \pmod{n} \text{ for all } n \in \mathbb{N}.$$

It is obvious that if $g(B) = 0$, then (21) gives $Cf(B)f(n) + D \equiv 0 \pmod{n}$, and so we get from Lemma 2 that $f(n) = 1$ for all $n \in \mathbb{N}$. Thus (20) is true in the case $g(B) = 0$.

In the following we assume that

$$(22) \quad g(B) \neq 0, (p, B) = 1.$$

For every $M \in \mathbb{N}$, we have

$$\begin{aligned} Eg(AMn+B)g(An+1) &= Eg[An(AMn+B+M)+B] \equiv \\ &\equiv Cf(AMn+B+M)f(n) + D \pmod{n} \end{aligned}$$

and

$$\begin{aligned} &(Cf(M)f(n) + D)(Cf(B)f(n) + D) = \\ &= C^2f(B)f(M)f(n)^2 + CD(f(B) + f(M))f(n) + D^2, \end{aligned}$$

therefore we get from (7) and (21) that

$$(23) \quad \begin{aligned} &CEg(B)f(AMn+B+M)f(n) \equiv \\ &\equiv C^2f(B)f(M)f(n)^2 + CD(f(B) + f(M))f(n) + D^2 - DEg(B) \pmod{n}. \end{aligned}$$

By applying $M = pm$, ($m \in \mathbb{N}$) in (23), using (19), we get

$$f(Apmn+B+pm) = \chi_p(Apmn+B+pm) = \chi_p(B) = f(B)$$

and

$$(24) \quad CEg(B)f(B)f(n) \equiv$$

$$\equiv C^2 f(B) f(pm) f(n)^2 + CD \left(f(B) + f(pm) \right) f(n) + D^2 - DEg(B) \pmod{n}.$$

Now, let us write $n(pt+1)$ in the place of n in (24), for every t . From (19) we get $f(pt+1) = 1$ and that

$$\begin{aligned} & CEg(B) f(B) f(n) \equiv \\ & \equiv C^2 f(B) f(pm) f(n)^2 + CD \left(f(B) + f(pm) \right) f(n) + D^2 - DEg(B) \pmod{pt+1} \end{aligned}$$

hold for all $n, m, t \in \mathbb{N}$, which gives that

$$\begin{aligned} (25) \quad & C \left[Eg(B) f(B) - D(f(B) + f(pm)) \right] f(n) = \\ & = C^2 f(B) f(pm) f(n)^2 + D^2 - DEg(B) \end{aligned}$$

hold for all $n, m \in \mathbb{N}$.

If there is an $m \in \mathbb{N}$ such that $Eg(B) f(B) - D(f(B) + f(pm)) \neq 0$, then we apply (25) for the case $n \in \mathbb{N}_p$, we have $f(n)^2 = 1$ and

$$f(n) = \frac{C^2 f(B) f(pm) + D^2 - DEg(B)}{CEg(B) f(B) - CD(f(B) + f(pm))},$$

consequently $f(n) = 1$ for all $n \in \mathbb{N}_p$. We apply (25) again for the case when $n = p^\nu$, $n \in \mathbb{N}$. We have $f(p) = \pm 1$ and

$$f(p)^\nu = \frac{C^2 f(B) f(pm) + D^2 - DEg(B)}{CEg(B) f(B) - CD(f(B) + f(pm))}.$$

This shows that $f(p) = 1$. Thus (20) is proved in this case.

If $Eg(B) f(B) = D(f(B) + f(pm))$ for all $m \in \mathbb{N}$, then

$$f(m) = \frac{Eg(B) f(B) - Df(B)}{Df(p)} \quad \text{for all } m \in \mathbb{N},$$

which implies $f(m) = 1$ for all $m \in \mathbb{N}$.

Finally, we infer from (7) and (20) that

$$Eg(An + B) \equiv Cf(n) + D = C + D \pmod{n}.$$

If $C + D \neq 0$, then Theorem 1 implies that there are a non-negative integer α and a real-valued Dirichlet character χ_A such that

$$g(n) = \chi_A(n) n^\alpha \quad \text{for all } n \in \mathbb{N}_A.$$

If $C + D = 0$, then $Eg(An + B) \equiv 0 \pmod{n}$. Assume that $g(A\ell + B) \neq 0$ for some $\ell \in \mathbb{N}$. Let $q \in \mathcal{P}$, $(q, A) = 1$. Then $(A\ell + B)q^{\varphi(A)t} \equiv B \pmod{A}$ and

$$g\left((A\ell + B)q^{\varphi(A)t}\right) = g(A\ell + B)g(q)^{\varphi(A)t} \equiv 0 \pmod{\frac{(A\ell + B)q^{\varphi(A)t} - B}{A}}$$

for all $t \in \mathbb{N}$. This implies that $P(t) \mid g(A\ell + B)g(q)$, where

$$P(t) = \text{the greatest prime factor of } \frac{(A\ell + B)q^{\varphi(A)t} - B}{A}.$$

This is impossible, because well-known that $P(t)$ cannot be bounded.

Lemma 4 and the part (II) of Theorem 2 is proved.

5. Proof of (III) of Theorem 2

Assume that all conditions of Theorem 2 are satisfied and

$$(26) \quad g(An + B)f(n) \neq 0 \text{ for all } n \in \mathbb{N}.$$

From (7) we have

$$Eg(B)g(An + 1) \equiv Cf(B)f(n) + D \pmod{n} \text{ for all } n \in \mathbb{N},$$

and so

$$(27) \quad \mathcal{E}g(An + 1) \equiv Cf(n) + D \pmod{n} \text{ for all } n \in \mathbb{N},$$

where $\mathcal{E} := Eg(B) \neq 0$, $\mathcal{C} := Cf(B) \neq 0$.

First we infer from (27) that

$$\mathcal{E}^2g(ANn + 1)g(AMn + 1) \equiv (\mathcal{C}f(N)f(n) + D)(\mathcal{C}f(M)f(n) + D) \pmod{n}$$

and

$$\begin{aligned} \mathcal{E}^2g(ANn + 1)g(AMn + 1) &= \mathcal{E}^2g\left[An(ANMn + N + M) + 1\right] \equiv \\ &\equiv \mathcal{C}\mathcal{E}f(ANMn + N + M)f(n) + D\mathcal{E} \pmod{n} \end{aligned}$$

are satisfied for all $n, N, M \in \mathbb{N}$. Consequently

$$\begin{aligned} & \mathcal{CE}f(ANMn + N + M)f(n) \equiv \\ & \equiv \mathcal{C}^2 f(N)f(M)f(n)^2 + \mathcal{CD}\left(f(N) + f(M)\right)f(n) + D^2 - D\mathcal{E} \pmod{n} \end{aligned}$$

holds for all $n, N, M \in \mathbb{N}$. By writing $n(N + M)$ in the place of n in the above congruence, we have

$$(28) \quad \begin{aligned} & a(N, M)f(ANMn + 1)f(n) \equiv \\ & \equiv b(N, M)f(n)^2 + c(N, M)f(n) + d \pmod{n}, \end{aligned}$$

where

$$a(N, M) := \mathcal{CE}f(N + M)^2, \quad b(N, M) := \mathcal{C}^2 f(N)f(M)f(N + M)^2,$$

and

$$c(N, M) := \mathcal{CD}\left(f(N) + f(M)\right)f(N + M), \quad d := D^2 - D\mathcal{E}.$$

By applying (28) with $N = M = 1$, we have

$$a(1, 1)f(An + 1)f(n) \equiv b(1, 1)f(n)^2 + c(1, 1)f(n) + d \pmod{n},$$

and if we substitute n by NMn , then

$$\begin{aligned} & a(1, 1)f(NM)f(ANMn + 1)f(n) \equiv \\ & \equiv b(1, 1)f(NM)^2 f(n)^2 + c(1, 1)f(NM)f(n) + d \pmod{n}. \end{aligned}$$

Hence, this congruence with (28) implies

$$(29) \quad \begin{aligned} & f(2)^2 f(NM)[b(N, M)f(n)^2 + c(N, M)f(n) + d] \equiv \\ & \equiv f(N + M)^2 [b(1, 1)f(NM)^2 f(n)^2 + c(1, 1)f(NM)f(n) + d] \pmod{n}. \end{aligned}$$

Let

$$\begin{aligned} & \lambda(N, M) := f(2)^2 f(NM)c(N, M) - f(N + M)^2 c(1, 1)f(NM) = \\ & = \mathcal{CD}f(N + M)f(2)f(NM) \left[f(2)\left(f(N) + f(M)\right) - 2f(N + M) \right] \end{aligned}$$

and

$$d(N, M) := d[f(2)^2 f(NM) - f(N + M)^2] =$$

$$= (D^2 - D\mathcal{E})[f(2)^2f(NM) - f(N + M)^2].$$

Since

$$f(2)^2f(NM)b(N, M) - f(N + M)^2b(1, 1)f(NM)^2 = 0$$

hold for all $N, M \in \mathbb{N}$, we infer from (29) that

$$(30) \quad \lambda(N, M)f(n) + d(N, M) \equiv 0 \pmod{n}.$$

Next we prove that (30) with (26) implies that

$$(31) \quad \text{either } f(n) = 1 \text{ or } n|f(n) \text{ for all } n \in \mathbb{N}.$$

We separate the cases $\lambda(N, M) = 0$ for every N, M , and $\lambda(N, M) \neq 0$ for some $N, N \in \mathbb{N}$.

Case A. $\lambda(N, M) = 0$ for all $N, M \in \mathbb{N}$

In this case (26) and (30) imply that

$$(32) \quad f(2)\left(f(N) + f(M)\right) - 2f(N + M) = 0$$

and

$$(33) \quad d(N, M) = (D^2 - D\mathcal{E})[f(2)^2f(NM) - f(N + M)^2] = 0$$

for all $N, M \in \mathbb{N}$.

By (26) we have $f(2) \neq 0$. Then by using (32), we have $2f(3) = 2f(2+1) = f(2)^2 + f(2)$ and $2f(2)^2 = 2f(3+1) = f(2)f(3) + f(2)$. These with $f(2) \neq 0$ imply that either $f(2) = 1$ or $f(2) = 2$. We apply (32) for the case when $N = 1$ and $M = m, m \in \mathbb{N}$. We have

$$2f(m + 1) = f(2)f(m) + f(2) \text{ for all } m \in \mathbb{N},$$

from which we obtain that $f(n) = 1$ in the case $f(2) = 1$ and $f(n) = n$ in the case $f(2) = 2$.

Case B. $\lambda(N, M) \neq 0$ for some $N, M \in \mathbb{N}$

In this case, if $d(N, M) \neq 0$, then we get from Lemma 2 that $f(n) = 1$ for all $n \in \mathbb{N}$. But this is impossible, because

$$d(N, M) = (D^2 - D\mathcal{E})[f(2)^2f(NM) - f(N + M)^2] \neq 0.$$

Thus, in this case we have $d(N, M) = 0$, which by using (26) and Lemma 2 gives that $n|f(n)$ for all $n \in \mathbb{N}$. Consequently we proved (31) for all cases.

In the following we assume that (7), (26) and (31) hold.

First we assume that $f(n) = 1$ for all $n \in \mathbb{N}$. Then we infer from (7) that

$$(34) \quad Eg(An + B) \equiv Cf(n) + D = C + D \pmod{n} \text{ for all } n \in \mathbb{N}.$$

We prove that in this case $C + D = \mathcal{E} = Eg(B)$ and $g(n) = n^\alpha \chi_A(n)$ is satisfied for all $n \in \mathbb{N}_A$.

Suppose that $C + D = 0$. Then $Eg(An + B) \equiv 0 \pmod{n}$ for all n . By (26) we have $g(AN + B) \neq 0$ is true for all $N \in \mathbb{N}$. Then $(AN + B)^{\varphi(A)t+1} \equiv B \pmod{A}$ and

$$\begin{aligned} & Eg(AN + B)^{\varphi(A)t+1} = \\ & = Eg[(AN + B)^{\varphi(A)t+1}] \equiv 0 \pmod{\frac{(AN + B)^{\varphi(A)t+1} - B}{A}} \end{aligned}$$

hold for all $t \in \mathbb{N}$. This is impossible, because well-known that $P(t)$ cannot be bounded, where

$$P(t) := \text{the largest prime divisor of } \frac{(AN + B)^{\varphi(A)t+1} - B}{A}.$$

Thus, we proved that in the case $f(n) = 1$ for all n , we have $C + D \neq 0$. This with (34) shows that $g(n) = n^\alpha \chi_A(n)$ for all $n \in \mathbb{N}_A$, where $\alpha \geq 0$ is an integer. Finally we infer from (34) that

$$\begin{aligned} C + D &\equiv Eg(An + B) = E(An + B)^\alpha \chi_A(An + B) \equiv \\ &\equiv EB^\alpha \chi_A(B) = Eg(B) \pmod{n}, \end{aligned}$$

and so $C + D = Eg(B)$.

Now assume that $n|f(n)$ for all $n \in \mathbb{N}$. In this case we get from (7), (33) and Theorem 1 that $D = \mathcal{E} = Eg(B)$ and $g(n) = n^\alpha \chi_A(n)$ ($n \in \mathbb{N}_A$) with some a non-negative integer α .

The proof of the part (III) is completed and Theorem 2 is proved.

References

- [1] **Elliott, P.D.T.A.**, *Arithmetic functions and integers products*, Grundle Math. Wiss. **272**, Springer-Verlag, 1985.

- [2] **Iványi, I.**, On multiplicative functions with congruence property, *Annales Univ. Sci. Budapest. Sect. Math.*, **15** (1972), 133-137.
- [3] **Phong, B.M.**, Multiplicative functions satisfying a congruence property V., *Acta Math. Hungar.*, **62** (1993), 81-87.
- [4] **Phong, B.M.**, On arithmetical functions satisfying congruence properties, *Annales Univ. Sci. Budapest. Sect. Comp.*, **22** (2003), 49-67.
- [5] **Phong, B.M.**, On integer-valued arithmetical functions satisfying congruence properties, *Annales Univ. Sci. Budapest. Sect. Comp.*, **27** (2007), 229-244.
- [6] **Phong, B.M. and Fehér J.**, Note on multiplicative functions satisfying congruence property, *Annales Univ. Sci. Budapest. Sect. Math.*, **33** (1990), 261-265.
- [7] **Phong, B.M. and Fehér J.**, Note on multiplicative functions satisfying congruence property II., *Math. Pannon.*, **10** (1) (1999), 133-138.
- [8] **Subbarao, M.V.**, Arithmetic functions satisfying congruence property, *Canad. Math. Bull.*, **9** (1966), 143-146.

Bui Minh Phong

Department of Computer Algebra

Eötvös Loránd University

Pázmány Péter sét. 1/C

H-1117 Budapest, Hungary

bui@compalg.inf.elte.hu