

POWER SERIES – THE STRUCTURE OF H.-O.-GAPS

Wolfgang Luh (Trier, Germany)

*Dedicated to an excellent mathematician
and my real good friend Karl-Heinz Indlekofer*

Communicated by Imre Kátai

(Received December 13, 2012; accepted January 24, 2013)

Abstract. Suppose that a power series $f(z) = \sum_{\nu=0}^{\infty} a_{\nu}(z - z_0)^{\nu}$ with positive radius of convergence has a sequence of H.-O.-gaps. Then there exists a neighborhood $U(z_0)$ of z_0 , such that the rearrangement $f(z) = \sum_{\nu=0}^{\infty} b_{\nu}(z - \zeta)^{\nu}$ also has H.-O.-gaps for all $\zeta \in U(z_0)$.

1. Introduction

Let be given a power series $\sum_{\nu=0}^{\infty} a_{\nu}(z - z_0)^{\nu}$ with radius of convergence R , where $0 < R < \infty$. We say that this series has a sequence $\{p_k, q_k\}$ of Hadamard–Ostrowski-gaps (H.-O.-gaps for short) if the following conditions hold:

- p_k, q_k are natural numbers with $p_1 < q_1 < p_2 < q_2 < \dots$,
- there exists $\lambda > 1$ such that $\frac{q_k}{p_k} > \lambda$ for all $k \in \mathbb{N}$,
- for $I := \bigcup_{k=1}^{\infty} [p_k, q_k]$ we have $\overline{\lim}_{\nu \rightarrow \infty, \nu \in I} |a_{\nu}|^{1/\nu} < \frac{1}{R}$.

Key words and phrases: H.-O.-gaps, overconvergence, rearrangements.
2010 Mathematics Subject Classification: 30B30, 30B40.

In the case that

$$\frac{q_k}{p_k} \rightarrow \infty \text{ and } \lim_{\substack{\nu \rightarrow \infty \\ \nu \in I}} |a_\nu|^{1/\nu} = 0$$

we say that this series has Ostrowski-gaps (O.-gaps for short).

Approximately 90 years ago it was shown by Ostrowski ([3], [4], [5], [6]) that there exists a deep interdependence with the occurrence of those gaps and the phenomenon of overconvergence i. e. the convergence of a subsequence of the partial sums $s_n(z) = \sum_{\nu=0}^n a_\nu(z-z_0)^\nu$ in a bigger domain than the circle of convergence.

Ostrowski's main results on overconvergence are the following.

Theorem O₁ *Suppose that the power series $f(z) = \sum_{\nu=0}^{\infty} a_\nu(z-z_0)^\nu$ with radius of convergence $R, 0 < R < \infty$ possesses H.-O.-gaps $\{p_k, q_k\}$ and let the function f be holomorphic at $z_1, |z_1 - z_0| = R$.*

Then there exists a neighborhood $U(z_1)$ of z_1 such that the sequence $\{s_{p_k}(z)\}$ of partial sums converges compactly on $U(z_1)$.

If the power series has O.-gaps $\{p_k, q_k\}$, then the function f is holomorphic in a simply connected domain $G(f) \supset \{z : |z - z_0| < R\}$ and $\{s_{p_k}(z)\}$ converges compactly on $G(f)$.

Theorem O₂ *Suppose that the power series $f(z) = \sum_{\nu=0}^{\infty} a_\nu(z-z_0)^\nu$ has radius of convergence $R, 0 < R < \infty$ and that $\{s_{p_k}(z)\}$ is an overconvergent subsequence of its partials sums.*

Then the series has a sequence of H.-O.-gaps of the type $\{p_k, q_k\}$.

For a good treatise on the theory of overconvergence we refer to Hille's book [2], section 16.7.

In recent years the investigation of overconvergence has got a revival since its connection with universal properties of power series has been detected. For details we refer to the excellent survey of Große-Erdmann [1], where also a synopsis of the relevant literatur is given.

In this paper we are dealing with a long outstanding problem: If $f(z) = \sum_{\nu=0}^{\infty} a_\nu(z-z_0)^\nu$ has a sequence of H.-O.-gaps, does there exist a neighborhood $U(z_0)$ of z_0 , such that the series $f(z) = \sum_{\nu=0}^{\infty} b_\nu(z-\zeta)^\nu$ has also H.-O.-gaps for all $\zeta \in U(z_0)$?

It is obvious that the corresponding problem cannot hold for power series with Hadamard-gaps or Fabry-gaps.

2. Power series with H.-O.-gaps

We first deal with power series having H.-O.-gaps and prove the following result.

Theorem 1. *Suppose that the power series $f(z) = \sum_{\nu=0}^{\infty} a_{\nu}(z - z_0)^{\nu}$ with radius of convergence R , $0 < R < \infty$ has a sequence of H.-O.-gaps.*

Then there is a neighborhood $U(z_0)$ of z_0 such that $f(z) = \sum_{\nu=0}^{\infty} b_{\nu}(z - \zeta)^{\nu}$ has also H.-O.-gaps for all $\zeta \in U(z_0)$.

Proof. Without loss of generality we may assume that $z_0 = 0$ and $R = 1$ and that $z = 1$ is a singularity of the function f . We denote by $\{p_k, q_k\}$ the sequence of H.-O.-gaps and in addition we can suppose that $a_{\nu} = 0$ for $p_k \leq \nu \leq q_k$.

We choose another sequence $\{p_k^*, q_k^*\}$ of H.-O.-gaps and constants $\gamma > 1$ and $\lambda^* > 1$ in the following way

$$p_k < p_k^* < q_k^* < q_k; \quad \frac{p_k^*}{p_k} \geq \gamma, \quad \frac{q_k}{q_k^*} \geq \gamma; \quad \frac{q_k^*}{p_k^*} \geq \lambda^*.$$

The partial sums of the considered power series are denoted by

$$s_n(z) = \sum_{\nu=0}^n a_{\nu} z^{\nu}.$$

Let be given a point $\zeta \in \mathbb{D}$, then the expansion of f around ζ :

$$f(z) = \sum_{\nu=0}^{\infty} b_{\nu}(z - \zeta)^{\nu}$$

has radius of convergence at most $|1 - \zeta|$.

1. For $p_k \leq n \leq q_k$ we have

$$s_n(z) = \sum_{\nu=0}^n b_{\nu}^{(\zeta)}(z - \zeta)^{\nu}$$

and we obtain for $0 < r < 1 - |\zeta|$ and $p_k \leq \nu \leq q_k$

$$b_{\nu} - b_{\nu}^{(\zeta)} = \frac{1}{2\pi i} \int_{|z-\zeta|=r} \frac{f(z) - s_{\nu}(z)}{(z - \zeta)^{\nu+1}} dz = \frac{1}{2\pi i} \int_{|z-\zeta|=r} \frac{f(z) - s_{q_k}(z)}{(z - \zeta)^{\nu+1}}.$$

We choose $0 < \varepsilon < \min\{1 - |\zeta| - r, r^{1/\gamma} - r\}$ and for all sufficiently great k it follows

$$\begin{aligned} |b_\nu - b_\nu^{(\zeta)}| &\leq \frac{1}{r^\nu} \cdot \max_{|z-\zeta| \leq r} |f(z) - s_{q_k}(z)| \leq \frac{1}{r^\nu} \cdot \max_{|z| \leq |\zeta|+r} |f(z) - s_{q_k}(z)| \leq \\ &\leq \frac{1}{r^\nu} \cdot (|\zeta| + r + \varepsilon)^{q_k}. \end{aligned}$$

Therefore, we have for $p_k^* \leq \nu \leq q_k^*$

$$\begin{aligned} |b_\nu - b_\nu^{(\zeta)}|^{1/\nu} &\leq \frac{1}{r} \cdot (|\zeta| + r + \varepsilon)^{\frac{q_k}{\nu}} \leq \frac{1}{r} \cdot (|\zeta| + r + \varepsilon)^{\frac{q_k}{p_k^*}} \leq \\ &\leq \frac{1}{r} (|\zeta| + r + \varepsilon)^\gamma. \end{aligned}$$

For $\zeta = 0$ we have $(r + \varepsilon)^\gamma < r$ and by continuity there exists $\zeta_1 \neq 0$ such that

$$(1 + |\zeta|)^2 \cdot (|\zeta| + r + \varepsilon)^\gamma < r$$

for all ζ with $|\zeta| < |\zeta_1|$. We therefore get for $0 < |\zeta| < |\zeta_1|$ and sufficiently great k

$$|b_\nu - b_\nu^{(\zeta)}|^{1/\nu} < \frac{1}{(1 + |\zeta|)^2} < \frac{1}{1 + |\zeta|} \leq \frac{1}{|1 - \zeta|}.$$

2. For $p_k \leq \nu \leq q_k$ and $R > 1$ we have

$$b_\nu^{(\zeta)} = \frac{1}{2\pi i} \int_{|z-\zeta|=R} \frac{s_\nu(z)}{(z-\zeta)^{\nu+1}} dz = \frac{1}{2\pi i} \int_{|z-\zeta|=R} \frac{s_{p_k}(z)}{(z-\zeta)^{\nu+1}} dz.$$

If we choose $\zeta \neq 0$ and $\varepsilon > 0$ so small that $R + |\zeta| + \varepsilon < R^\gamma$, we get for all sufficiently large k

$$\begin{aligned} |b_\nu^{(\zeta)}| &\leq \frac{1}{R^\nu} \cdot \max_{|z-\zeta| \leq R} |s_{p_k}(z)| \leq \frac{1}{R^\nu} \cdot \max_{|z| \leq |\zeta|+R} |s_{p_k}(z)| \leq \\ &\leq \frac{1}{R^\nu} \cdot (R + |\zeta| + \varepsilon)^{p_k}. \end{aligned}$$

Therefore, we have for $p_k^* \leq \nu \leq q_k^*$

$$\begin{aligned} |b_\nu^{(\zeta)}|^{1/\nu} &\leq \frac{1}{R} \cdot (R + |\zeta| + \varepsilon)^{\frac{p_k}{\nu}} \leq \frac{1}{R} \cdot (R + |\zeta| + \varepsilon)^{\frac{p_k}{p_k^*}} \leq \\ &\leq \frac{1}{R} \cdot (R + |\zeta| + \varepsilon)^{1/\gamma} \end{aligned}$$

and again as above we find a $\zeta_2 \neq 0$ such that for $0 < |\zeta| < |\zeta_2|$ and all sufficiently great k

$$|b_\nu^{(\zeta)}|^{1/\nu} < \frac{1}{(1 + |\zeta|)^2} < \frac{1}{1 + |\zeta|} \leq \frac{1}{|1 - \zeta|}.$$

3. It follows that for $I^* = \bigcup_{k=1}^\infty [p_k^*, q_k^*]$ and all ζ with

$$0 < |\zeta| < \min \{|\zeta_1|, |\zeta_2|\} =: |\zeta_0|$$

$$\overline{\lim}_{\substack{\nu \rightarrow \infty \\ \nu \in I^*}} |b_\nu|^{1/\nu} \leq \frac{1}{(1 + |\zeta|)^2} < \frac{1}{|1 - \zeta|},$$

so, that the series $f(z) = \sum_{\nu=0}^\infty b_\nu(z - \zeta)^\nu$ has H.-O.-gaps. ■

Remark 2.1. By slightly changing the proof one can choose $\zeta_0 \neq 0$ so small, that for $m \geq 2$

$$\overline{\lim}_{\substack{\nu \rightarrow \infty \\ \nu \in I^*}} |b_\nu|^{1/\nu} \leq \frac{1}{(1 + |\zeta|)^m}$$

for all ζ with $0 < |\zeta| < |\zeta_0|$. Therefore, the coefficients in the intervals $[p_k^*, q_k^*]$ become arbitrary small, when ζ is near to zero.

3. Power series with O.-gaps

We now consider power series $f(z) = \sum_{\nu=0}^\infty a_\nu(z - z_0)^\nu$ with radius of convergence $R, 0 < R < \infty$ and O.-gaps. It is well known [4] that such a series defines a function f , which is holomorphic in a simply connected domain $G(f)$ and every boundary point of $G(f)$ is a singularity for f .

We prove the following result.

Theorem 2. *Suppose that the power series $f(z) = \sum_{\nu=0}^\infty a_\nu(z - z_0)^\nu$ with radius of convergence $R, 0 < R < \infty$, has a sequence of O.-gaps.*

Then the power series $f(z) = \sum_{\nu=0}^\infty b_\nu(z - \zeta)^\nu$ has also O.-gaps for all $\zeta \in G(f)$.

Proof. Similar to the situation in the proof of Theorem 1 we again may assume that $z_0 = 0$ and $R = 1$, so that $f(z) = \sum_{\nu=0}^\infty a_\nu z^\nu$. The sequence of O.-gaps is denoted by $\{p_k, q_k\}$ and we may suppose that $a_\nu = 0$ for $p_k \leq \nu \leq q_k$, where $\frac{q_k}{p_k} \rightarrow \infty$ for $k \rightarrow \infty$.

We choose p_k^*, q_k^* with $p_k < p_k^* < q_k^* < q_k$ in such way that

$$\frac{p_k^*}{p_k} \rightarrow \infty, \quad \frac{q_k^*}{p_k^*} \rightarrow \infty, \quad \frac{q_k}{q_k^*} \rightarrow \infty \quad \text{for } k \rightarrow \infty.$$

Let be given a point $\zeta \in G(f)$, consider the power series

$$f(z) = \sum_{\nu=0}^{\infty} b_{\nu}(z - \zeta)^{\nu}$$

and suppose that its radius of convergence is $2r$.

1. We consider

$$s_{q_k^*}(z) = \sum_{\nu=0}^{q_k^*} a_{\nu} z^{\nu} = \sum_{\nu=0}^{q_k^*} b_{\nu}^{(\zeta)} (z - \zeta)^{\nu}.$$

For all sufficiently great k we get with a constant $\vartheta, 0 < \vartheta < 1$

$$\max_{|z-\zeta| \leq r} |f(z) - s_{q_k^*}(z)| = \max_{|z-\zeta| \leq r} |f(z) - s_{q_k}(z)| \leq \vartheta^{q_k}.$$

For $p_k \leq \nu \leq q_k$ we have

$$b_{\nu} - b_{\nu}^{(\zeta)} = \frac{1}{2\pi i} \int_{|z-\zeta|=r} \frac{f(z) - s_{q_k^*}(z)}{(z - \zeta)^{\nu+1}} dz$$

and it follows for $p_k^* \leq \nu \leq q_k^*$

$$|b_{\nu} - b_{\nu}^{(\zeta)}|^{1/\nu} \leq \frac{1}{r} \cdot \vartheta^{\frac{q_k}{\nu}}$$

which implies that for $I^* = \bigcup_{k=1}^{\infty} [p_k^*, q_k^*]$

$$\lim_{\substack{\nu \rightarrow \infty \\ \nu \in I^*}} |b_{\nu} - b_{\nu}^{(\zeta)}|^{1/\nu} = 0.$$

2. For $R > 1$ and $p_k^* \leq \nu \leq q_k^*$ we have for sufficiently great k

$$b_{\nu}^{(\zeta)} = \frac{1}{2\pi i} \int_{|z-\zeta|=R} \frac{s_{q_k^*}(z)}{(z - \zeta)^{\nu+1}} dz = \frac{1}{2\pi i} \int_{|z-\zeta|=R} \frac{s_{p_k}(z)}{(z - \zeta)^{\nu+1}} dz$$

and therefore we get for sufficiently great k

$$\begin{aligned} |b_{\nu}^{(\zeta)}|^{1/\nu} &\leq \frac{1}{R} \cdot \max_{|z| \leq |\zeta| + R} |s_{p_k}(z)|^{1/\nu} \leq \frac{1}{R} \cdot (|\zeta| + R + 1)^{\frac{p_k}{\nu}} \leq \\ &\leq \frac{1}{R} \cdot (|\zeta| + R + 1)^{\frac{p_k}{p_k^*}} \end{aligned}$$

which implies

$$\overline{\lim}_{\substack{\nu \rightarrow \infty \\ \nu \in I^*}} |b_\nu^{(\zeta)}|^{1/\nu} \leq \frac{1}{R}$$

and, since $R > 1$ was arbitrary, it follows

$$\lim_{\substack{\nu \rightarrow \infty \\ \nu \in I^*}} |b_\nu^{(\zeta)}|^{1/\nu} = 0.$$

3. It follows that for all $\zeta \in G(f)$ the power series $f(z) = \sum_{\nu=0}^{\infty} b_\nu(z - \zeta)^\nu$ has a sequence of O.-gaps $\{p_k^*, q_k^*\}$ (which is independent of ζ). In addition the sequence $s_{p_k^*}(z) = \sum_{\nu=0}^{p_k^*} b_\nu^{(\zeta)}(z - \zeta)^\nu$ converges to $f(z)$ compactly on $G(f)$. ■

References

- [1] **Große-Erdmann, K.-G.**, Universal families and hypercyclic operators, *Bull. Amer. Math. Soc.*, **36** (1999), 345–381.
- [2] **Hille, E.**, *Analytic Function Theory*, vol. II, Chelsea Publishing Company (New York, 1987).
- [3] **Ostrowski, A.**, Über eine Eigenschaft gewisser Potenzreihen mit unendlich vielen verschwindenden Koeffizienten, *S.-B. Preuss. Akad. Wiss.*, (Berlin) Phys.-Math. Kl. (1921), 557–565.
- [4] **Ostrowski, A.**, Über vollständige Gebiete gleichmäßiger Konvergenz von Folgen analytischer Funktionen, *Abh. Math. Sem. Hamburg. Univ.*, **1** (1922), 327–450.
- [5] **Ostrowski, A.**, Über Potenzreihen, die überkonvergente Abschnittsfolgen besitzen, *S.-B. Preuss. Akad. Wiss.*, (Berlin) Phys.-Math. Kl. (1923), 185–192.
- [6] **Ostrowski, A.**, Zur Theorie der Überkonvergenz, *Math. Ann.*, **103** (1930), 15–27.

W. Luh

Fachbereich IV – Mathematik

Universität Trier

D-54286 Trier

Germany

luh@uni-trier.de