

ON A THEOREM OF FEICHTINGER AND WEISZ

Péter Simon (Budapest, Hungary)

Dedicated to the 70th birthday of Professor Karl-Heinz Indlekofer

Communicated by Ferenc Schipp

(Received June 07, 2012; accepted November 27, 2012)

Abstract. The so-called θ -summation is well-known in the theory of approximation. A remarkable result gives a necessary and sufficient condition for uniformly or L^1 -norm convergence of θ -means if θ has compact support. This condition is nothing else but the integrability of the (trigonometric) Fourier transform of θ . Later this theorem was improved by Feichtinger and Weisz showing the same result for θ 's belonging to a suitable Wiener algebra $W(C, \ell_1)$. If θ is compactly supported then $\theta \in W(C, \ell_1)$ holds evidently but there are functions $\theta \in W(C, \ell_1)$ with unbounded support. In this work we extend the statement of Feichtinger and Weisz. To this end a new space $S(C, \ell_1)$ of functions will be constructed for which we prove the validity of the integrability condition. A simple consideration leads to the proper inclusion $W(C, \ell_1) \subset S(C, \ell_1)$.

1. Introduction

The so-called θ -summation, as a general method of summation generated by a single function θ is an intensively investigated area of approximation. (For this see e.g. [1], [5], [8] and references in [2], [6], [7] as illustration.) In

Key words and phrases: Fourier transform, inversion, periodization, θ -summation.

2010 Mathematics Subject Classification: Primary 42A10, 42A24, 42A38.

This research was supported by the European Union and co-financed by the European Social Fund (grant agreement no. TAMOP 4.2.1./B-09/1/KMR-2010-0003).

this paper we consider θ -means of trigonometric Fourier series and investigate the question: for what functions θ do we have convergence result. To this end we summarize briefly the most important concepts, definitions and well-known facts about θ -summation (for historical background see also the references).

Next we denote by L^1 the set of the functions $f : \mathbb{R} \rightarrow \mathbb{R}$ integrable in the sense of Lebesgue. Furthermore, let $\|f\|_1 := \int |f(x)| dx := \int_{-\infty}^{+\infty} |f(x)| dx$. If $g \in L^1$ then

$$Pg(t) := \sum_{k=-\infty}^{+\infty} g(t + 2k\pi) \quad t \in ([-\pi, \pi])$$

is the so-called periodization of g . Since

$$\int_{-\pi}^{\pi} \sum_{k=-\infty}^{+\infty} |g(t + 2k\pi)| dt = \int |g(t)| dt < +\infty,$$

the series $\sum_{k=-\infty}^{+\infty} g(t + 2k\pi)$ is absolutely convergent a.e. $t \in \mathbb{R}$. It is clear that $\int_{-\pi}^{\pi} |Pg(t)| dt \leq \|g\|_1$ and $\int_{-\pi}^{\pi} Pg(t) dt = \int g(t) dt$. Moreover, Pg is periodic by 2π .

Now let $f \in L^1[-\pi, \pi]$. We take it as $f : \mathbb{R} \rightarrow \mathbb{R}$ periodic by 2π and for $g \in L^1$ define $f \star g$ as the usual convolution of f and Pg :

$$f \star g(x) := f * Pg = \int_{-\pi}^{\pi} \sum_{k=-\infty}^{+\infty} f(x-t)g(t+2k\pi) dt \quad (x \in \mathbb{R}).$$

Then $f \star g \in L^1[-\pi, \pi]$ and $\int_{-\pi}^{\pi} |f \star g(x)| dx \leq \int_{-\pi}^{\pi} |f(x)| dx \cdot \|g\|_1 =: \|f\|_1 \cdot \|g\|_1$. For example if $f(t) := e^{ijt} \quad (j \in \mathbb{Z}, t \in [-\pi, \pi])$ then

$$\begin{aligned} e_j \star g(x) &= e^{ijx} \int_{-\pi}^{\pi} \sum_{k=-\infty}^{+\infty} e^{-ijt} g(t+2k\pi) dt = \\ &= e^{ijx} \int_{-\pi}^{\pi} \sum_{k=-\infty}^{+\infty} e^{-ij(t+2k\pi)} g(t+2k\pi) dt = e^{ijx} \int g(t) e^{-ijt} dt \quad (x \in \mathbb{R}), \end{aligned}$$

i.e. $e_j \star g(x) = e^{ijx} \hat{g}(-j) \quad (x \in \mathbb{R})$. Here \hat{g} stands for the (trigonometric) Fourier transform:

$$\hat{g}(x) := \int g(t) e^{itx} dt \quad (x \in \mathbb{R}).$$

We remember to the so-called inversion formula: if $g, \hat{g} \in L^1$ then

$$g(x) = \frac{1}{2\pi} \hat{g}(-x) = \frac{1}{2\pi} \int \hat{g}(t) e^{-itx} dt \quad (\text{a.e. } x \in \mathbb{R}).$$

(It is well-known that \hat{g} is continuous, so in this case it can be assumed that g is also continuous. Hence the above equality holds for all $x \in \mathbb{R}$.) Let $\theta \in L^1$ be given such that $\hat{\theta} \in L^1$ and with a natural number $m = 1, 2, \dots$ we take

$$\theta_m(t) := \frac{m}{2\pi} \cdot \hat{\theta}(mt) \quad (t \in \mathbb{R}).$$

By means of θ_m let us considered the operators

$$T_m^\theta f := f \star \theta_m \quad (f \in L^1[-\pi, \pi]).$$

Then for all $f \in L^1[-\pi, \pi]$ we have

$$\begin{aligned} \|T_m^\theta f\|_1 &= \|f \star \theta_m\|_1 \leq \|f\|_1 \cdot \|\theta_m\|_1 = \frac{m\|f\|_1}{2\pi} \int |\hat{\theta}(mt)| dt = \\ &= \frac{\|f\|_1}{2\pi} \int |\hat{\theta}(t)| dt = \frac{\|\hat{\theta}\|_1}{2\pi} \|f\|_1. \end{aligned}$$

In other words the sequence of (obviously linear) operators

$$T_m^\theta : L^1[-\pi, \pi] \rightarrow L^1[-\pi, \pi] \quad (0 < m \in \mathbb{N})$$

are uniformly bounded with respect to the norm $\|\cdot\|_1$ of the Banach space $L^1[-\pi, \pi]$. Special (see above) $T_m^\theta e_j = e_j \widehat{\theta}_m(-j)$ ($j \in \mathbb{Z}$), where by the inversion formula

$$\widehat{\theta}_m(-j) = \frac{m}{2\pi} \int \hat{\theta}(mt) e^{-\imath jt} dt = \frac{1}{2\pi} \int \hat{\theta}(t) e^{-\imath jt/m} dt = \theta(j/m) \quad (x \in \mathbb{R}).$$

Therefore $T_m^\theta e_j = \theta(j/m) e_j$ ($j \in \mathbb{Z}$).

Further we assume that the function θ is also continuous and satisfies the condition

$$C_m := \sum_{k=-\infty}^{+\infty} |\theta(k/m)| < +\infty \quad (0 < m \in \mathbb{N}).$$

Under these assumptions we consider the mappings σ_m^θ ($0 < m \in \mathbb{N}$) as follows:

$$\sigma_m^\theta f := \sum_{k=-\infty}^{+\infty} \theta(k/m) c_k(f) e_k \quad (f \in L^1[-\pi, \pi]),$$

where $c_k(f) := (2\pi)^{-1} \int_{-\pi}^{\pi} f(t) e^{-\imath kt} dt$ is the usual k -th Fourier coefficient of f . Since $|c_k(f)| \leq \|f\|_1 / (2\pi)$ ($k \in \mathbb{Z}$), i.e. $\sum_{k=-\infty}^{+\infty} |\theta(k/m) c_k(f) e_k| \leq$

$\leq C_m \|f\|_1 / (2\pi)$, thus the series in question converges uniformly and for every $f \in L^1[-\pi, \pi]$ we get

$$\|\sigma_m^\theta f\|_1 \leq \sum_{k=-\infty}^{+\infty} |\theta(k/m)| \cdot |c_k(f)| \cdot \|e_k\|_1 \leq C_m \|f\|_1.$$

This means that for all $m = 1, 2, \dots$ the (linear) operator $\sigma_m^\theta : L^1[-\pi, \pi] \rightarrow L^1[-\pi, \pi]$ is also bounded. Furthermore,

$$c_k(e_j) = \delta_{kj} = \begin{cases} 1 & (k = j) \\ 0 & (k \neq j) \end{cases} \quad (k, j \in \mathbb{Z}),$$

which involves (see above) $\sigma_m^\theta e_j = \theta(j/m)e_j = T_m^\theta e_j$ ($j \in \mathbb{Z}, 0 < m \in \mathbb{N}$). From this it follows immediately the analogous equality for all trigonometric polynomials. They form a dense set in $L^1[-\pi, \pi]$ with respect to $\|\cdot\|_1$, therefore

$$\sigma_m^\theta f = T_m^\theta f \quad (f \in L^1[-\pi, \pi], 0 < m \in \mathbb{N}).$$

A simple calculation shows that

$$\sigma_m^\theta f(x) = \int_{-\pi}^{\pi} f(t) K_m^\theta(x-t) dt \quad (x \in [-\pi, \pi]),$$

where the (2π periodic) kernel K_m^θ is defined as follows:

$$K_m^\theta := \frac{1}{2\pi} \sum_{k=-\infty}^{\infty} \theta(k/m) e_k \quad (0 < m \in \mathbb{N}).$$

The assumption $C_m < +\infty$ guaranties that K_m^θ is continuous. On the other hand

$$\begin{aligned} \|\sigma_m^\theta f\|_\infty &:= \max_{x \in [-\pi, \pi]} |\sigma_m^\theta f(x)| = \|T_m^\theta f\|_\infty \leq \\ &\leq \|f\|_\infty \cdot \|\theta_m\|_1 = \frac{1}{2\pi} \|\hat{\theta}\|_1 \|f\|_\infty \quad (f \in C[-\pi, \pi]). \end{aligned}$$

This leads by standard argument to the inequality $\|K_m^\theta\|_1 \leq \|\hat{\theta}\|_1 / (2\pi)$ ($m = 1, 2, \dots$), i.e.

$$\sup_{0 < m \in \mathbb{N}} \|K_m^\theta\|_1 \leq \frac{1}{2\pi} \|\hat{\theta}\|_1.$$

A remarkable result in the theory of approximation (see e.g. [8]) says that in the above estimation we can write equality:

$$\sup_{0 < m \in \mathbb{N}} \|K_m^\theta\|_1 = \frac{1}{2\pi} \|\hat{\theta}\|_1.$$

Moreover, the "reverse" implication was also investigated. Namely (see e.g. [3], [4], [8]), if a continuous and integrable function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ has compact support, then the next implication is true:

$$(*) \quad \sup_{0 < m \in \mathbb{N}} \|K_m^\theta\|_1 < +\infty \implies \hat{\theta} \in L^1.$$

Here the assumption on the compactness of the support of θ can be weakened. To this end let for a function $f : \mathbb{R} \rightarrow \mathbb{R}$

$$\|f\|_W := \sum_{k=-\infty}^{\infty} \sup_{x \in [0,1)} |f(k+x)|$$

and denote $W(C, \ell_1)$ the Wiener algebra of all continuous functions $f : \mathbb{R} \rightarrow \mathbb{R}$ for which $\|f\|_W < +\infty$. Then the following statement holds (Feichtinger and Weisz [2]): the assumption $\theta \in W(C, \ell_1)$ is enough to the implication (*). It is clear that every continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$ with compact support belongs to $W(C, \ell_1)$. Furthermore, a simple example can be constructed to show that there are functions in $W(C, \ell_1)$ with unbounded supports.

2. The space $(S(C, \ell_1), \|\cdot\|_S)$

Next we prove that the just mentioned result of Feichtinger and Weisz can be improved. In other words the space $W(C, \ell_1)$ can be so enlarged that the implication (*) remains true. For this purpose we introduce for a continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$ a new norm $\|f\|_S$ as follows:

$$\|f\|_S := \sum_{j=-\infty}^{+\infty} \sup_{0 < m \in \mathbb{N}} \frac{1}{m} \sum_{l=0}^{m-1} |f(j+l/m)|$$

and the space

$$S(C, \ell_1) := \{f \in C : \|f\|_S < +\infty\}.$$

Since $m^{-1} \sum_{l=0}^{m-1} |f(j+l/m)|$ ($f \in S(C, \ell_1), 0 < m \in \mathbb{N}, j \in \mathbb{Z}$) is a Riemann sum of the integral $\int_j^{j+1} |f(t)| dt$ thus

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{l=0}^{m-1} |f(j+l/m)| = \int_j^{j+1} |f(t)| dt.$$

Therefore

$$\sup_{0 < m \in \mathbb{N}} \frac{1}{m} \sum_{l=0}^{m-1} |f(j+l/m)| \geq \int_j^{j+1} |f(t)| dt$$

and we get

$$\begin{aligned} \int |f(t)| dt &= \sum_{j=-\infty}^{+\infty} \int_j^{j+1} |f(t)| dt \leq \\ &\leq \sum_{j=-\infty}^{+\infty} \sup_{0 < m \in \mathbb{N}} \frac{1}{m} \sum_{l=0}^{m-1} |f(j+l/m)| = \|f\|_S < +\infty. \end{aligned}$$

Hence $S(C, \ell_1) \subset L^1$. Furthermore, if $f \in S(C, \ell_1)$ and $m = 1, 2, \dots$ then

$$\sum_{k=-\infty}^{+\infty} |f(k/m)| \leq m \cdot \sum_{j=-\infty}^{+\infty} \frac{1}{m} \sum_{l=0}^{m-1} |f(j+l/m)| \leq m \cdot \|f\|_S < +\infty.$$

Next we list some basic properties of $(S(C, \ell_1), \|\cdot\|_S)$.

1° First of all, a simple consideration shows that $(S(C, \ell_1), \|\cdot\|_S)$ is a normed space. Indeed, $\|0\|_S = 0$ is trivial. If $f \in S(C, \ell_1)$ and $\|f\|_S = 0$ then for every $j \in \mathbb{Z}$, $0 < m \in \mathbb{N}$ we have $f(j+l/m) = 0$ ($l = 0, \dots, m-1$). Let $x \in \mathbb{R}$, $\varepsilon > 0$. By the continuity of f there are $j \in \mathbb{Z}$, $0 < m \in \mathbb{N}$, $l = 0, \dots, m-1$ such that with $y := j+l/m$ the inequality $|f(x) - f(y)| = |f(x)| < \varepsilon$ holds. Hence $f(x) = 0$, i.e. $f \equiv 0$. Furthermore, the equality $\|\lambda f\|_S = |\lambda| \cdot \|f\|_S$ ($f \in S(C, \ell_1), \lambda \in \mathbb{R}$) and the inequality $\|f+g\|_S \leq \|f\|_S + \|g\|_S$ ($f, g \in S(C, \ell_1)$) are obvious.

2° Now, we take a sequence $f_n \in S(C, \ell_1)$ ($n \in \mathbb{N}$) of functions which is convergent in $S(C, \ell_1)$. In other words there exists $f \in S(C, \ell_1)$ such that

$$\|f_n - f\|_S \rightarrow 0 \quad (n \rightarrow \infty).$$

This means that

$$\sum_{j=-\infty}^{+\infty} \sup_{0 < m \in \mathbb{N}} \frac{1}{m} \sum_{l=0}^{m-1} |f_n(j+l/m) - f(j+l/m)| \rightarrow 0 \quad (n \rightarrow \infty).$$

If $0 \neq r \in \mathbb{R}$ is a rational number then with suitable $j_0 \in \mathbb{Z}$, $0 < m_0 \in \mathbb{N}$, $l_0 = 0, \dots, m_0 - 1$ the equality $r = j_0 + l_0/m_0$ holds. It is clear that

$$|f_n(r) - f(r)| \leq \sum_{l=0}^{m_0-1} |f_n(j_0+l/m_0) - f(j_0+l/m_0)| \leq m_0 \|f_n - f\|_S \quad (n \in \mathbb{N}),$$

i.e. $f(r) = \lim_{n \rightarrow \infty} f_n(r)$.

3° The space $(S(C, \ell_1), \|\cdot\|_S)$ does not form a Banach space. Indeed, if $n = 1, 2, \dots$ and

$$f_n(x) := \begin{cases} \sin(\pi/x) & (1/n \leq x \leq 1) \\ 0 & (x \in \mathbb{R} \setminus (1/n, 1)), \end{cases}$$

then $f_n \in C$ and

$$\|f_n\|_S = \sup_{0 < m \in \mathbb{N}} \frac{1}{m} \sum_{l=0}^{m-1} |f_n(l/m)| \leq 1$$

implies $f_n \in S(C, \ell_1)$. Furthermore, if $n, k, m = 1, 2, \dots$ and $k > n$, then

$$\begin{aligned} \sum_{l=0}^{m-1} |f_n(l/m) - f_k(l/m)| &= \sum_{l=0, 1/k < l/m < 1/n}^{m-1} |f_n(l/m) - f_k(l/m)| = \\ &= \sum_{l=0, m/k < l < m/n}^{m-1} |f_k(l/m)| \leq \frac{m}{n} - \frac{m}{k}. \end{aligned}$$

From this it follows that

$$\begin{aligned} \|f_n - f_k\|_S &= \sup_{0 < m \in \mathbb{N}} \frac{1}{m} \sum_{l=0}^{m-1} |f_n(l/m) - f_k(l/m)| \leq \\ &\leq \frac{1}{n} - \frac{1}{k} \rightarrow 0 \quad (n, k \rightarrow \infty). \end{aligned}$$

Hence the sequence (f_n) is a Cauchy sequence with respect to $\|\cdot\|_S$. Assume the existence $f \in S(C, \ell_1)$ such that $\|f_n - f\|_S \rightarrow 0$ ($n \rightarrow \infty$). Then it would be true by 2° for all rational $r \in (0, 1)$ that

$$f(r) = \lim_{n \rightarrow \infty} f_n(r) = \sin(\pi/r).$$

However, such a continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$ does not exist.

4° Let $f \in C[0, 1]$ and

$$s_m(f) := \frac{1}{m} \sum_{l=0}^{m-1} |f(l/m)| \quad (m = 1, 2, \dots).$$

If

$$f_{ml} := \max\{|f(t)| : l/m \leq t \leq (l+1)/m\} \quad (l = 0, \dots, m-1)$$

and

$$S_m(f) := \frac{1}{m} \sum_{l=0}^{m-1} f_{ml} \quad (m = 1, 2, \dots),$$

then

$$s(f) := \sup_m s_m(f) \leq S(f) := \sup_m S_m(f).$$

Now let $n = 1, 2, \dots$ be given and let us consider the function $f_n \in C[0, 1]$ in the following way: $f_n(0) := f_n(t) := 0 \quad (1/n \leq t \leq 1)$, $f_n(1/(2n)) := 1$ and the graph of f_n over $[0, 1/n]$ is a triangle. Then $S_1(f_n) = 1$ which implies $S(f_n) \geq 1$. On the other hand for $m = 1, \dots, n$ it follows $s_m(f_n) = 0$ but

$$s_m(f_n) = \frac{1}{m} \sum_{l=1}^{[m/n]} f_n(l/m) \leq \frac{1}{m} \sum_{l=1}^{[m/n]} 1 \leq \frac{1}{n} \quad (m = n + 1, n + 2, \dots).$$

Therefore $s(f_n) \leq 1/n \quad (0 < n \in \mathbb{N})$, i.e. it does not exist constant $q \geq 0$ such that $S(f) \leq q \cdot s(f) \quad (f \in C[0, 1])$.

5° Define for $f \in S(C, \ell_1)$ the symbol $\|f\|_{SW}$ as follows:

$$\|f\|_{SW} := \sum_{j=-\infty}^{+\infty} \sup_{0 < m \in \mathbb{N}} \frac{1}{m} \sum_{l=0}^{m-1} \|f \chi_{[j+l/m, j+(l+1)/m]}\|_\infty.$$

It is not hard to see that $\|\cdot\|_{SW}$ is norm and $\|\cdot\|_S \leq \|\cdot\|_{SW}$. However, the functions $f_n \quad (n = 1, 2, \dots)$ from 4° show that $\|\cdot\|_S, \|\cdot\|_{SW}$ are not equivalent.

6° It is clear that for all continuous functions $\theta : \mathbb{R} \rightarrow \mathbb{R}$

$$\frac{1}{m} \sum_{l=0}^{m-1} |\theta(j + l/m)| \leq \sup_{0 \leq x < 1} |\theta(j + x)| \quad (j \in \mathbb{Z}),$$

which means that $W(C, \ell_1) \subset S(C, \ell_1)$. A simple example proves that this inclusion is proper, i.e. $W(C, \ell_1) \neq S(C, \ell_1)$. Indeed, let $\theta : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function such that $\|\theta \chi_{(j, j+1/j)}\|_\infty = 1/j \quad (j = 1, 2, \dots)$ and $\theta(t) = 0 \quad (t \in \mathbb{R} \setminus A)$, where $A := \bigcup_{j=1}^\infty (j, j + 1/j)$ (see also Feichtinger and Weisz [2]). Then $\theta \in L^1$ and for all $m = 1, 2, \dots$ and $j \in \mathbb{Z}$

$$\frac{1}{m} \sum_{l=0}^{m-1} |\theta(j + l/m)| \leq \begin{cases} 0 & (j \leq 0) \\ m^{-1} \sum_{l=1}^{[m/j]} 1/j \leq j^{-2} & (j > 0). \end{cases}$$

Hence $\theta \in S(C, \ell_1)$. However,

$$\sum_{j=-\infty}^{+\infty} \sup_{0 \leq x < 1} |\theta(j + x)| = \sum_{j=1}^{+\infty} \frac{1}{j} = +\infty,$$

in other words $\theta \notin (W, \ell_1)$.

3. Main result

Now, we prove the main result of this work.

Theorem 3.1. *For all functions $\theta \in S(C, \ell_1)$ the implication $(*)$ is true.*

Proof. Let m, M, N be positive natural numbers and assume $M \leq m\pi$. Then $\|K_m^\theta\|_1$ can be considered as follows (the basic idea in the first steps is due to [8]):

$$\begin{aligned} 2\pi\|K_m^\theta\|_1 &= \int_{-\pi}^{\pi} \left| \sum_{k=-\infty}^{+\infty} \theta(k/m)e^{ikt} \right| dt = \int_{-m\pi}^{m\pi} \left| \frac{1}{m} \sum_{k=-\infty}^{+\infty} \theta(k/m)e^{ikt/m} \right| dt \geq \\ &\geq \int_{-M}^M \left| \frac{1}{m} \sum_{k=-\infty}^{+\infty} \theta(k/m)e^{ikt/m} \right| dt \geq \\ &\geq \int_{-M}^M \left| \frac{1}{m} \sum_{k=-mN}^{mN} \theta(k/m)e^{ikt/m} \right| dt - \int_{-M}^M \left| \frac{1}{m} \sum_{|k|>mN} \theta(k/m)e^{ikt/m} \right| dt \geq \\ &\geq \int_{-M}^M \left| \frac{1}{m} \sum_{k=-mN}^{mN} \theta(k/m)e^{ikt/m} \right| dt - \frac{2M}{m} \sum_{|k|>mN} |\theta(k/m)|. \end{aligned}$$

Here the sum $m^{-1} \sum_{k=-mN}^{mN} \theta(k/m)e^{ikt/m}$ ($-M \leq t \leq M$) is nothing else but a Riemann sum of the continuous function $[-N, N] \ni x \mapsto \theta(x)e^{itx}$. Therefore

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{k=-mN}^{mN} \theta(k/m)e^{ikt/m} = \int_{-N}^N \theta(x)e^{itx} dx \quad (-M \leq t \leq M).$$

However, if $-M \leq t \leq M$ then

$$\left| \frac{1}{m} \sum_{k=-mN}^{mN} \theta(k/m)e^{ikt/m} \right| \leq (2N + 1) \max_{|x| \leq N} |\theta(x)|,$$

i.e. by the Lebesgue's dominated convergence theorem

$$\lim_{m \rightarrow \infty} \int_{-M}^M \left| \frac{1}{m} \sum_{k=-mN}^{mN} \theta(k/m)e^{ikt/m} \right| dt = \int_{-M}^M \left| \int_{-N}^N \theta(x)e^{itx} dx \right| dt.$$

Furthermore,

$$\frac{1}{m} \sum_{|k| > mN} |\theta(k/m)| \leq \frac{1}{m} \sum_{|j| \geq N} \sum_{l=0}^{m-1} |\theta(j+l/m)| \leq \sum_{|j| \geq N} \gamma_j,$$

where

$$\gamma_j := \sup_{0 < n \in \mathbb{N}} \frac{1}{n} \sum_{l=0}^{n-1} |\theta(j+l/n)|.$$

We remark that $\theta \in S(C, \ell_1)$ implies $\sum_{j=-\infty}^{+\infty} \gamma_j = \|\theta\|_S < +\infty$, i.e. $\sum_{|j| \geq N} \gamma_j \rightarrow 0$ ($N \rightarrow \infty$).

Summarizing the above facts we get

$$2\pi \|K_m^\theta\|_1 \geq \int_{-M}^M \left| \frac{1}{m} \sum_{k=-mN}^{mN} \theta(k/m) e^{ikt/m} \right| dt - 2M \sum_{|j| \geq N} \gamma_j,$$

from which it follows that

$$\begin{aligned} 2\pi \sup_{0 < m \in \mathbb{N}} \|K_m^\theta\|_1 &\geq \lim_{m \rightarrow \infty} \int_{-M}^M \left| \frac{1}{m} \sum_{k=-mN}^{mN} \theta(k/m) e^{ikt/m} \right| dt - 2M \sum_{|j| \geq N} \gamma_j = \\ &= \int_{-M}^M \left| \int_{-N}^N \theta(x) e^{itx} dx \right| dt - 2M \sum_{|j| \geq N} \gamma_j. \end{aligned}$$

Taking into account $\left| \int_{-N}^N \theta(x) e^{itx} dx \right| \leq \|\theta\|_1$ ($|t| \leq M$) the above mentioned theorem of Lebesgue guaranties that

$$\begin{aligned} \lim_{N \rightarrow \infty} \int_{-M}^M \left| \int_{-N}^N \theta(x) e^{itx} dx \right| dt &= \int_{-M}^M \left| \lim_{N \rightarrow \infty} \int_{-N}^N \theta(x) e^{itx} dx \right| dt = \\ &= \int_{-M}^M \left| \int_{-\infty}^{+\infty} \theta(x) e^{itx} dx \right| dt = \int_{-M}^M |\hat{\theta}(t)| dt. \end{aligned}$$

Thus

$$\sup_{0 < m \in \mathbb{N}} \|K_m^\theta\|_1 \geq \lim_{N \rightarrow \infty} \int_{-M}^M \left| \int_{-N}^N \theta(x) e^{itx} dx \right| dt - 2M \lim_{N \rightarrow \infty} \sum_{|j| \geq N} \gamma_j = \int_{-M}^M |\hat{\theta}(t)| dt,$$

from which

$$\|\hat{\theta}\|_1 = \lim_{M \rightarrow \infty} \int_{-M}^M |\hat{\theta}(t)| dt \leq 2\pi \sup_{0 < m \in \mathbb{N}} \|K_m^\theta\|_1 < +\infty,$$

i.e., $\hat{\theta} \in L^1$ follows.

4. Remarks

1° It is clear that for compactly supported θ the proof can be simplified (see e.g. [8]). Namely, in this case $\text{supp } \theta \subset [-N, N]$ can be supposed in the above proof. Then $\sum_{|k| > mN} \theta(k/m)e^{ikt/m} = 0$ ($|t| \leq M$) holds trivially, hence

$$2\pi \|K_m^\theta\|_1 \geq \int_{-M}^M \left| \frac{1}{m} \sum_{k=-mN}^{mN} \theta(k/m)e^{ikt/m} \right| dt.$$

Therefore

$$\begin{aligned} 2\pi \sup_{0 < m \in \mathbb{N}} \|K_m^\theta\|_1 &\geq \lim_{m \rightarrow \infty} \int_{-M}^M \left| \frac{1}{m} \sum_{k=-mN}^{mN} \theta(k/m)e^{ikt/m} \right| dt = \\ &= \int_{-M}^M \left| \int_{-N}^N \theta(x)e^{itx} dx \right| dt = \int_{-M}^M |\hat{\theta}(x)| dt \end{aligned}$$

and the proof can be finished as above.

2° If $\theta \in S(C, \ell_1)$ then

$$\sup_{0 < m \in \mathbb{N}} \frac{1}{m} \sum_{k=-\infty}^{+\infty} |\theta(k/m)| \leq \sum_{j=-\infty}^{+\infty} \sup_{0 < m \in \mathbb{N}} \frac{1}{m} \sum_{l=0}^{m-1} |\theta(j + l/m)| = \|\theta\|_S < +\infty.$$

So the next estimation holds:

$$(**) \quad \sup_{0 < m \in \mathbb{N}} \frac{1}{m} \sum_{k=-\infty}^{+\infty} |\theta(k/m)| < +\infty.$$

On the other hand a continuous function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ can be constructed such that

$$\sum_{l=0}^{m-1} |\theta(j + l/m)| \sim |j|^{-1-1/m} \quad (0 < m, |j| \in \mathbb{N}).$$

For this θ it follows that $\sum_{k=-\infty}^{+\infty} |\theta(k/m)| \sim m$. This means that $(**)$ holds but θ does not belong to $S(C, \ell_1)$. Indeed, then $\gamma_j \sim 1/(|j| \ln |j|)$ ($1 < |j| \in \mathbb{N}$), i.e. $\|\theta\|_S = +\infty$.

3° The following question remains open: is the assumption $(**)$ on a continuous and integrable function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ enough to the implication $(*)$?

4° Let $\theta \in S(C, \ell_1)$ and $(X, \|\cdot\|_*)$ be defined as follows:

$$(X, \|\cdot\|_*) := \begin{cases} (C[-\pi, \pi], \|\cdot\|_\infty) \\ \text{or} \\ (L^1[-\pi, \pi], \|\cdot\|_1). \end{cases}$$

It is well-known that the norm of the operator $\sigma_m^\theta : X \rightarrow X$ is nothing else but $\|K_m^\theta\|_1$ ($m = 1, 2, \dots$). Assume that the function θ satisfies also $\theta(0) = 1$. Then a simple calculation shows that $\|\sigma_m^\theta e_j - e_j\|_* \rightarrow 0$ ($m \rightarrow \infty$), from which the same convergence follows for all trigonometric polynomials. Since the set of the trigonometric polynomials is dense in X , the theorem of Banach and Steinhaus (taking into account also our theorem) implies the following corollary:

$$\lim_{m \rightarrow \infty} \|\sigma_m^\theta f - f\|_* = 0 \quad (f \in X) \iff \hat{\theta} \in L^1.$$

5° We remark that (see Feichtinger and Weisz [2]) if $\theta \in W(C, \ell_1)$, $\theta(0) = 1$, then

$$\lim_{m \rightarrow \infty} \|\sigma_m^\theta f - f\|_2 = 0 \quad (f \in L^2[-\pi, \pi]).$$

However, let $\theta : \mathbb{R} \rightarrow \mathbb{R}$ be continuous such that $\|\theta \chi_{(j, j+1/j^2)}\|_\infty = \sqrt{j}$ ($j = 1, 2, \dots$) and $\theta(t) = 0$ ($t \in \mathbb{R} \setminus B$), where $B := \bigcup_{j=1}^\infty (j, j + 1/j^2)$ and $\theta(j + 1/(2j^2)) = \sqrt{j}$ ($0 < j \in \mathbb{N}$). For this function the relation $\theta \in S(C, \ell_1)$ follows with $j^{-3/2}$ instead of j^{-2} ($0 < j \in \mathbb{N}$) analogously as above in similar situation (see our example for the illustration of $W(C, \ell_1) \neq S(C, \ell_1)$). Furthermore for $m := 2j^2$, $k := jm + 1$ ($0 < j \in \mathbb{N}$) we get

$$\frac{k}{m} = j + \frac{1}{2j^2},$$

hence $|\theta(k/m)| = \sqrt{j}$. This implies

$$\sup_{0 < m \in \mathbb{N}} \sup_{k \in \mathbb{Z}} |\theta(k/m)| = +\infty.$$

The norm of the operator $\sigma_m^\theta : L^2[-\pi, \pi] \rightarrow L^2[-\pi, \pi]$ ($0 < m \in \mathbb{N}$) is $\|\sigma_m^\theta\| = \sup_{k \in \mathbb{Z}} |\theta(k/m)|$ (see [2]). Hence $\sup_{0 < m \in \mathbb{N}} \|\sigma_m^\theta\| = +\infty$ and the theorem of Banach and Steinhaus gives $f \in L^2$ such that the sequence $(\sigma_m^\theta f)$ diverges in $\|\cdot\|_2$ norm. In other words the last mentioned Feichtinger and Weisz's theorem on L^2 -convergence cannot be extended to the space $S(C, \ell_1)$.

References

- [1] **Butzer, P.L. and R.J. Nessel**, *Fourier Analysis and Approximation*, Birkhäuser, Basel, 1971.
- [2] **Feichtinger, H.G. and F. Weisz**, The Segal algebra $\mathbf{S}_0(\mathbb{R}^d)$ and norm summability of Fourier series and Fourier transforms, *Monatsh. Math.*, **148** (2006), 333–349.
- [3] **Trigub, R.M.**, A connection between summability and absolute convergence of Fourier series and transforms, *Dokl. Akad. Nauk SSSR*, **217** (1974), 34–37.
- [4] **Trigub, R.M.**, Integrability of the Fourier transform of a function with compact support, *Teor. Funkcii Funkcional. Anal. i Prilozen. Vyp.*, **23** (1975), 124–131.
- [5] **Trigub, R.M. and E.S. Belinsky**, *Fourier Analysis and Approximation of Functions*, Kluwer Academic Publishers, Dordrecht, Boston, London, 2004.
- [6] **Weisz, F.**, *Summability of Multi-Dimensional Fourier Series and Hardy Spaces. Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, Boston, London, 2002, 350 pp.
- [7] **Weisz, F.**, *Summability of Multi-Dimensional Trigonometric Fourier Series*, *Surveys in Approximation Theory*, **7** (2012), 1–179.
- [8] **Zhuk, V.V. and G.I. Natanson**, *Trigonometric Fourier Series and Elements of Approximation Theory*, Leningrad. Univ., Leningrad, 1983.

P. Simon

Department of Numerical Analysis
Eötvös Loránd University
Pázmány Péter sétány 1/C
H-1117 Budapest, Hungary
simon@ludens.elte.hu