

A GENERALIZATION OF THE ROOT FUNCTION

Tamás Dózsa and Ferenc Schipp

(Budapest, Hungary)

Communicated by László Szili

(Received May 1, 2021; accepted June 26, 2021)

Abstract. We consider the interpretation and the numerical construction of the inverse branches of n factor Blaschke-products on the disk \mathbb{D} and show that these provide a generalization of the n -th root function. The inverse branches can be defined on pairwise disjoint regions, whose union provides the disk. An explicit formula can be given for the n factor Blaschke-products on the torus, which can be used to provide the inverse branches on \mathbb{T} . The inverse branches can be thought of as the solutions $z = z_t(r)$ ($0 \leq r \leq 1$) to the equation $B(z) = re^{it}$, where B denotes an n factor Blaschke-product. We show that starting from a known value $z_t(1)$, any $z_t(r)$ point of the solution trajectory can be reached in finite steps. The appropriate grouping of the trajectories leads to two natural interpretations of the inverse branches (see Figure 2). We introduce an algorithm which can be used to find the points of the trajectories.

1. Introduction



In this work we consider a generalization of the n -th root function $w \rightarrow \sqrt[n]{w}$ ($w \in \mathbb{C}$). Using polar coordinates $w = re^{it}$ ($0 \leq r < \infty$, $t \in \mathbb{I} := [-\pi, \pi)$), we can express the branches of the root function as

$$(1.1) \quad \begin{aligned} \varphi_k(r, t) &:= r^{1/n} e^{i(t/n + 2k\pi/n)} \\ (r > 0, t \in \mathbb{I}, k = 0, 1, \dots, n-1). \end{aligned}$$

Key words and phrases: Blaschke-functions, Blaschke-products, root-function, Newton-iteration.

2010 Mathematics Subject Classification: Primary 30J10, Secondary 26C15, 30J99, 65T99.
EFOP-3.6.3-VEKOP-16-2017-00001: Talent Management in Autonomous Vehicle Control Technologies – The Project is supported by the Hungarian Government and co-financed by the European Social Fund.

One can acquire all of the branches from a single one, using rotations of angle $2\pi/n$. The roots' absolute values $r^{1/n}$ and angles can be found with Newton's method and division respectively. Figure (1a) shows the range of the branches of the n -th root function for $n = 3$.

(a) Ranges of φ_k 

(b) Initial points of Newton's method

Figure 1: Branches of the n -th root function

The solution of the equation $z^n = w$ ($w \in \mathbb{R}$) is equivalent to the n -th root function's value at w . In figure (1b) we illustrate the roots of w for $n = 3$ on the unit disk. Each color represents a single root denoted by z_1, z_2, z_3 . The roots can be identified as the limit of the Newton iteration

$$(1.2) \quad v_{k+1} = v_k - \frac{g(v_k)}{g'(v_k)}, \quad g(v) = v^n - w \quad (v_k \in \mathbb{D}, k = 0, 1, \dots).$$

Any $v_0 \in \mathbb{D}$ initial points for which the method diverges are colored black, the rest of the colors show the basins of attractions for each root z_k . The figure shows that choosing an initial point close to a particular root will result in Newton's method (quadratically) converging to that root.

In this paper we propose a generalization of the above problem by considering n factor Blaschke-products instead of the n -th root function [2]. Blaschke-functions have proven to be fundamental tools in the factorization of analytic functions and the construction of rational bases, while playing an important role for many applications in signal processing and control theory [3, 4, 5, 11, 12, 13]. The Blaschke-functions

$$(1.3) \quad B_a(z) := \frac{z - a}{1 - \bar{a}z} \quad (z \in \mathbb{C}, a \in \mathbb{D} := \{z \in \mathbb{C} : |z| < 1\})$$

are bijections on the disk \mathbb{D} and torus $\mathbb{T} := \{z \in \mathbb{C} : |z| = 1\}$. They can also be used to describe congruent transformations in the Poincaré model of the

Bolyai–Lobachevskian geometry [5, 14]. The n -factor Blaschke-products

$$(1.4) \quad \begin{aligned} B(z) &:= \epsilon \prod_{k=1}^n \frac{z - a_k}{1 - \bar{a}_k z} \\ (z \in \mathbb{C}, \epsilon \in \mathbb{T}, A &:= \{a_k : 1 \leq k \leq n\} \subset \mathbb{D}) \end{aligned}$$

map \mathbb{D} , \mathbb{T} and their complements into themselves. We refer to the set $A' := \{b \in \mathbb{C} : B'(b) = 0\}$ as the set of *critical points* and to the values $B(b)$ $b \in A'$ as *critical values*. If a line segment $S_t := \{re^{it} : 0 \leq r \leq 1\}$ takes a critical value for some r , then t is referred to as a critical parameter. The set of critical parameters will be denoted by T_0 . We note that if $a_1 = \dots = a_n = 0, \epsilon = 1$, then $B(z) = z^n$ ($z \in \mathbb{C}$) therefore the inverse of B can be considered as a generalization of the n -th root function.

The Riesz-factorization provides the basis for several algorithms which deal with the rational representation of signals [6, 7, 17]. It can be considered as a generalization of polar coordinates for functions in $H^p(\mathbb{D})$ ($0 < p \leq \infty$) Hardy-spaces. In this work instead of $H^p(\mathbb{D})$, we consider the class of functions \mathcal{A}_R , which are analytic on $\mathbb{D}_R := \{z \in \mathbb{C} : |z| < R\}$. For this class, if the roots of a function f are known, its Riesz-factorization can be easily acquired. Namely, let a_1, \dots, a_n denote the zeros of $f \in \mathcal{A}_R$. Then the factorization is given as

$$(1.5) \quad \begin{aligned} f(z) &= f_1(z) \prod_{k=1}^n (z - a_k) = \\ &= B(z) f_1(z) \prod_{k=1}^n (1 - \bar{a}_k z) = B(z) S(z) \quad (z \in \mathbb{D}_R), \end{aligned}$$

where the function $S \in \mathcal{A}_R$ has no zeros on \mathbb{D} . The existence of such a factorization, when $f \in H^p(\mathbb{D})$ is however not easy to prove [13]. Since $S(z) \neq 0$ ($z \in \mathbb{D}$) and $|B(z)| = 1$ ($z \in \mathbb{T}$), this factorization is often considered a generalization of polar coordinates.

It is well known that $B : \mathbb{T} \rightarrow \mathbb{T}$ provides an n -fold mapping of \mathbb{T} , moreover there exists a $\beta = \beta_n : \mathbb{R} \rightarrow \mathbb{R}$ strictly increasing function $\beta(t + 2\pi) = \beta(t) + 2\pi$ ($t \in \mathbb{R}$) for which

$$(1.6) \quad B(e^{it}) = e^{in\beta(t)} \quad (t \in \mathbb{R}).$$

The function β can be expressed in an explicit form if the parameters of the Blaschke-product are known. As a result, we can easily produce the inverse of a Blaschke-product on the torus [14]. Namely, for the function

$$\tau_k(t) := \beta^{-1}(t/n + 2k\pi/n) \quad (t \in \mathbb{R}, k = 0, \dots, n-1),$$

the following holds

$$B(e^{i\tau_k(t)}) = e^{in\beta(\beta^{-1}(t/n+2k\pi/n))} = e^{it}.$$

From the above, we get that $\varphi_k(1, t) := e^{i\tau_k(t)}$ ($t \in \mathbb{R}$) are the inverse branches of B on the torus.

In general, finding the inverse branches can be reduced to finding the roots of an n -th degree polynomial. Unfortunately, when considering this form of the problem, accurate geometric description of the results seems impossible. In the case of two factor Blaschke-products, the inverse branches can be expressed explicitly using the complex root function [8]. These results provide a basis for the description of the inverse branches for n factor Blaschke-products.

In this paper, we consider a description of the inverse branches and propose a numerical algorithm to produce them. We plan to extend these results to functions in \mathcal{A}_R in a future work, hence providing a numerical method to produce their Riesz-factorization. Notice, that when considering the inverse branches of B , for each $w = re^{it} \in \overline{\mathbb{D}}$, the equation $B(z) = w$ is equivalent to finding the roots of an n degree polynomial, therefore (considering multiplicities as well) has exactly n solutions. When grouping these solutions into n classes, there are several ways we can define (continuous) inverse branches. In Section 2, we shall prove the following theorem about the branches:

Theorem 1.1. *For all $t \in \mathbb{T}, t \notin T_0$ the equation $B(z) = re^{it}$ has a single, unique (continuous in r) solution $\varphi_k(r, t)$, for which*

$$(1.7) \quad \begin{aligned} B(\varphi_k(r, t)) &= re^{it}, \varphi_k(1, t) = e^{i\tau_k(t)} \\ (0 \leq r \leq 1, k &= 0, \dots, n-1), \end{aligned}$$

furthermore these solution curves are disjoint for $r > 0$ and their union provides the closed disk.

There are other ways of defining the inverse lines of the generalized root function as well. For example one could consider all trajectories ending in the same zero to be an inverse branch. Figure 2 depicts the inverse branches for a 3 pole Blaschke-product. In Figures (2b) and (2c), small circles show the zeros of B , while larger circles refer to the critical points. Figure (2a) shows the branches of the function $w \rightarrow \sqrt[3]{w}$, in which case $a_1 = \dots = a_3 = 0$, or in other words 0 is a root with a multiplicity of 3. Figure (2b) illustrates the inverse branches as defined in Theorem 1, while in (2c) a branch is made up of the trajectories ending in the same zero points. The dots show the positions of the inverse poles and critical points.




Figure 2: Classification of inverse branches

2. Differential equation form of the problem

The Blaschke-product [16] B has $2n - 2$ critical points which fall in the hyperbolic convex hull of the zeros. All other critical points fall outside \mathbb{D} and are the reflections of the ones in \mathbb{D} with respect to \mathbb{T} [16]. Let $R := \min\{1/|a| : a \in A\}$ and set of critical values by

$$K := B(A') := \{B(z) : z \in A'\}.$$

We show that if $w = re^{it_0} \notin K$ ($r \in [r_1, r_2]$) and $B(z_0) = w_0$, then the implicit equation

$$B(\varphi(r)) = w = re^{it_0}, \quad \varphi(r_0) = z_0 \quad (r \in [r_1, r_2] \subset I := (0, R))$$

is equivalent to an initial value problem corresponding to a first order differential equation [1]. Since for all $r \in [r_1, r_2]$, $w = re^{it_0} \notin K$, the function $B(z) = w$ has a locally unique inverse, piecing these together yields the differentiable solution $\varphi(r) = \varphi(r, t_0)$ ($r \in [r_1, r_2]$) of the implicit equation, for

which

$$(2.1) \quad B(\varphi(r)) = re^{it_0}, \quad \varphi'(r)B'(\varphi(r)) = e^{it_0} \quad (r \in [r_1, r_2]).$$

We note that if the roots of B have a multiplicity of 1, then the choice $r_1 = 0$ is also possible.

Let us introduce the function

$$f(z) := \frac{B(z)}{B'(z)} \quad (z \in \Omega := \mathbb{D}_R \setminus (A \cup A')).$$

We show that if $z_0 \in \Omega$, then the problem (2.1) is equivalent to the initial value problem for the function $\chi : I \rightarrow \Omega$, $I \subset (0, \infty)$:

$$(2.2) \quad \chi'(r) = \frac{f(\chi(r))}{r}, \quad \chi(r_0) = z_0 \quad (r_0 \in I, z_0 \in \Omega)$$

where $B(z_0) = w_0 = r_0 e^{it_0} \notin K$. Indeed, by (2.1)

$$\varphi'(r) = \frac{e^{it_0}}{B'(\varphi(r))} = \frac{B(\varphi(r))}{rB'(\varphi(r))},$$

therefore φ is a solution of (2.2). Conversely, for any χ solution of (2.2)

$$\frac{d}{dr} \log r = \frac{1}{r} = \chi'(r) \frac{B'(\chi(r))}{B(\chi(r))} = \frac{d}{dr} \log B(\chi(r))$$

holds, therefore $B(\chi(r)) = cr$ ($r \in (r_1, r_2)$), where by the initial values, for the constant c we get

$$r_0 e^{it_0} = B(z_0) = B(\chi(r_0)) = c \cdot r_0,$$

from which $c = e^{it_0}$ follows.

The right side of (2.2) fulfills the usual conditions for the existence and uniqueness of the solution, therefore for any $(r_0, z_0) \in I \times \Omega$, the differential equation (2.2) has a unique extended solution $\chi_{z_0}(r)$ ($r \in J \subset I$) to the boundary [1] (pp. 50–57). The ranges of these solutions will be referred to as trajectories henceforth. From the existence and uniqueness theorem, it follows that two trajectories are either disjoint or are the same. Furthermore, for any point in Ω there is a trajectory which passes through it, therefore taking the union of these trajectories yields Ω .

In order to describe the maximal solutions, let us introduce the partitioning of the line segments S_t induced by the finite set K . For

$$S_t \cap K := \{\rho_j e^{it} : j = 1, \dots, m_t\}$$

let $\rho_0 := 0, \rho_{m_t+1} := R$, furthermore let

$$\begin{aligned} \mathcal{J}_t &:= \{J_j := (\rho_{j-1}e^{it}, \rho_j e^{it}) : j = 1, \dots, m_t + 1\}, \\ \mathcal{J}_t &:= \{(0, R)\}, \text{ if } S_t \cap K = \emptyset. \end{aligned}$$


Denote line segments $J = (\rho e^{it}, \sigma e^{it}) \in \mathcal{J}_t$ by $\widehat{J} := (\rho, \sigma)$. Then the solutions $\varphi = \varphi_J$ ($J \in \mathcal{J}_t$) of the implicit equation

$$B(\varphi(r)) = e^{it}r \quad (r \in \widehat{J}, J \in \mathcal{J}_t, t \in \mathbb{I})$$


match the maximal solutions of (2.2).

Now we can describe the inverse branches in the following manner. For any $w = re^{it_0}, t_0 \notin T_0$ take the initial values $z_{0k} = e^{i\tau_k(t_0)}$ ($k = 0, \dots, n - 1$). We note that for these values $B(z_{0k}) = e^{it_0}$ holds. Using these initial values, let us consider the maximal solutions of (2.2) $\varphi_k = \chi_{z_{0k}}$. If the zeros have a multiplicity of one and $t_0 \notin T_0$, the domain of these solutions is $[0, R)$, furthermore the solutions match the trajectories from Theorem 1.

Figure 3 illustrates the partitioning of the trajectories. Figure (3a) shows the set $K = B(A')$ (for $|K| = 2$) and the partitioning of the critical line segments S_t ($t \in T_0$), which in this case consists of 2 segments. The inverse images of these are illustrated in Figure (3b).



(a) Critical lines S_t



(b) Inverse images of S_t and the trajectories

Figure 3: Critical lines and their inverse images

Taking nested circles instead of line segments, we get the following theorem.

Theorem 2.1. *Suppose that for the circle $C_r := \{re^{it} : t \in \mathbb{I}\}$ ($0 < r \leq 1$), $C_r \cap K = \emptyset$ holds. Let $w_0 = re^{it_0} \in C_r$ and denote by z_{0j} the inverse images of w_0 : $B(z_{0j}) = w_0$ ($j = 0, \dots, n - 1$). Then, the smooth solutions of the implicit equation*

$$(2.3) \quad B(\psi_j(t)) = re^{it} \quad (t \in \mathbb{I}), \quad \psi_j(t_0) = z_{0j}$$

exist uniquely.

Figures (4a) and (4b) show the nested circles and the corresponding inverse images.




Figure 4: Nested circles and the corresponding inverse images

3. Inverse algorithm

In this section we consider a numerical solution to the problem outlined above. The problem of finding the inverse of a Blaschke-product can be posed in three equivalent forms. Namely, the inverse images are solutions to the following implicit equation, differential equation and line integral problems.

- i) $B(z) = w = re^{it}$, $B(z_0) = w_0$ ($z = \varphi(r)$),
- ii) $\chi'(r) = f(\chi(r))/r$, $\chi(r_0) = z_0$ ($f(z) = B(z)/B'(z)$),
- iii) $F(z) = \int_{z_0}^z \frac{B'(\zeta)}{B(\zeta)} d\zeta$.

Any of these forms can be solved with various efficient numerical algorithms [10]. In this section we introduce an algorithm based on Newton's method which solves the implicit equation form of the problem. We generalize the findings of the previous section and examine the solutions of

$$(3.1) \quad B(z) = w \in \Gamma,$$

where $\Gamma \subset \mathbb{D}_R$, $\Gamma \cap K = \emptyset$ is a simple smooth curve. We show that starting from the initial solution $B(z_0) = w_0$, one can acquire a finite number of (z_k, w_k) ($k = 1, \dots, N$) solutions for which $(z, w) = (z_N, w_N)$ and z_{k+1} is a limit point of a Newton-iteration starting from z_k . These results are a consequence of the following theorem.

Theorem 3.1. *There exist numbers $r_0 > 0$, $r_1 > 0$, such that for any $B(z_0) = w_0$, $|z_0| \leq 1$ initial solution and any $w \in K_{r_1}(w_0) \cap \Gamma$ a unique $z \in K_{r_0}(z_0)$, $|z| \leq 1$, $B(z) = w$ solution exists of (3.1), which is the limit of the quadratically convergent sequence*

$$z_{j+1} = g(z_j), \quad (j \in \mathbb{N}),$$

where

$$(3.2) \quad g(z) := g_w(z) = (z - z_0) - \frac{B(z) - w}{B'(z)} \quad (z \in K_{r_0}(z_0)),$$

$$z = \lim_{j \rightarrow \infty} z_j.$$

Proof. We show the existence of $r > 0$ (independent of z_0), such that the mapping g is contractive on the disk $Z := \{z \in \mathbb{C} : |z - z_0| \leq r\}$:

$$g : Z \rightarrow Z, \quad |g(z_1) - g(z_2)| \leq \frac{1}{2}|z_1 - z_2| \quad (z_1, z_2 \in Z).$$

The function g is twice differentiable on $\overline{\mathbb{D}}_R$ and its derivatives satisfy

$$g'(z) = \frac{(B(z) - w)B''(z)}{(B'(z))^2},$$

$$g''(z) = \frac{B''(z)}{B'(z)} + \frac{(B(z) - w)B^{(3)}(z)}{(B'(z))^2} - 2 \frac{(B(z) - w)(B''(z))^2}{(B'(z))^3}.$$

First we prove that there exists $\rho_0 > 0$, independent of z_0 , so

$$(3.3) \quad w_0 = B(z_0) \in \Gamma \quad \Rightarrow \quad \forall \kappa \in K : |\kappa - z_0| \geq \rho_0.$$

Indeed, by the mean value theorem $B'(\kappa) = 0$, $M := \sup_{|z| \leq R} |B''(z)| < \infty$ we get

$$0 < d := \min_{\kappa \in K, w \in \Gamma} |\kappa - w| \leq |B(\kappa) - B(z_0)| \leq M|\kappa - z_0|^2,$$

from which (3.3) holds with the constant $\rho_0 := \sqrt{d/M}$. Let

$$H := \overline{\mathbb{D}}_R \setminus \bigcup_{\kappa \in K} K_{\rho_1}(\kappa) \quad (\rho_1 := \rho_0/2).$$

The functions g', g'' are bounded on H , furthermore $\overline{K}_{\rho_1}(z_0) \subset H$ holds. Let

$$m := \min_{z \in H} |B'(z)|, \quad M_1 := \max_{|z| \leq R} |g''(z)|.$$

Then by $M_1 < \infty$, $m > 0$ and the mean value theorem

$$(3.4) \quad |g'(z)| \leq |g'(z) - g'(z_0)| + |g'(z_0)| \leq$$

$$\leq M_1|z - z_0| + \frac{M}{m^2}|w_0 - w| < \frac{1}{2} \quad (z \in K_{\rho_1}(z_0))$$

holds if, for example

$$|w - w_0| \leq \frac{m^2}{4M}, \quad |z - z_0| \leq \frac{1}{2M_1}.$$

By the above, for $r = \min\{\rho_1, 1/(2M_1)\}$ the mapping g is contractive:

$$|g(z_1) - g(z_2)| \leq \frac{1}{2} |z_1 - z_2| \quad (z_1, z_2 \in \overline{K}_r(z_0)).$$

Since

$$(3.5) \quad \begin{aligned} |g(z) - z_0| &\leq |g(z) - g(z_0)| + |g(z_0) - z_0| \leq \\ &\leq \frac{1}{2} |z - z_0| + \frac{|w_0 - w|}{m}, \end{aligned}$$

choosing the parameters r, r_0, r_1 according to

$$(3.6) \quad \begin{aligned} r = r_0 &\leq \min\{\rho_1, 1/(2M_1)\}, \quad K_r(z_0) \subset K_R(0), \\ |w - w_0| &\leq r_1 := \min\{mr/2, m/(2M)\} \end{aligned}$$

the mapping $g : Z := \overline{K}_r(z_0) \rightarrow Z$ is indeed contractive, with a contraction coefficient of $1/2$.

By the fixed-point theorem the limit

$$z := \lim_{j \rightarrow \infty} z_j \in Z$$

exists and

$$g(z) = z \iff B(z) = w.$$

Finally, $|z| \leq 1$ holds, since $w \in \Gamma \subset \mathbb{D}$ and B maps \mathbb{D} onto itself. Since $g(z) = z$, the derivative $g'(z) = 0$ and by the mean value theorem

$$|z_{j+1} - z| = |g(z_j) - g(z)| \leq \max_{\zeta \in Z} |g''(\zeta)| \cdot |z_j - z|^2/2 \leq M_1 |z_j - z|^2/2,$$

meaning that the method is quadratically convergent, which concludes our proof. ■

We would like to draw attention to the fact, that in applications calculating the derivative function $B'(z)$ in (3.2) may pose some difficulty. In [15] the authors describe a method with which the derivatives of a polynomial can be calculated from its roots and the function value. The same idea can be easily extended for Blaschke-products and a simple formula for $B'(z)$ expressed with the function value and the poles can be given:

$$(3.7) \quad B'(z) = B(z) \cdot \left[\sum_{i=1}^n \frac{1}{z - a_i} + \sum_{i=1}^n \frac{\overline{a_i}}{1 - \overline{a_i}z} \right].$$

Suppose we have w_0, z_{0i} such that $B(z_{0i}) = w_0$, ($i = 1, \dots, n$). Then, z_{0i} are the roots of the polynomial

$$P(z) = \prod_{i=1}^n (z - a_i) - w_0 \prod_{i=1}^n (1 - \bar{a}_i z),$$

furthermore $P(z_{0i}) = B(z_{0i}) - w_0 = 0$. In a future work we hope to develop a method to calculate the derivative values from only the known z_{0i} , ($i = 1, \dots, n$) based on the above observations.

References

- [1] **Arnold, V.I.**, *Ordinary Differential Equation*, MIT Press Cambridge, Massachusetts and London, England, 1998.
- [2] **Blaschke, W.**, Eine Erweiterung des Satzes von Vitali über Folgen analytischer Funktionen, *Math. Phys. Kl. Sächs. Gessel. der Wiss.*, Leipzig, **67** (1915), 194–200.
- [3] **Bokor, J., F. Schipp and A. Soumelidis**, Pole structure estimation from Laguerre representation using hyperbolic metric on the unite disc, in: *50th IEEE Conf. on Decision and Control an European Control Conf.*, Orlando, Florida, December 12–15, 2011, 2136–2141.
- [4] **Bokor, J., F. Schipp and A. Soumelidis**, Applying hyperbolic wavelets in frequency domain identification, in: *Int. Conference in Control Automation and Robotics, ICINCO 2012*, Rome, Italy, July 28–30, 2012, 532–535.
- [5] **Bokor, J. and Z. Szabó**, *Non-euclidean Geometries in Modelling and Control*, Széchenyi University Press, Győr, 2015.
- [6] **Coifman, R.R. and J. Peyriere**, Phase unwinding, or invariant subspace decompositions of Hardy spaces, *J. of Fourier Anal. Appl.*, **25** (2019), 684–695.
- [7] **Coifman, R.R. and S. Steinerberger**, Nonlinear phase unwinding of functions, 2016. <https://arxiv.org/pdf/1508.01241.pdf>
- [8] **Dózsa, T. and F. Schipp**, Hyperbolic geometry and Blaschke functions, *Annales Univ. Sci. Budapest., Sect. Comp.*, **51** (2020), 59–68.
- [9] **Fridli, S. and F. Schipp**, Discrete rational biorthogonal systems on the disc, *Annales Univ. Sci. Budapest., Sect. Comp.*, **50** (2020), 127–134.
- [10] **Henrici, P.**, *Applied and Computational Complex Analysis, Vol. I, II*, John Wiley and Sons, New York, London, Sidney, Toronto, 1974.

- [11] **Heuberger, P., P. Van den Hof and B. Wahlberg**, *Modelling and Identification with Rational Orthogonal Basis Functions*, Springer, 2005.
- [12] **de Hoog, T.**, *Rational Orthonormal Bases and Related Transforms in Linear System Modeling*, Ponsen and Looijen, Netherlands, 2001. ISBN 90-6464-357-1
- [13] **Móricz, F.**, *Harmonic Analysis on the Disc* (Hungarian), Polygon, Szegedi Egyetemi Kiadó, 2013.
- [14] **Schipp, F.**, *Rational Orthogonal Systems* (Hungarian), Eötvös Loránd University, Faculty of Informatics, 2016.
<https://www.inf.elte.hu/karidigitaliskonyvtar>
- [15] **Sendov, Bl., A. Andreev and N. Kjurkchiev**, Numerical solution of polynomial equations, *Handbook of Numerical Analysis*, Vol. III, 629–777.
- [16] **Sheil-Small, T.**, *Complex Polynomials*, Cambridge Studies in Adv. Math. 73, Cambridge University Press.
- [17] **Tan, C., L. Zhang and H-T. Wu**, A novel Blaschke unwinding adaptive Fourier decomposition based signal Compression algorithm with application on ECG signals, 2018. <https://arxiv.org/pdf/1803.06441.pdf>

T. Dózsa and F. Schipp

Department of Numerical Analysis

Eötvös Loránd University

H-1117 Budapest

Pázmány Péter sétány 1/C

Hungary

dotuaai@inf.elte.hu

schipp@numanal.inf.elte.hu