

EXPANDING ON RESULTS OF WIRSING AND KLURMAN

Jean-Marie De Koninck (Québec, Canada)

Imre Kátai and Bui Minh Phong (Budapest, Hungary)

Communicated by Karl-Heinz Indlekofer

(Received July 20, 2022; accepted August 15, 2022)

Abstract. Building on two conjectures formulated by the second author in the 1980's regarding characterisations of multiplicative functions and their respective proofs by Wirsing in the late 1990's and by Klurman in 2017, we expand on the achievements of these two authors by proving new results.

1. Introduction

As is common, we let \mathbb{N} , \mathbb{R} , \mathbb{C} stand the sets of positive integers, real and complex numbers, respectively. Let also \mathcal{M} (resp. \mathcal{M}^*) stand for the set of complex-valued multiplicative (resp. completely multiplicative) functions. We say that $f \in \mathcal{M}_1$ (resp. \mathcal{M}_1^*), if $f \in \mathcal{M}$ (resp. \mathcal{M}^*) and $|f(n)| = 1$ for every $n \in \mathbb{N}$.

Let us first recall two classical conjectures due to the second author.

Conjecture 1. (Kátai [3]) *If $g \in \mathcal{M}$ and*

$$g(n+1) - g(n) \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

then either $g(n) \rightarrow 0$ as $n \rightarrow \infty$, or $g(n) = n^s$ for some complex number s with $\Re s < 1$.

Key words and phrases: Arithmetical function, multiplicative functions.

2010 Mathematics Subject Classification: primary 11A07, 11A25, secondary 11N25, 11N64.

Conjecture 2. (Kátai [3]) *If $g \in \mathcal{M}_1$ and*

$$\lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} |g(n+1) - g(n)| = 0 \quad \text{or} \quad \lim_{x \rightarrow \infty} \frac{1}{\log x} \sum_{n \leq x} \frac{|g(n+1) - g(n)|}{n} = 0,$$

then there exists a real number τ for which

$$g(n) = n^{i\tau} \quad \text{for all } n \in \mathbb{N}.$$

Conjecture 1 was proved by E. Wirsing in 1996. The proof is given in Wirsing, Yuansheng and Pintsung [8] as well as in Wirsing and Zagier [9].

A particular case of Conjecture 2 was proved by O. Klurman [4] in 2017 and we state it as follows.

Theorem A. (Klurman) *Conjecture 2 is true for $g \in \mathcal{M}_1^*$.*

Here, we expand on the above results of Wirsing and Klurman.

2. A generalisation of Wirsing's result

In a recent paper [1], we proved that if $f \in \mathcal{M}_1^*$ and if c_0, c_1, c_2 are three complex numbers satisfying $(c_0, c_1, c_2) \neq (0, 0, 0)$ for which

$$\lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} |c_0 f(n) + c_1 f(n+1) + c_2 f(n+2)| = 0,$$

then there exists a real number τ for which $f(n) = n^{i\tau}$ for all positive integers n .

In the same paper [1], we formulated the following conjecture.

Conjecture 3. *Given $f \in \mathcal{M}_1^*$ and complex numbers c_0, c_1, \dots, c_k such that $(c_0, c_1, \dots, c_k) \neq (0, 0, \dots, 0)$. Assuming that*

$$c_0 f(n) + c_1 f(n+1) + \dots + c_k f(n+k) \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

then there exists a real number τ for which $f(n) = n^{i\tau}$ for all positive integers n .

In [1], we also formulated the following weaker conjecture.

Conjecture 4. *Given $f \in \mathcal{M}_1^*$ and complex numbers c_0, c_1, \dots, c_k such that $(c_0, c_1, \dots, c_k) \neq (0, 0, \dots, 0)$. Assuming that*

$$\frac{1}{x} \sum_{n \leq x} |c_0 f(n) + c_1 f(n+1) + \dots + c_k f(n+k)| \rightarrow 0 \quad \text{as } x \rightarrow \infty,$$

then $c_0 + c_1 + \dots + c_k = 0$ and there exists a real number τ for which $f(n) = n^{i\tau}$ for all positive integers n .

Observe that in [1] we were able to prove Conjecture 4 in the particular cases $k = 1, 2$. However, we were unable to prove the more general case $k \geq 3$.

Now, consider the three operators E , I and Δ defined on the sequence $(x_n)_{n \in \mathbb{N}}$ by

$$E x_n = x_{n+1}, \quad I x_n = x_n, \quad \Delta x_n = E x_n - I x_n = x_{n+1} - x_n.$$

We also consider iterations of $\Delta f(n)$. For this, we let

$$\Delta^2 f(n) := \Delta \Delta f(n) = \Delta(f(n+1) - f(n)) = f(n+2) - 2f(n+1) + f(n),$$

and for an arbitrary integer $k \geq 3$, we let $\Delta^k f(n) := \Delta \Delta^{k-1} f(n)$.

Our purpose in this section and the next one is the following generalisation of the result of Wirsing mentioned in the Introduction.

Theorem 1. *Let $f \in \mathcal{M}_1^*$, set $s_f(n) := \Delta^7 f(n)$ and assume that $\lim_{n \rightarrow \infty} s_f(n) = 0$. Then, there exists a real number τ such that $f(n) = n^{i\tau}$ for all $n \in \mathbb{N}$.*

3. Proof of Theorem 1

In our recent paper, we proved (see Theorem 10 in [2]) that, assuming that

$$|\Delta^7 f(n)| \leq K := 12 - \delta,$$

then there exists some real number t such that $f(n) = n^{it} F(n)$ for all $n \in \mathbb{N}$, where $F^k(n) = 1$ for all $n \in \mathbb{N}$, and $|\Delta^7 F(n)| \leq K + \varepsilon$ provided $n \geq n_0(\varepsilon)$.

As we will see, Theorem 1 is an easy consequence of our Theorem 10 in [2]. Indeed, since we assumed that $s_f(n) \rightarrow 0$ as $n \rightarrow \infty$ and since $n^{i\tau} - (n+1)^{i\tau} \rightarrow 0$ as $n \rightarrow \infty$, it follows that $s_F(n) \rightarrow 0$ as $n \rightarrow \infty$. But since $s_F(n)$ can assume only finitely many distinct values, it follows that

$$s_F(n) = \Delta^7 F(n) = 0 \quad \text{for every large } n.$$

On the other hand, since the roots of the polynomial $(x-1)^7$ are $\omega_\ell := e^{2\pi i \ell / 7}$ ($i = 0, 1, \dots, 6$), we have that

$$(3.1) \quad F(m+n_0) = \sum_{\ell=0}^6 c_\ell \omega_\ell^m \quad (m = 0, 1, 2, \dots).$$

Observe that the right-hand side of (3.1) is periodic mod 7, that is, $F(n_1) = F(n_2)$ if $n_1 \equiv n_2 \pmod{7}$ with $n_1, n_2 \geq n_0$. From this, it follows that $F(7n) = F(7(n+1))$, and since $F(7) \neq 0$, this implies that $F(n) = 1$ for all positive integers n , thereby completing the proof of Theorem 1. \blacksquare

In concluding this topic, we formulate yet another conjecture.

Conjecture 5. *Let $f \in \mathcal{M}_1^*$ and assume that*

$$\lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} |\Delta^\tau f(n)| = 0.$$

Then, there exists a real number τ for which $f(n) = n^{i\tau}$ for every $n \in \mathbb{N}$.

4. Consequences of Klurman's theorem

We now move to some interesting consequences of Klurman's theorem, namely by establishing four new results.

Theorem 2. *Assume that $a, b, c \in \mathbb{N}$ and $d \in \mathbb{C} \setminus \{0\}$. Then functions $g_1 \in \mathcal{M}_1$ and $g_2 \in \mathcal{M}_1$ satisfy the condition*

$$\lim_{x \rightarrow \infty} \frac{1}{\log x} \sum_{n \leq x} \frac{|g_1(an + b) - dg_2(cn)|}{n} = 0$$

if and only if there are functions $G_1, G_2 \in \mathcal{M}_1$ and a number $\tau \in \mathbb{R}$ such that

$$g_1(n) = n^{i\tau} G_1(n), \quad g_2(n) = n^{i\tau} G_2(n)$$

and

$$G_1(an + b) - d \left(\frac{c}{a} \right)^{i\tau} G_2(cn) = 0$$

are satisfied for every $n \in \mathbb{N}$.

Theorem 3. *Assume that $a, b, c \in \mathbb{N}$ and $d \in \mathbb{C} \setminus \{0\}$. Then functions $g_1 \in \mathcal{M}_1$ and $g_2 \in \mathcal{M}_1$ satisfy the condition*

$$\lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} |g_1(an + b) - dg_2(cn)| = 0$$

if and only if there are functions $G_1, G_2 \in \mathcal{M}_1$ and a number $\tau \in \mathbb{R}$ such that

$$g_1(n) = n^{i\tau} G_1(n), \quad g_2(n) = n^{i\tau} G_2(n)$$

and

$$G_1(an + b) - d \left(\frac{c}{a} \right)^{i\tau} G_2(cn) = 0$$

are satisfied for every $n \in \mathbb{N}$.

Theorem 4. *Assume that $a > 0$, $b, A > 0$, $B \in \mathbb{N}$ are integers with $(a, A) = 1$, $\Delta = Ab - aB \neq 0$ and $C \in \mathbb{C} \setminus \{0\}$. If $g \in \mathcal{M}_1^*$ satisfies the condition*

$$\lim_{x \rightarrow \infty} \frac{1}{\log x} \sum_{n \leq x} \frac{|g(an + b) - Cg(An + B)|}{n} = 0,$$

then there is a real number τ such that $g(n) = n^{i\tau}$ for every $n \in \mathbb{N}$.

Theorem 5. *Assume that $a > 0$, $b, A > 0$, $B \in \mathbb{N}$ are integers with $(a, A) = 1$, $\Delta = Ab - aB \neq 0$ and $C \in \mathbb{C} \setminus \{0\}$. If $g \in \mathcal{M}_1^*$ satisfies the condition*

$$\lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} |g(an + b) - Cg(An + B)| = 0,$$

then there is a real number τ such that $g(n) = n^{i\tau}$ for every $n \in \mathbb{N}$.

5. Proofs of Theorems 2 through 5

For the proofs of Theorems 2 and 3, we will be using the following result.

Theorem B. (Phong [6], Theorems 1 and 2) *Assume that $a, b, c \in \mathbb{N}$ and $d \in \mathbb{C} \setminus \{0\}$. Then functions $g_1 \in \mathcal{M}_1$ and $g_2 \in \mathcal{M}_1$ satisfy the condition either*

$$\lim_{x \rightarrow \infty} \frac{1}{\log x} \sum_{n \leq x} \frac{|g_1(an + b) - dg_2(cn)|}{n} = 0$$

or

$$\lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} |g_1(an + b) - dg_2(cn)| = 0$$

if and only if there are functions $g^ \in \mathcal{M}_1^*$ and $G_1, G_2 \in \mathcal{M}_1$ such that*

$$g_1(n) = g^*(n)G_1(n), \quad g_2(n) = g^*(n)G_2(n),$$

$$G_1(an + b) - d \left(\frac{g^*(c)}{g^*(a)} \right) G_2(cn) = 0$$

and either

$$(5.1) \quad \lim_{x \rightarrow \infty} \frac{1}{\log x} \sum_{n \leq x} \frac{|g^*(n+1) - g^*(n)|}{n} = 0$$

or

$$(5.2) \quad \lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} |g^*(n+1) - g^*(n)| = 0$$

are satisfied for every $n \in \mathbb{N}$.

Now, we infer from Theorem A that the conditions (5.1)–(5.2) imply that there is a number $\tau \in \mathbb{R}$ such that $g^*(n) = n^{i\tau}$ for every $n \in \mathbb{N}$.

Hence, Theorems 2 and 3 are proved.

In similar way, we prove Theorem 4. This time, we will be using the following result.

Theorem C. (Phong [7], Theorem 1) *Assume that $a > 0$, $b, A > 0$, $B \in \mathbb{N}$ be integers with $(a, A) = 1$, $\Delta = Ab - aB \neq 0$ and $C \in \mathbb{C} \setminus \{0\}$. If $g \in \mathcal{M}_1^*$ satisfies the condition*

$$\lim_{x \rightarrow \infty} \frac{1}{\log x} \sum_{n \leq x} \frac{|g(an + b) - Cg(An + B)|}{n} = 0$$

then

$$(5.3) \quad \lim_{x \rightarrow \infty} \frac{1}{\log x} \sum_{n \leq x} \frac{|g(n+1) - g(n)|}{n} = 0.$$

We also infer from Theorem A that the condition (5.3) implies that there is a number $\tau \in \mathbb{R}$ such that $g(n) = n^{i\tau}$ for every $n \in \mathbb{N}$, thereby completing the proof of Theorem 4.

We now move to the proof of Theorem 5. By adopting the method used in the proof of Theorem 1 in [7], we have the following result.

Theorem D. *Assume that $a > 0$, $b, A > 0$, $B \in \mathbb{N}$ be integers with $(a, A) = 1$, $\Delta = Ab - aB \neq 0$ and $C \in \mathbb{C} \setminus \{0\}$. If $g \in \mathcal{M}_1^*$ satisfies the condition*

$$\lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} |g(an + b) - Cg(An + B)| = 0,$$

then

$$(5.4) \quad \lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} |g(n+1) - g(n)| = 0.$$

The condition (5.4) with Theorem A implies the proof of Theorem 5.

References

- [1] **De Koninck, J.-M., I. Kátai and B.M. Phong**, Characterising some triplets of completely multiplicative functions, *Ann. Univ. Sci. Budapest., Sect. Comput.* **50** (2020), 101–112.

- [2] **De Koninck, J.-M., I. Kátai and B.M. Phong**, On the variations of completely multiplicative functions at consecutive arguments, *Publicationes Mathematicae Debrecen*, **98(1-2)** (2021), 219–230.
- [3] **Kátai, I.**, Some problems in number theory, *Studia Sci. Math. Hungar.*, **16(3-4)** (1983), 289–295.
- [4] **Klurman, O.**, Correlations of multiplicative functions and applications, *Compositio Math.*, **153(8)** (2017), 1620–1657.
- [5] **Phong, B.M.**, A characterization of some arithmetical multiplicative functions, *Acta Math. Hungar.*, **63(1)** (1994), 29–43.
- [6] **Phong, B.M.**, A characterization of some unimodular multiplicative functions, *Publ. Math. Debrecen*, **57(3-4)** (2000), 339–366.
- [7] **Phong, B.M.**, A characterization of the some unimodular multiplicative functions II, *Ann. Univ. Sci. Budapest., Sect. Comput.*, **21** (2004), 187–194.
- [8] **Wirsing, E., T. Yuansheng and S. Pintsung**, On a conjecture of Kátai for additive functions, *J. Number Theory*, **56** (1996), 391–395.
- [9] **Wirsing, E. and D. Zagier**, Multiplicative functions with difference tending to zero, *Acta Arith.*, **100** (2001), 75–78.

J.-M. De Koninck

Département de mathématiques

Université Laval

Québec

Québec G1V 0A6

Canada

jmdk@mat.ulaval.ca

I. Kátai and B. M. Phong

Department of Computer Algebra

Faculty of Informatics

Eötvös Loránd University

H-1117 Budapest, Pázmány Péter sétány 1/C

Hungary

katai@inf.elte.hu, bui@inf.elte.hu

