

ON APPROXIMATION OF HYDROLOGICAL FUNCTIONS

by

P. VÉRTESI

(Budapest)

(Received July 4, 1979)

1. Introduction

In their paper [1] L. Góczán and A. F. Szász introduced some hydrological functions of new type to investigate the agronomical and technical economy of water-supplies.

In many cases they had to determine a good approximating polynomial for the hydrological functions $f(x)$ under the following conditions:

(a) $f'(x)$ is continuous (say) in $[-1, 1]$ (i.e., $f' \in C$).

(b) One can measure the data $f(1)$, $f'(1)$ and

$$f(x_k) \quad (k = 1, 2, \dots, n, \quad -1 \leq x_k < 1, \quad x_k \neq x_j \text{ if } k \neq j).$$

As an easy calculation shows the uniquely determined polynomial $G_n(f, x)$ of degree $\leq n+1$ satisfying the conditions

(a) and (b) has the form

$$(1.1) \quad G_n(f, x) = \sum_{k=1}^n f(x_k) l_k(x) \frac{(x-1)^2}{(x_k-1)^2} + f(1) \frac{\Omega(x)}{\Omega(1)} \left[1 - \frac{\Omega'(1)}{\Omega(1)} (x-1) \right] + f'(1) \frac{\Omega(x)}{\Omega(1)} (x-1)$$

where

$$(1.2) \quad \begin{cases} \Omega(x) = \Omega_n(x) = c_n \prod_{k=1}^n (x-x_k) & (c_n \neq 0), \\ l_k(x) = l_{kn}(x) = \frac{\Omega(x)}{\Omega'(x_k)(x-x_k)} & (k = 1, \dots, n). \end{cases}$$

A very natural question (which was raised in [1], too) is how to choose the nodes $x_k = x_{kn}$ ($k = 1, 2, \dots, n$; $n = 1, 2, \dots$) to ensure that

$$(1.3) \quad \lim_{n \rightarrow \infty} \|G_n(f, x) - f(x)\| = 0 \quad \text{whenever} \quad f' \in C.$$

Here, as usual, $\|g\| = \max_{-1 \leq x \leq 1} |g(x)|$ for $g \in C$.

2. Results

2.1. It is easy to see that using the equidistant nodes $x_{kn} = -1 + k(n+1)^{-1}$ (which were applied in [1]), the "Lebesgue constant" of the process (1.1)

$$\lambda_n = \|\lambda_n(x)\| \stackrel{\text{def}}{=} \left\| \sum_{k=1}^n |l_k(x)| \frac{(x-1)^2}{(x_k-1)^2} + \frac{|\Omega(x)|}{|\Omega(1)|} \left| 1 - \frac{\Omega'(1)}{\Omega(1)} (x-1) \right| + \left| \frac{\Omega(x)}{\Omega(1)} (x-1) \right| \right\|$$

very rapidly tends to infinity with n , from where we shall obtain, that for the equidistant nodes (1.3) generally does not hold (see 3.6).

But considering the Jacobi nodes, i.e. the roots

$$(2.1) \quad -1 < x_{nn}^{(\alpha, \beta)} < x_{n-1, n}^{(\alpha, \beta)} < \dots < x_{1n}^{(\alpha, \beta)} < 1 \quad (\alpha, \beta > -1)$$

of the polynomial $P_n^{(\alpha, \beta)}(x)$ of degree n defined by

$$(1-x)^\alpha (1+x)^\beta P_n^{(\alpha, \beta)}(x) = \frac{(-1)^n}{2^n n!} \left(\frac{d}{dx} \right)^n [(1-x)^{n+\alpha} (1+x)^{n+\beta}]$$

we can state as follows

THEOREM 2.1. *If $3 < \alpha = \beta + 4 \leq 4.5$, for the $G_n(\alpha, \beta)$ (f, x) = $G_n(f, x)$ process defined on the nodes (2.1) the relation (1.3) is valid. More exactly, if $\omega(f', t)$ denotes the modulus of continuity of $f'(x) \in C$, then*

$$(2.2) \quad \|G_n(\alpha, \beta)(f, x) - f(x)\| = \begin{cases} O\left(\frac{\ln n}{n}\right) \omega\left(f', \frac{1}{n}\right) & \text{if } 3 < \alpha \leq 3.5, \\ O(n^{\alpha-4.5}) \omega\left(f', \frac{1}{n}\right) & \text{if } \alpha > 3.5. \end{cases}$$

2.2. Here we state, the Theorem 2.1, in certain sense, is not far from the best possible one. This can be formulated as follows.

THEOREM 2.2. *If $\alpha = \beta + 4$ and $\alpha > 4.5$, then one can construct a function $f_1(x)$ such that $f_1'(x) \in C$ and*

$$(2.3) \quad \lim_{n \rightarrow \infty} \|G_n(\alpha, \beta)(f_1, x) - f_1(x)\| = \infty.$$

2.3. Similar results can be obtained for $\beta = 1$. We omit the details.

3. Proofs

3.1. We shall use the following relations (sometimes omitting the superfluous notations).

$$(3.1) \quad P_n^{(\alpha, \beta)}(x) = (-1)^n P_n^{(\beta, \alpha)}(-x).$$

if $x_{kn} = x_{kn}^{(\alpha, \beta)} = \cos \vartheta_{kn}^{(\alpha, \beta)}$, $x_{0n} = 1$, $x_{n+1, n} = -1$,

$$(3.2) \quad \vartheta_{k+1, n}^{(\alpha, \beta)} - \vartheta_{kn}^{(\alpha, \beta)} \sim \frac{1}{n} \quad (k = 0, 1, \dots, n)$$

moreover, if $x_{j(n), n}$ is the nearest root to x ($1 \leq j \leq n$) then

$$(3.3) \quad |x - x_{kn}| \sim \frac{|j^2 - k^2|}{n^2} \quad (k \neq j; k = 0, 1, 2, \dots, n+1),$$

$$(3.4) \quad |P_n^{(\alpha, \beta)}(x_{kn})| \sim k^{-\alpha-3/2} n^{\alpha+2} \quad (0 < \vartheta_k \leq \pi - \varepsilon),$$

$$(3.5) \quad |P_n^{(\alpha, \beta)}(x)| \sim |x - x_j| \vartheta_j^{-\alpha-3/2} n^{1/2} \sim |\vartheta - \vartheta_j| \vartheta_j^{-\alpha-1/2} n^{1/2}$$

uniformly for $\vartheta \in [0, \pi - \varepsilon]$.

At these formulae $x = \cos \vartheta$, the „ \sim ” depends on α and β (see [2], (1.1), (4.1.3), (8.9.2) and [3], [4]).

Denoting by $Q_n(x) = Q_n(f, x)$ the polynomial of degree $\leq n$ for which

$$(3.6) \quad |f^{(i)}(x) - Q_n^{(i)}(x)| = O(1) \left(\frac{\sin \vartheta}{n} \right)^{1-i} \omega \left(f', \frac{\sin \vartheta}{n} \right) \quad (i = 0, 1)$$

(see [5]) we can write by (1.1)

$$(3.7) \quad \begin{aligned} G_n(f, x) - f(x) &= G_n(f, x) - Q_n(f, x) + Q_n(f, x) - f(x) = \\ &= G_n(f - Q_n, x) + Q_n(f, x) - f(x) = \\ &= \sum_{k=1}^n [f(x_k) - Q_n(x_k)] l_k(x) \left(\frac{x-1}{x_k-1} \right)^2 + O(1) \frac{\sin \vartheta}{n} \omega \left(f', \frac{\sin \vartheta}{n} \right). \end{aligned}$$

3.2. For the sum, using the formulae (3.1)–(3.6), we can write if $x \geq 0$

$$\begin{aligned} \sum_{k=1}^n \dots &= O(1) \left[\sum_{k=1}^n \frac{\sin \vartheta_k}{n} \omega \left(\frac{\sin \vartheta_k}{n} \right) \left| \frac{P_n(x)}{P'_n(x_k)(x - x_k)} \right| \left(\frac{x-1}{x_k-1} \right)^2 + \right. \\ &+ \frac{\sin \vartheta_j}{n} \omega \left(\frac{\sin \vartheta_j}{n} \right) \frac{\vartheta_j^{-\alpha-3/2} n^{1/2}}{j^{-\alpha-3/2} n^{\alpha+2}} \left. \right] = O(1) \left[\sum_{k=1}^{\left[\frac{3n}{4} \right]} \frac{\vartheta_k}{n} \omega \left(\frac{\vartheta_k}{n} \right) \cdot \right. \\ &\cdot \frac{\vartheta_j^{-\alpha-1/2} n^{-1/2} n^2}{k^{-\alpha-3/2} n^{\alpha+2} |k+j| |k-j|} \frac{j^4 n^{-4}}{k^4 n^{-4}} + O(1) \sum_{k=1}^{\left[\frac{n}{4} \right]} \frac{\vartheta_k}{n} \omega \left(\frac{\vartheta_k}{n} \right) \cdot \\ &\cdot \frac{\vartheta_j^{-\alpha-1/2} n^{-1/2}}{k^{-\beta-3/2} n^{\beta+2}} \frac{j^4}{n^4} \stackrel{\text{def}}{=} S_1 + S_2 \end{aligned}$$

where Σ' means that $k \neq j$.

$$\begin{aligned} S_1 &= O(1) \frac{1}{n^2} \sum_{k=1}^n{}' \omega\left(\frac{k}{n^2}\right) \frac{j^{3,5-\alpha} k^{\alpha-1,5}}{|k+j||k-j|} = \\ &= O(1) \left(n^{-2} \sum_{k < \frac{j}{2}} \dots + n^{-2} \sum_{\frac{j}{2} \leq k \leq 2j}{}' \dots + n^{-2} \sum_{k > 2j} \right) \stackrel{\text{def}}{=} O(1)(I_1 + I_2 + I_3). \end{aligned}$$

Now

$$I_1 = O(n^{-2}) j^{1,5-\alpha} \omega\left(\frac{1}{n}\right) \sum_{k=1}^j k^{\alpha-1,5}$$

$$\text{so } I_1 = O(n^{-1}) \omega\left(\frac{1}{n}\right) \text{ if } \alpha > 3.$$

For I_2 we can write

$$I_2 = O(n^{-2}) j \omega\left(\frac{1}{n}\right) \sum_{k=j+1}^{2j} (k-j)^{-1} = O\left(\frac{\ln n}{n}\right) \omega\left(\frac{1}{n}\right).$$

As I_3 , one can estimate as follows

$$I_3 = O(n^{-2}) j^{3,5-\alpha} \sum_{k=2j}^n \omega\left(\frac{k}{n^2}\right) k^{\alpha-3,5}$$

$$\text{which is } O\left(\frac{1}{n}\right) \omega\left(\frac{1}{n}\right) \text{ if } 3 < \alpha \leq 3,5 \text{ and } O(n^{\alpha-4,5}) \omega\left(\frac{1}{n}\right) \text{ if } \alpha > 3,5.$$

Let us estimate S_2 . We have

$$S_2 = O(1) \frac{j^{3,5-\alpha}}{n^{8-\alpha-\beta}} \sum_{k=1}^n \omega\left(\frac{k}{n^2}\right) k^{\beta+2,5} = O(1) \frac{j^{3,5-\alpha}}{n^4} \sum_{k=1}^n \omega\left(\frac{k}{n^2}\right) k^{\alpha-1,5}$$

$$\begin{aligned} &\text{which can be estimated by } O\left(\frac{1}{n}\right) \omega\left(\frac{1}{n}\right) \text{ if } 3 < \alpha \leq 3,5 \text{ and by } O(n^{\alpha-4,5}). \\ &+ \omega\left(\frac{1}{n}\right) \text{ if } \alpha > 3,5. \end{aligned}$$

3.3. Let now $x < 0$. Then the corresponding expression with $t = n+1-j$ are as follows

$$\begin{aligned} S_1^* &= O(1) \frac{1}{n^2} \sum_{k=1}^n{}' \omega\left(\frac{k}{n^2}\right) \frac{t^{-\beta-1/2} k^{\beta+2,5}}{|k+t||k-t|}, \\ S_2^* &= O(1) \frac{t^{-\beta-1/2}}{n^{\alpha-\beta}} \sum_{k=1}^n \omega\left(\frac{k}{n^2}\right) k^{\alpha-1,5}, \end{aligned}$$

i.e. S_1^* and S_2^* has the same form as S_1 and S_2 , respectively, if we consider $\alpha = \beta + 4$. These estimations give Theorem 2.1.

3.4. To prove Theorem 2.2, we apply the following statement, which is a special case of [6], **Theorem 3.1.**

If for the sequence of linear operators $T_n(f, x)$ ($f \in C$) and the functions $g_n(x)$ ($n = 1, 2, \dots$) we have

$$(a1) \quad g_n(x) \in C,$$

$$(a2) \quad T_n(g_n, z_n) \equiv c_n \mu_n(z_n) \text{ for certain } \{z_n\} \subset [-1, 1] \text{ (definition of } \mu_n(z_n));$$

$$(B^*) \quad \tilde{f}(x) \stackrel{\text{def}}{=} \sum_{i=1}^{\infty} e_{n_i} g_{n_i}(x) \in C \text{ for any } \{n_i\} \subset \{p_r\}$$

where $\{p_r\}$ is a certain fixed sequence of indices and $0 < e_n \leq e_{n+1} \leq 1$;

$$(C^*) \quad c_5 \mu_n(z_n) > \sum_{i=k+1}^{\infty} e_{n_i} |T_{n_k}(g_{n_i}, z_{n_k})| + \sum_{i=k}^{\infty} e_{n_i} |(g_{n_i} z_{n_k})|$$

($k = 1, 2, \dots$; $0 < C_4 < C_5$ for arbitrary $\{n_i\} \subset \{p_r\}$; then with a suitable c_8 and $\{s_i\} \subset \{p_r\}$ with $f_1(x) = c_8 \tilde{f}(x)$)

$$(3.8) \quad T_n(f_1, z_n) - f_1(z_n) > e_n \mu_n(z_n) \quad (n = s_1, s_2, \dots).$$

(Here c_i are fixed constants.)

3.5. Let $\alpha - 4,5 = \varepsilon > 0$, $z_4 = (1 + x_{1n})/2$ and

$$(3.9) \quad g_n(x_{kn}) = \begin{cases} \frac{\sin \vartheta_k}{n} \omega_1 \left(\frac{\sin \vartheta_k}{n} \right) \operatorname{sign} l_k(z_n) & \text{if } -1 < x_k \leq 0, \\ 0 & \text{if } k = n+1 \text{ or } x_k > 0, \end{cases}$$

where $\omega_1(t)$ is a modulus of continuity.

In the interval (x_{k+1}, x_k) let $g_n(x)$ be the Hermite interpolatory polynomial of degree ≤ 3 for which

$$g'_n(x_{kn}) = g'_n(x_{k+1, n}) = 0 \quad (k = 0, 1, \dots, n).$$

Let $T_n = G_n$. By (3.9) and (1.1)

$$G_n(g_n, z_n) = \sum_{-1 < x_k \leq 0} \frac{\sin \vartheta_k}{n} \omega_1 \left(\frac{\sin \vartheta_k}{n} \right) |l_k(z_n)| \left(\frac{z_n - 1}{x_k - 1} \right)^2 \stackrel{\text{def}}{=} \mu_n(z_n).$$

By (3.1)–(3.5) we obtain, as at the estimation of S_2

$$(3.10) \quad \mu_n(z_n) \sim \frac{1}{n^4} \sum_{k=1}^n \omega_1 \left(\frac{k}{n^2} \right) k^{x-1,5} \equiv c \omega_1 \left(\frac{1}{n^2} \right) n^\varepsilon \quad (n = 1, 2, \dots)$$

i.e. we can suppose $\mu_n(z_n) \nearrow \infty$ for a „bad” $\omega_1(t)$. Let us see now (B*). By definition it is easy to verify, that $\omega(g'_n, t) \sim \omega_1(t)$, i.e. supposing $\sum_{i=1}^{\infty} e_{p_i} < \infty$, we obtain for any $\{n_i\} \subset \{p_r\}$

$$(3.11) \quad \omega(\tilde{f}', t) \leq c \sum_{i=1}^{\infty} e_{n_i} \omega(g'_{n_i}, t) = 0(\omega_1(t)),$$

which is more than (B*).

To prove (C*), we remark, that $|g_n(x)| \leq 1/n$. So, if we choose the sequence $\{p_i\}$ such that

$$\frac{\lambda_{p_k}}{p_{k+1}} \leq 1 \quad (k = 1, 2, \dots)$$

we obtain (C*).

So by (3.8)–(3.11)

$$G_n(f_1, z_n) - f_1(z_n) > e_n \omega_1\left(\frac{1}{n^2}\right) n^\varepsilon \quad (n = s_1, s_2, \dots)$$

which, with a suitable $\{e_i\}$, is more than (2.3).

3.6. For the equidistant nodes one can use analogous argument, using the fact that for certain $\{k_n\}$ $\|l_{k_n, n}(x)\| > (1,5)^{n^2}$ ($n \geq n_0$) (see e.g. [7] Part 3, Chapter 2, § 3). We omit the further details.

REFERENCES

- [1] L. Góczán, A. F. Szász, Approximations of hydrological functions, Acta Sci. Hungar. Inst. Geogr. **19** (1970), 233–260 (Hungarian).
- [2] G. Szegő, Orthogonal polynomials, AMS Coll. Publ. Vol. XXIII (New York, 1959).
- [3] G. I. Natanson, Two-sided estimate for the Lebesgue function of the Lagrange interpolation with Jacobi nodes, Izv. Vyss. Ucebn. Zaved., Matematika, **11** (1967), 67–74 (Russian).
- [4] P. Vértesi, On Lagrange interpolation, Period. Math. Hungar., (1981).
- [5] I. E. Gopengauz, On a theorem of A. F. Timan, Mat. Zametki, **1(2)** (1967), 163–172 (Russian).
- [6] P. Vértesi, On certain linear operators, VIII, Acta Math. Acad. Sci. Hungar. **25** (1974), 171–187, 449–450.
- [7] I. P. Natanson, Constructive Theory of Functions, Gostehizdat, 1949 (Russian).

Mathematical Institute of the Hungarian Academy of Sciences
1053 Budapest, Reáltanoda u. 13–15.
Hungary