

ON DOMAIN OF ATTRACTION OF THE DOUBLE EXPONENTIAL DISTRIBUTION*

JÁNOS C. FODOR

1. Introduction

For a sequence of independent, identically distributed random variables X_1, X_2, \dots with common distribution function (d.f.) $F(x)$ and for $n \geq 1$, let

$$Z_n = \max(X_1, X_2, \dots, X_n).$$

Then, obviously,

$$P(Z_n < x) = F^n(x).$$

We say that F belongs to the domain of attraction of a nondegenerate d.f. H , if there exist sequences of constants $\{a_n\}_{n=1}^{\infty}$, $\{b_n\}_{n=1}^{\infty}$ with $b_n > 0$, such that

$$(1) \quad \lim_{n \rightarrow \infty} F^n(a_n + b_n x) = H(x)$$

holds at all continuity points of H . The relation (1) will be denoted by $F \in D(H)$.

The limit laws for Z_n were fully characterized in [4]. They are the so-called extreme value distributions. We employ the notation $H_{1,\gamma}(x) = \exp(-x^{-\gamma})$, $x > 0$, $H_{2,\gamma}(x) = \exp(-(-x)^\gamma)$, $x < 0$

* This work has been partially supported by OTKA-1049.

and $H_{3,0}(x) = \exp(-e^{-x})$, $-\infty < x < +\infty$, where γ is a positive parameter. A d.f. F can belong only to the domain of attraction of one of the three types $H_{1,\gamma}$, $H_{2,\gamma}$ and $H_{3,0}$ (see [4] or [3], for example).

Necessary and sufficient conditions for $F \in D(H)$ are well-known (see [5] and references above). In a recent paper [2] we gave characterizations of domains of attraction of extreme value distributions in terms of ultimately convexity (concavity) properties of some functions $[1 - G(x)]^q$ ($q \in \mathbf{R} \setminus \{0\}$), where G is an appropriate d.f. depending only on F . This approach seems to be new.

In [4] Gnedenko remarked that the characterization of $H_{3,0}$ (the double exponential distribution) cannot be regarded as final and simple enough. In this paper we show that each d.f. F in the domain of attraction of $H_{3,0}$ is asymptotic to some twice differentiable d.f. G , where $[1 - G(x)]^q$ is ultimately convex for every $q \in \mathbf{R} \setminus \{0\}$. This result is closely related to the Theorem of [1].

2. The result

For a given d.f. F , let

$$\omega(F) = \sup\{x; F(x) < 1\}.$$

We say that the function $[1 - F(x)]^q$ ($q \in \mathbf{R} \setminus \{0\}$ is fixed) is ultimately convex if for the given q there exists an $x(q) < \omega(F)$ such that $[1 - F(x)]^q$ is convex on the interval $(x(q), \omega(F))$.

Theorem. $F \in D(H_{3,0})$ if and only if there exists a twice differentiable d.f. G such that

- a.) $\omega(G) = \omega(F) = \omega$
- b.) $[1 - G(x)]^q$ is ultimately convex for every $q \in \mathbf{R} \setminus \{0\}$
- c.) $\lim_{x \uparrow \omega} \frac{1 - F(x)}{1 - G(x)} = 1. \quad \square$

The proof will be based on the following lemmas.

Lemma 1. For a given twice differentiable d.f. G , $[1 - G(x)]^q$

is ultimately convex for every $q \in \mathbf{R} \setminus \{0\}$, if and only if

$$\lim_{x \uparrow \omega} \left[\frac{1 - G(x)}{G'(x)} \right]' = 0,$$

where commas denote derivatives. \square

Proof. The statement follows directly from the Proposition of [2]. \square

The basic idea we use to construct the d.f. G is due to Balkema and de Haan, see [1]. Assume that $F \in D(H_{3,0})$. Let $\bar{F}_0(x) = 1 - F(x)$ and

$$F_{k+1}(x) = \max(0, 1 - \int_x^\omega \bar{F}_k(t) dt)$$

for $k = 0, 1, 2$ and $\bar{F}_k(x) = 1 - F_k(x)$. It follows from [5] that the integrals in the definition of $F_{k+1}(x)$ are finite and $F_{k+1}(x)$ is a d.f. also for $k = 0, 1, 2$ with $\omega(F_{k+1}) = \omega(F) = \omega$. In [5] it was proved also that if $F \in D(H_{3,0})$ then $F_{k+1} \in D(H_{3,0})$ ($k = 0, 1, 2$) and the following relation is valid:

$$(2) \quad \lim_{x \uparrow \omega} \frac{\bar{F}_k(x) \cdot \bar{F}_{k+2}(x)}{\{\bar{F}_{k+1}(x)\}^2} = 1, \quad k = 0, 1.$$

Let us define the function $A(x)$ by

$$(3) \quad A(x) = \frac{\{\bar{F}_2(x)\}^3}{\{\bar{F}_3(x)\}^2} \quad \text{for } x \in (x_0, \omega),$$

where $x_0 < \omega$ is such that $\bar{F}_3(x_0) > 0$.

Lemma 2. If $F \in D(H_{3,0})$ then there exists an x_1 such that $x_1 \in (x_0, \omega)$ and $A(x)$ is strictly decreasing on the interval (x_1, ω) .

\square

Proof. It is obvious that

$$(4) \quad A'(x) = A(x) \cdot \frac{\bar{F}_2(x)}{\bar{F}_3(x)} \cdot \left[-3 \cdot \frac{\bar{F}_1(x) \cdot \bar{F}_3(x)}{\{\bar{F}_2(x)\}^2} + 2 \right].$$

By relation (2) it follows that there exists an $x_1 \in (x_0, \omega)$ such that $A'(x) < 0$ for $x \in (x_1, \omega)$. \square

Lemma 3. *If $F \in D(H_{3,0})$ then $\{A(x)\}^q$ is ultimately convex for every $q \in \mathbb{R} \setminus \{0\}$.* \square

Proof. It will be sufficient to show that

$$\lim_{x \uparrow \omega} \left[\frac{A(x)}{A'(x)} \right]' = 0.$$

From (4) we get

$$\begin{aligned} \frac{A(x)}{A'(x)} &= \frac{\bar{F}_3(x) [\bar{F}_2(x)]^{-1}}{2 - 3 \cdot \bar{F}_1(x) \cdot \bar{F}_3(x) \cdot [\bar{F}_2(x)]^{-2}}, \text{ whence} \\ \left[\frac{A(x)}{A'(x)} \right]' &= \frac{-1 + \bar{F}_3(x) \cdot \bar{F}_1(x) \cdot [\bar{F}_2(x)]^{-2}}{2 - 3 \cdot \bar{F}_1(x) \cdot \bar{F}_3(x) \cdot [\bar{F}_2(x)]^{-2}} - \\ &- 3 \cdot \frac{-\bar{F}_0(x) \cdot [\bar{F}_3(x)]^2 \cdot [\bar{F}_2(x)]^{-3} - \bar{F}_1(x) \cdot \bar{F}_3(x) \cdot [\bar{F}_2(x)]^{-2}}{\{2 - 3 \cdot \bar{F}_1(x) \cdot \bar{F}_3(x) \cdot [\bar{F}_2(x)]^{-2}\}^2} \\ &+ \frac{2 \cdot [\bar{F}_1(x)]^2 \cdot [\bar{F}_3(x)]^2 \cdot [\bar{F}_2(x)]^{-4}}{\{2 - 3 \cdot \bar{F}_1(x) \cdot \bar{F}_3(x) \cdot [\bar{F}_2(x)]^{-2}\}^2}. \end{aligned}$$

By relation (2) the denominators tend to $-1, 1, 1$, respectively, and the numerators tend to $0, -2, 2$, respectively. This completes the proof. \square

Lemma 4. *Assume that $F \in D(H_{3,0})$. Then*

$$(5) \quad \lim_{x \uparrow \omega} \frac{1 - F(x)}{A(x)} = 1.$$

Proof. By (3) we have that

$$\frac{1 - F(x)}{A(x)} = \frac{\bar{F}_0(x) \cdot \bar{F}_2(x)}{[\bar{F}_1(x)]^2} \cdot \left[\frac{\bar{F}_1(x) \cdot \bar{F}_3(x)}{[\bar{F}_2(x)]^2} \right]^2.$$

So by (2) we get the relation (5). \square

COROLLARY. If $F \in D(H_{3,0})$ then $\lim_{x \uparrow \omega} A(x) = 0$. \square

Proof. The statement is evident from (5). \square

Let us turn to the proof of our Theorem. The "if" part is a consequence of Theorem 2.7.2. of [3], Lemma 2.5 of [6] and our Lemma 1.

Assume now that $F \in D(H_{3,0})$ and let

$$(6) \quad G(x) = \begin{cases} \max(0, 1 - A(x)) & \text{for } x > x_1 \\ 0 & \text{for } x \leq x_1. \end{cases}$$

$G(x)$ is a d.f. because of Lemma 2 and the Corollary. It is obvious by (3) and (6) that $\omega(G) = \omega(F)$. The part b.) follows from Lemma 3 and Lemma 1. The part c.) is a direct consequence of Lemma 4. This completes the proof. \square

References

- [1] BALKEMA, A.A. and de HAAN L., On R. von Mises' condition for the domain of attraction of $\exp(-e^{-x})$. *The Annals of Mathematical Statistics*, **43** (1972) 1352-1354.
- [2] FODOR, J.C., On domains of attraction of extreme value distributions via generalized concavity - convexity. To appear in : *Annales Univ. Sci. Budapest, Sect. Comp.*

- [3] GALAMBOS, J., The Asymptotic Theory of Extreme Order Statistics. John Wiley and Sons, New York, (1978).
- [4] GNEDENKO, B.V., Sur la distribution limite du terme maximum d'une série aléatoire. *Ann. of Math.*, **44**(1943), 423-453.
- [5] de HAAN L., A form of regular variation and its application to the domain of attraction of the double exponential distribution. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, **17**(1971), 241-258.
- [6] RESNICK, S.I., Tail equivalence and its applications. *J. Appl. Probability*, **8**(1971), 136-156

(Received June 24, 1987)

JÁNOS C. FODOR

*Computer Center, Eötvös Loránd University
H-1502 Budapest, 112, P.O.Box 157*

HUNGARY