

A MATKOWSKI–SUTÔ–TYPE PROBLEM FOR WEIGHTED QUASI–ARITHMETIC MEANS

Z. Daróczy and Zs. Páles (Debrecen, Hungary)

*Dedicated to the 60th birthday of
Professor Karl-Heinz Indlekofer*

Abstract. The aim of this paper is to solve the functional equation

$$\begin{aligned} \lambda\varphi^{-1}(\lambda\varphi(x) + (1-\lambda)\varphi(y)) + (1-\lambda)\psi^{-1}(\lambda\psi(x) + (1-\lambda)\psi(y)) = \\ = \lambda x + (1-\lambda)y, \end{aligned}$$

where φ, ψ are strictly monotone continuous real functions defined on an open real interval I and $\lambda \in]0, 1[$ is a fixed number. The case $\lambda = \frac{1}{2}$ has recently been completely solved by the authors in [6]. The main result of the paper offers a complete solution for the case $\lambda \neq \frac{1}{2}$ and it states that if $\lambda \neq \frac{1}{2}$ then φ, ψ are solutions of the above equation if and only if there exist constants a, b, c, d with $ac \neq 0$ such that $\varphi(x) = ax + b$ and $\psi(x) = cx + d$ for all $x \in I$.

1. Introduction

Let $I \subset \mathbb{R}$ be a nonvoid open interval. A function $M : I^2 \rightarrow I$ is called a *strict mean* on I if it is continuous and $\min\{x, y\} < M(x, y) < \max\{x, y\}$ for

This research has been supported by the Hungarian Scientific Research Fund (OTKA) grant T043080 and by the Higher Education, Research and Development Fund (FKFP) grant 0215/2001.

all $x, y \in I$, $x \neq y$. Let $M_i : I^2 \rightarrow I$ ($i = 1, 2$) be strict means. For any fixed $x, y \in I$, we define the Gauss-iteration in the following way

$$\begin{aligned} x_1 &:= x, & y_1 &:= y, \\ x_{n+1} &:= M_1(x_n, y_n), & y_{n+1} &:= M_2(x_n, y_n) \quad (n \in \mathbb{N}). \end{aligned}$$

It is known ([1],[6]) that, for any $x, y \in I$, the limit $M_3(x, y) = \lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} y_n$ exists, and $M_1 \otimes M_2 := M_3 : I^2 \rightarrow I$ is a strict mean on I called the *Gauss-composition* of M_1 and M_2 .

Denote by $\mathcal{CM}(I)$ the class of continuous and strictly monotone functions defined on the interval I . A function $M : I^2 \rightarrow I$ is called a *weighted quasi-arithmetic mean* on I if there exist $0 < \lambda < 1$ and $\varphi \in \mathcal{CM}(I)$ such that

$$(1.1) \quad M(x, y) = \varphi^{-1}(\lambda\varphi(x) + (1 - \lambda)\varphi(y)) =: A_\varphi(x, y; \lambda)$$

for all $x, y \in I$ (see [8], [15], [6]). The number λ in (1.1) is called the *weight* and the function φ is said to be the *generating function*. Let $0 < \lambda < 1$ be a fixed number and M_i ($i = 1, 2, 3$) be *weighted quasi-arithmetic means* on I with the same weight λ . Our main concern is to find conditions so that

$$(1.2) \quad M_3 = M_1 \otimes M_2$$

be an identity on I^2 . In the particular case $\lambda = \frac{1}{2}$ we have recently determined all the solutions in full generality in [6].

In order to solve the problem (1.2), we need to study the functional equation

$$(1.3) \quad \lambda A_\varphi(x, y; \lambda) + (1 - \lambda) A_\psi(x, y; \lambda) = \lambda x + (1 - \lambda)y \quad (x, y \in I),$$

where $\varphi, \psi \in \mathcal{CM}(I)$ are unknown functions. The case $\lambda = \frac{1}{2}$ is called the Matkowski-Sutô problem (cf. [16], [17], [11], [3], [4], [5]). When $\lambda \neq \frac{1}{2}$, the continuously differentiable solutions of (1.3) were determined in [7].

Our approach is analogous to that of followed in [6]. First we prove certain regularity properties of the functions satisfying (1.3). Based on this and also applying the extension theorem known from [2], we then obtain the complete solution of the problem described above.

2. The locally Lipschitz property of solutions

Let $0 < \lambda < 1$ and let $\varphi, \psi \in \mathcal{CM}(I)$ be solutions for (1.3). Our aim is to prove that $\varphi, \varphi^{-1}, \psi, \psi^{-1}$ are locally Lipschitz functions on their domains.

Definition 2.1. Let $J \subset \mathbb{R}$ be a nonvoid open interval and $f : J \rightarrow \mathbb{R}$. We say that the function f is *locally Lipschitz* in J if, for any $u_0 \in J$, there exist constants $\delta > 0$ and $L > 0$ such that $U :=]u_0 - \delta, u_0 + \delta[\subset J$ and, for all $u, v \in U$,

$$|f(u) - f(v)| \leq L|u - v|.$$

The following theorem plays an important role in our investigations.

Theorem 2.2. *Let $f : J \rightarrow \mathbb{R}$ ($J \subset \mathbb{R}$ is a nonvoid open interval) be a strictly monotone increasing and continuous function such that, for all $v \in J$, the map*

$$u \longmapsto f(u) - f(\lambda u + (1 - \lambda)v) \quad (u \in J)$$

is strictly monotone increasing. Then f and its inverse f^{-1} are Lipschitz functions on their domains J and $f(J)$, respectively.

Proof. For the case $\lambda = \frac{1}{2}$, the proof can be found in [6]. In the more general case $0 < \lambda < 1$ (including also the case $\lambda = \frac{1}{2}$) the result follows from a more general result stated in [14, Theorem 3].

Theorem 2.3. *Let $0 < \lambda < 1$ and $\varphi, \psi \in \mathcal{CM}(I)$ be solutions for the functional equation (1.3). Then $\varphi, \varphi^{-1}, \psi, \psi^{-1}$ are locally Lipschitz functions on their domains.*

Proof. It is sufficient to prove the statement for the functions φ, φ^{-1} because the role of functions φ and ψ can be interchanged. Applying (1.3) with the substitutions $u = \varphi(x)$, $v = \varphi(y)$ ($u, v \in J := \varphi(I)$) we deduce the equation

$$\begin{aligned} (2.1) \quad & (1 - \lambda)\psi^{-1}(\lambda\psi \circ \varphi^{-1}(u) + (1 - \lambda)\psi \circ \varphi^{-1}(v)) = \\ & = \lambda\varphi^{-1}(u) + (1 - \lambda)\varphi^{-1}(v) - \lambda\varphi^{-1}(\lambda u + (1 - \lambda)v) \end{aligned}$$

for all $u, v \in J$. Without loss of generality, we can assume that φ and ψ are strictly *increasing* functions. Then, for each fixed v , the left hand side of (2.1) is strictly increasing function of u , which results that the right hand side of (2.1) should also be strictly increasing in u . Therefore, for each fixed $v \in J$,

$$u \longmapsto \varphi^{-1}(u) - \varphi^{-1}(\lambda u + (1 - \lambda)v) \quad (u \in J)$$

is a strictly increasing function. Hence, in virtue of Theorem 2.2, φ^{-1} and φ are locally Lipschitz functions on J and on $I = \varphi^{-1}(J)$, respectively.

Corollary 2.4. *If $\varphi, \psi \in \mathcal{CM}(I)$ are solutions of (1.3) and φ (or ψ) is differentiable at a point $x_0 \in I$, then $\varphi'(x_0) \neq 0$ (or $\psi'(x_0) \neq 0$).*

Proof. See Corollary 4.4 of Theorem 4.3 in [6].

3. Differentiability of solutions

Suppose that $\varphi, \psi \in \mathcal{CM}(I)$ are *increasing* functions satisfying equation (1.3). Let

$$x := \varphi^{-1}(t + (1 - \lambda)s), \quad y := \varphi^{-1}(t - \lambda s),$$

where $t \in J := \varphi(I)$ and

$$s \in \left(\frac{J - t}{1 - \lambda} \right) \cap \left(\frac{t - J}{\lambda} \right) := J_{t,\lambda}$$

are arbitrary elements. Then equation (1.3) yields that, for any $t \in J := \varphi(I)$ and for any $s \in J_{t,\lambda}$,

$$(3.1) \quad \begin{aligned} \lambda \varphi^{-1}(t) &= \lambda \varphi^{-1}(t + (1 - \lambda)s) + (1 - \lambda) \varphi^{-1}(t - \lambda s) - \\ &\quad - (1 - \lambda) \psi^{-1}[\lambda h(t + (1 - \lambda)s) + (1 - \lambda)h(t - \lambda s)], \end{aligned}$$

where $h := \psi \circ \varphi^{-1}$.

Definition 3.1. Let $f : J \rightarrow \mathbb{R}$ be an arbitrary function and $0 < \lambda < 1$ be fixed. An element $t \in J$ is said to be a *point of λ -symmetry* for f , in notation $t \in \sigma_\lambda(f)$, if the identity

$$(3.2) \quad \lambda f(t + (1 - \lambda)s) + (1 - \lambda)f(t - \lambda s) = f(t)$$

holds true for all $s \in J_{t,\lambda}$.

Lemma 3.2. *If $F : J \rightarrow \mathbb{R}$ is a continuous function then $\sigma_\lambda(F)$ is closed in J .*

Proof. The proof of this obvious statement is analogous to that of [6, Lemma 4.6] concerning the case $\lambda = \frac{1}{2}$.

Lemma 3.3. *Let $0 < \lambda < 1$ and $\varphi, \psi \in \mathcal{CM}(I)$ be solutions of (1.3). Then $\sigma_\lambda(h) = \sigma_\lambda(\varphi^{-1})$, where $h := \psi \circ \varphi^{-1}$.*

Proof. We have that, for any $t \in J := \varphi(I)$ and $s \in J_{t,\lambda}$, (3.1) holds.

If $t \in \sigma_\lambda(h)$ then, by (3.1), we obtain

$$\lambda\varphi^{-1}(t) = \lambda\varphi^{-1}(t + (1 - \lambda)s) + (1 - \lambda)\varphi^{-1}(t - \lambda s) - (1 - \lambda)\psi^{-1} \circ h(t),$$

and since $\psi^{-1} \circ h(t) = \varphi^{-1}(t)$ holds, it follows that $t \in \sigma_\lambda(\varphi^{-1})$.

Conversely, if $t \in \sigma_\lambda(\varphi^{-1})$, then, by (3.1),

$$\varphi^{-1}(t) = \psi^{-1}(\lambda h(t + (1 - \lambda)s) + (1 - \lambda)h(t - \lambda s)),$$

whence we obtain that $t \in \sigma_\lambda(h)$.

Theorem 3.4. *If $0 < \lambda < 1$ and $\varphi, \psi \in \mathcal{CM}(I)$ are solutions of (1.3) then φ^{-1} is differentiable at any point $t_0 \in J \setminus \sigma_\lambda(\varphi^{-1})$.*

Proof. Without loss of generality we can assume that φ and ψ are increasing functions. If $J \setminus \sigma_\lambda(\varphi^{-1}) \neq \emptyset$ then let $t_0 \in J \setminus \sigma_\lambda(\varphi^{-1})$ be arbitrary. For an arbitrary function $g : J_{t_0,\lambda} \rightarrow \mathbb{R}$ denote by N_g the set of points $s \in J_{t_0,\lambda}$ at which g is not differentiable. Define the following functions

$$g_1(s) := \varphi^{-1}(t_0 + (1 - \lambda)s),$$

$$g_2(s) := \varphi^{-1}(t_0 - \lambda s),$$

$$g_3(s) := h(t_0 + (1 - \lambda)s),$$

$$g_4(s) := h(t_0 - \lambda s)$$

for all values $s \in J_{t_0,\lambda}$. Since φ^{-1} and h are strictly monotone functions, therefore, by Lebesgue's theorem on the almost everywhere differentiability of monotone functions, each N_{g_i} ($i = 1, 2, 3, 4$) is a null set, that is, the set

$$N := \bigcup_{i=1}^4 N_{g_i} \subset J_{t_0,\lambda}$$

is of measure zero. Since $t_0 \notin \sigma_\lambda(\varphi^{-1})$ thus, by Lemma 3.3, $t_0 \notin \sigma_\lambda(h)$. Therefore, the function

$$h_{t_0}(s) := \lambda h(t_0 + (1 - \lambda)s) + (1 - \lambda)h(t_0 - \lambda s) \quad (s \in J_{t_0,\lambda})$$

is not constant, which yields that its image is a proper interval $H_0 := h_{t_0}(J_{t_0,\lambda})$.

Let the set C be defined in the following way:

$$C := \{u \in H_0 \mid \psi^{-1} \text{ is not differentiable at } u\}.$$

Then, by Lebesgue's theorem, C is a null set. Therefore $H_0 \setminus C$ is a set of *positive* measure. Define D as follows:

$$D := h_{t_0}^{-1}(H_0 \setminus C) \subseteq J_{t_0, \lambda}.$$

Then $h_{t_0}(D) = H_0 \setminus C$. If D were a null set, then $h_{t_0}(D)$ would also be a null set since by Theorem 2.2, h_{t_0} is a locally Lipschitz function. Therefore D is a set of positive measure in $J_{t_0, \lambda}$. This implies that $D \setminus N$ is also a set of positive measure, hence $D \setminus N$ is not empty. Let $s_0 \in D \setminus N$ be arbitrarily fixed. Then g_i is differentiable at s_0 ($i = 1, 2, 3, 4$) and ψ^{-1} is differentiable at $h_{t_0}(s_0)$. Then, by (3.1), the equation

$$(3.3) \quad \begin{aligned} \lambda \varphi^{-1}(t) &= \lambda \varphi^{-1}(t + (1 - \lambda)s_0) + (1 - \lambda) \varphi^{-1}(t - \lambda s_0) - \\ &\quad - (1 - \lambda) \psi^{-1}(\lambda h(t + (1 - \lambda)s_0) + (1 - \lambda)h(t - \lambda s_0)) \end{aligned}$$

holds for all $t \in J$ such that $s_0 \in J_{t, \lambda}$ is also valid. This latter set of values of t is an open interval containing t_0 . Thus, ψ^{-1} is differentiable at $(t_0 + (1 - \lambda)s_0)$ and at $(t_0 - \lambda s_0)$; h is differentiable at $(t_0 + (1 - \lambda)s_0)$ and at $(t_0 - \lambda s_0)$, and ψ^{-1} is differentiable at $h_{t_0}(s_0)$, therefore, by the chain rule, the expression on the right side of (3.2) is differentiable at t_0 . Thus, we obtain that φ^{-1} is differentiable at t_0 .

Applying the previous result, we obtain the following important regularity theorem for the solutions of (1.3).

Theorem 3.5. *If $0 < \lambda < 1$ and $\varphi, \psi \in \mathcal{CM}(I)$ are solutions of (1.3), then there exists a nonvoid open interval $K \subset I$ on which φ and ψ are differentiable and $\varphi'(x) \neq 0$, $\psi'(x) \neq 0$ for all $x \in K$.*

Proof. Consider the function $\varphi^{-1} : J \rightarrow I$, where $J := \varphi(I)$. Then there are two possible cases:

- (i) either $\sigma_\lambda(\varphi^{-1}) = J$, that is, every $t \in J$ is a point of λ -symmetry for φ^{-1} ;
- (ii) or $\sigma_\lambda(\varphi^{-1}) \neq J$, that is, φ^{-1} has a point of non- λ -symmetry in J .

In case (i), for all $t \in J$ and $s \in J_{t, \lambda}$,

$$\varphi^{-1}(t) = \lambda \varphi^{-1}(t + (1 - \lambda)s) + (1 - \lambda) \varphi^{-1}(t - \lambda s)$$

holds. Since φ^{-1} is continuous, we have $\varphi^{-1}(u) = Au + B$ (where $A \neq 0$ and B are constants) for $u \in J$. This implies that $A_\varphi(x, y; \lambda) = \lambda x + (1 - \lambda)y$ for all $x, y \in I$, hence, by (1.3), $A_\psi(x, y; \lambda) = \lambda x + (1 - \lambda)y$ holds for all $x, y \in I$. Thus, ψ is also an affine function, therefore φ and ψ are differentiable functions with non-vanishing derivatives.

In case (ii), there exists $t_0 \notin \sigma_\lambda(\varphi^{-1})$. With the notation $G := \{t \in J \mid t \notin \sigma_\lambda(\varphi^{-1})\}$, due to Lemma 3.2, we have that G is a *nonvoid open set*. Thus, by Theorem 3.4, there exists a nonvoid open interval $\Delta \subset G \subset J$ such that φ^{-1} is differentiable on Δ . Hence φ is differentiable on some nonvoid open interval $K_0 \subset I$ and Corollary 2.4 implies that $\varphi'(x) \neq 0$ if $x \in K_0$. Now let us restrict equation (1.3) to the interval K_0 . Then the role of the functions φ and ψ can be interchanged and, by a similar argument, we obtain that there exists a nonvoid open interval $K \subset K_0 \subset I$ on which ψ is differentiable and $\psi'(x) \neq 0$ if $x \in K$. This completes the proof of the existence of the desired subinterval.

4. Continuous differentiability of solutions

If $\varphi, \psi \in \mathcal{CM}(I)$ are differentiable solutions of (1.3) with non-vanishing derivatives then (since the functions φ^{-1} and ψ^{-1} have the Darboux's property) we can assume that $\varphi'(x) > 0$ and $\psi'(x) > 0$ for every $x \in I$ without loss of generality.

Lemma 4.1. *If $0 < \lambda < 1$ is a fixed number and $\varphi, \psi \in \mathcal{CM}(I)$ are solutions of (1.3), moreover, φ and ψ are differentiable functions on I and $\varphi'(x) > 0$, $\psi'(x) > 0$ if $x \in I$ then, with the notation*

$$(4.1) \quad J := \varphi(I), \quad f := \varphi' \circ \varphi^{-1}, \quad g := \psi' \circ \varphi^{-1},$$

the functions $f, g : J \rightarrow \mathbb{R}_+$ satisfy the functional equation

$$(4.2) \quad f(\lambda u + (1 - \lambda)v)(g(v) - g(u)) = \lambda(f(u)g(v) - f(v)g(u))$$

for all $u, v \in J$.

Proof. Let us differentiate the functional equation (1.3) first with respect to x and then with respect to y . The conditions of the lemma ensure the differentiability, and we get the equations

$$\lambda \frac{\lambda \varphi'(x)}{\varphi'(A_\varphi(x, y; \lambda))} + (1 - \lambda) \frac{\lambda \psi'(x)}{\psi'(A_\psi(x, y; \lambda))} = \lambda$$

and

$$\lambda \frac{(1 - \lambda) \varphi'(y)}{\varphi'(A_\varphi(x, y; \lambda))} + (1 - \lambda) \frac{(1 - \lambda) \psi'(y)}{\psi'(A_\psi(x, y; \lambda))} = 1 - \lambda$$

for all $x, y \in I$. Multiplying the first equation by $(1 - \lambda)\psi'(y)$, the second by $\lambda\psi'(x)$, and subtracting the new equations from each other, we obtain

$$\frac{\lambda(\varphi'(x)\psi'(y) - \varphi'(y)\psi(x))}{\varphi'[A_\varphi(x, y; \lambda)]} = \psi'(y) - \psi'(x)$$

for all $x, y \in I$. Let $u = \varphi(x)$, $v = \varphi(y)$ ($u, v \in J := \varphi(I)$) be arbitrary then with the notations of (4.1), we obtain equation (4.2).

Definition 4.2. We say that $h : J \rightarrow \mathbb{R}_+$ is an element of the set $\mathcal{D}(J)$ if $h = d \circ c$, where $c \in \mathcal{CM}(J)$ and $d : I := c(J) \rightarrow \mathbb{R}_+$ is a derivative function, that is there exists a differentiable function $D : I \rightarrow \mathbb{R}_+$, such that $D'(x) = d(x)$ for all $x \in I$.

According to the previous definition, the functions f and g involved in the functional equation (4.2) are elements of the set $\mathcal{D}(J)$.

Theorem 4.3. *If the functions $f, g \in \mathcal{D}(J)$ satisfy the functional equation (4.2) for all $u, v \in J$ (where $0 < \lambda < 1$ is fixed), then there exists a nonvoid open interval $J_0 \subset J$ on which f is continuous.*

Proof. (i) If there exists a nonvoid open interval $J_0 \subset J$ on which f is continuous then the statement is true. If there exists a nonvoid open interval $J_0 \subset J$ on which g is constant then let $g(t) =: k$ for $t \in J_0$. Substituting arbitrary values $u, v \in J_0 (\subset J)$ into (4.2), we get that $f(u)k - f(v)k = 0$ for all $u, v \in J_0$. Hence, f must be constant on J_0 and consequently, f is continuous on J_0 .

Therefore, we may assume that $f, g \in \mathcal{D}(J)$ and that f and g are not constants on *any* nonvoid open subinterval $J_0 \subset J$. Denote by $\mathcal{D}_0(J)$ the set of functions in $\mathcal{D}(J)$ which are not constant on any proper subinterval of J .

(ii) Suppose that $f, g \in \mathcal{D}_0(J)$ satisfy (4.2) for all $y, v \in J$. Define the set $C(g)$ by

$$C(g) := \{t \mid t \in J, g \text{ is continuous at } t\}.$$

Then $g = d \circ c$, where c is continuous and strictly monotone, d is a derivative function; therefore g is continuous at each point $t \in J$ for which d is continuous at the point $c(t)$. Since the derivative function d is of Baire class 0 or 1, thus, according to Baire's theorem ([12], [13], [10]), the set of all points at which d is continuous is a *dense set of type G_δ* in $c(J)$, whence, because c is continuous and strictly monotone, $C(g)$ is also a dense G_δ set in J .

Now we will show that there exist points $u_0, v_0 \in C(g)$ such that $g(u_0) \neq g(v_0)$. Contrary to our assumption, suppose that $g(t) = k$ for every $t \in C(g)$, where $k > 0$ is a constant. Then substituting the values $u, v \in C(g)$ into (4.2), we obtain

$$f(u)k - f(v)k = 0,$$

whence $f(t) = l$ follows for every $t \in C(g)$, where $l > 0$ is a constant.

Because of the property of the set $C(g)$, for all $u \in J$, there exists $v \in C(g)$ such that $\lambda u + (1 - \lambda)v \in C(g)$. Thus, by (4.2),

$$l[k - g(u)] = \lambda(f(u)k - lg(u))$$

for every $u \in J$. This implies

$$f(u) = \frac{l(\lambda - 1)g(u) + lk}{\lambda k} \quad \text{if } u \in J.$$

If we substitute this form of the function f back into equation (4.2), after some calculations, we get

$$(4.3) \quad (k - g(\lambda u + (1 - \lambda)v))(g(v) - g(u)) = 0$$

for all $u, v \in J$.

Now let $v_0 \in J$ be fixed such that $c := g(v_0) \neq k$ holds. (Such a v_0 exists since g is non-constant.)

On the other hand, for any $t \in J$ and for any $\varepsilon > 0$ satisfying $]t - \varepsilon, t + \varepsilon[\subset J$, there exists $u \in]t - \varepsilon, t + \varepsilon[\subset J$ such that

$$g(\lambda u + (1 - \lambda)v_0) \neq k.$$

This last statement is valid because g is non-constant on any proper subinterval. Thus, by (4.3) it is obvious that $g(u) = g(v_0) = c$ holds. So in any neighborhood of *any* point $t \in J$ there exists u such that $g(u) = c$ and there exists s such that $g(s) = k \neq c$ which yields that g is not continuous anywhere and it is a contradiction.

(iii) We have proved in the previous part (ii) that there exist points $u_0, v_0 \in C(g)$ such that

$$g(u_0) \neq g(v_0)$$

holds. Then there exist a neighborhood $U \subset J$ of u_0 and a neighborhood $V \subset J$ of v_0 such that for any $u \in U$ and $v \in V$ we have $g(u) \neq g(v)$. Hence, by (4.2),

$$f(\lambda u + (1 - \lambda)v) = \lambda \frac{f(u)g(v) - f(v)g(u)}{g(v) - g(u)}$$

follows for all $u \in U$ and $v \in V$. This implies

$$(4.4) \quad f(t) = \lambda \frac{f\left(\frac{t - (1 - \lambda)v}{\lambda}\right)g(v) - f(v)g\left(\frac{t - (1 - \lambda)v}{\lambda}\right)}{g(v) - g\left(\frac{t - (1 - \lambda)v}{\lambda}\right)}$$

for every pair of values

$$(t, v) \in S := \{(t, v) \mid v \in V, t \in \lambda U + (1 - \lambda)v\},$$

where $t_0 := \lambda u_0 + (1 - \lambda)v_0 \in J$. By (4.4) and in view of Járαι's theorem ([9, Theorem 3.3]), we obtain that f is *continuous* in a neighborhood of the point $t_0 \in J$, that is, there exists a nonvoid open interval $t_0 \in J_0 \subset J$ on which f is continuous.

Finally we can state the following regularity theorem.

Theorem 4.4. *Let $0 < \lambda < 1$ and $\varphi, \psi \in \mathcal{CM}(I)$ be solutions of the functional equation (1.3). Then there exists a nonvoid open interval $K \subset I$ such that φ, ψ are continuously differentiable on K and $\varphi'(x) \neq 0$, $\psi'(x) \neq 0$ if $x \in K$.*

Proof. In virtue of Theorem 3.5, there exists a nonvoid open interval $K_1 \subset I$ on which φ and ψ are differentiable with non-vanishing derivatives. We can assume that $\varphi'(x) > 0$ and $\psi'(x) > 0$ if $x \in K_1$. Then by Lemma 4.1, with the notation of (4.1), we obtain that (4.2) holds, where $f, g \in \mathcal{D}(K_1)$. Thus, by Theorem 4.3, we obtain that there exists a nonvoid open interval $J_0 \subset J$ on which f is *continuous*. It means that $f := \varphi' \circ \psi^{-1} : J \rightarrow \mathbb{R}_+$ is continuous in J_0 . Consequently, $K_2 := \varphi^{-1}(J_0) \subset K_1 \subset I$ is a nonvoid open interval and

$$\varphi'(x) = \varphi' \circ \varphi^{-1}(s) = f(s) = f \circ \varphi(x)$$

for all $x \in K_2$. Hence φ' is *continuous* on the nonvoid open interval $K_2 \subset I$.

It is obvious that $\varphi, \psi \in \mathcal{CM}(K_2)$ and φ, ψ satisfy the functional equation (1.3) in K_2 , where φ is continuously differentiable on K_2 and $\varphi'(x) > 0$ if $x \in K_2$. Now apply our previous results for ψ . Then there exists a nonvoid open interval $K \subset K_2$ such that ψ is continuously differentiable on K and $\psi'(x) > 0$ if $x \in K$. Thus the statement of the theorem holds on the interval K .

5. Solution for the problem

The case $\lambda = \frac{1}{2}$ (which is the original Matkowski-Sutô problem) was treated and completely solved in [6]. Therefore, it remains to consider the case $0 < \lambda < 1$ and $\lambda \neq \frac{1}{2}$ only. Then the following statement holds.

Theorem 5.1. *Let $0 < \lambda < 1$ and $\lambda \neq \frac{1}{2}$. If $\varphi, \psi \in \mathcal{CM}(I)$ are solutions of the functional equation (1.3) then there exist constants $a, b, c, d \in \mathbb{R}$, $ac \neq 0$ such that*

$$(5.1) \quad \varphi(x) = ax + b \quad \text{and} \quad \psi(x) = cx + d$$

for all $x \in I$. Then

$$(5.2) \quad A_\varphi(x, y; \lambda) = A_\psi(x, y; \lambda) = \lambda x + (1 - \lambda)y$$

for all $x, y \in I$.

Proof. In virtue of the Theorem 4.4, there exists a nonvoid open interval $K \subset I$ such that φ and ψ are continuously differentiable on K and their derivatives do not vanish. Then, according to [7], (5.1) holds in K . Due to the extension theorem of the paper [2], (5.1) holds for all $x \in I$. This immediately yields that (5.2) is also true.

Now we are going to examine the solution of the general Matkowski-Sutô problem stated in the introduction for the class of weighted quasi-arithmetic means.

Theorem 5.2. *If $M_i : I^2 \rightarrow I$ ($i = 1, 2, 3$) are weighted quasi-arithmetic means with some weight λ ($0 < \lambda < 1; \lambda \neq \frac{1}{2}$) on I , then the identity $M_3 = M_1 \otimes M_2$ holds on I^2 if and only if there exists $f \in \mathcal{CM}(I)$ such that $M_i(x, y) = A_f(x, y; \lambda)$ ($i = 1, 2, 3$) holds for all $x, y \in I$.*

Proof. There exist generating functions $f_1, f_2, f_3 \in \mathcal{CM}(I)$ such that the invariance equation (c.f. [6], [1])

$$(5.3) \quad A_{f_3}(x, y; \lambda) = A_{f_3}(A_{f_1}(x, y; \lambda), A_{f_2}(x, y; \lambda); \lambda)$$

holds for all $x, y \in I$. Thus with the notations $u := f_3(x)$, $v := f_3(y)$, ($u, v \in f_3(I) =: J$), $\varphi := f_1 \circ f_3^{-1}$, $\psi := f_2 \circ f_3^{-1}$, (5.3) holds if and only if $\varphi, \psi \in \mathcal{CM}(J)$ satisfy the functional equation (1.3) for all $u, v \in J$. Then, by Theorem 5.1, we get that

$$\varphi(u) = au + b, \quad \psi(u) = cu + d \quad (ac \neq 0)$$

for all $u \in J$. Whence, with the notation $f := f_3$, $f \in \mathcal{CM}(I)$, and since $\varphi = f_1 \circ f_3^{-1} = f_1 \circ f^{-1}$, $\psi = f_2 \circ f_3^{-1} = f_2 \circ f^{-1}$, we have $f_1 = \varphi \circ f$, $f_2 = \psi \circ f$, where φ and ψ are given in (5.1). Thus, $M_i(x, y; \lambda) = A_f(x, y; \lambda)$ is valid for all $x, y \in I$ and $i = 1, 2, 3$.

References

- [1] **Borwein J.M. and Borwein P.B.**, *Pi and the AGM (A study in analytic number theory and computational complexity)*, John Wiley & Sons Inc., New York, 1987.
- [2] **Daróczy Z., Hajdu G. and Che Tat Ng**, An extension theorem for a Matkowski-Sutô problem, *Colloq. Math.*, (2003) (to appear)
- [3] **Daróczy Z. and Maksa Gy.**, On a problem of Matkowski, *Colloq. Math.*, **82** (1999), 117-123.
- [4] **Daróczy Z., Maksa Gy. and Páles Zs.**, Extension theorems for the Matkowski-Sutô problem, *Demonstratio Math.*, **33** (2000), 547-556.
- [5] **Daróczy Z. and Páles Zs.**, On means that are both quasi-arithmetic and conjugate arithmetic, *Acta Math. Hungar.*, **90** (2001), 271-282.
- [6] **Daróczy Z. and Páles Zs.**, Gauss-compositions of means and the solution of the Matkowski-Sutô problem, *Publ. Math. Debrecen*, **61** (2002), 157-218.
- [7] **Daróczy Z. and Páles Zs.**, The Matkowski-Sutô problem for weighted quasi-arithmetic means, *Acta Math. Hungar.*, (2003) (to appear)
- [8] **Hardy G.H., Littlewood J.E. and Pólya G.**, *Inequalities*, Cambridge University Press, Cambridge, 1934 (first edition), 1952 (second edition).
- [9] **Járai A.**, Baire property implies continuity for solutions of functional equations - even with few variables, *Acta Sci. Math. (Szeged)*, **66** (2000), 579-601.
- [10] **Laczkovich M.**, *Valós függvénytan (Real function theory)*, Eötvös Loránd Tudományegyetem Természettudományi Kar, Budapest, 1995.
- [11] **Matkowski J.**, Invariant and complementary quasi-arithmetic means, *Aequationes Math.*, **57** (1999), 87-107.
- [12] **Natanson I.P.**, *Theorie der Funktionen einer reellen Veränderlichen*, 5th ed., Akademie-Verlag, Berlin, 1981.
- [13] **Oxtoby J.C.**, *Measure and category. A survey of the analogies between topological and measure spaces*, Springer, New York, 1971. (Graduate Texts in Mathematics, Vol. 2.)
- [14] **Páles Zs.**, A regularity theorem for composite functional equations, *Acta Sci. Math. (Szeged)*, (2003) (to appear)
- [15] **Roberts A.W. and Varberg D.E.**, *Convex functions*, Academic Press, New York-London, 1973.
- [16] **Sutô O.**, Studies on some functional equations I., *Tôhoku Math. J.*, **6** (1914), 1-15.

- [17] **Sutô O.**, Studies on some functional equations II., *Tôhoku Math. J.*, **6** (1914), 82-101.

Z. Daróczy

Institute of Mathematics
University of Debrecen
H-4010 Debrecen, Pf. 12
Hungary
daroczy@math.klte.hu

Zs. Páles

Institute of Mathematics
University of Debrecen
H-4010 Debrecen, Pf. 12
Hungary
pales@math.klte.hu

