COMPUTATIONAL RESULTS AND QUERIES IN NUMBER THEORY

J.-M. De Koninck (Québec, Canada)

Dedicated to Professor Imre Kátai on his 65-th anniversary

1. Introduction

According to Moore's Law, the speed of modern computers doubles every 18 months. Hence it is no surprise that several old and new problems in number theory have gained special interest, mainly because one can study them using both a theoretical approach and a computational one. In this paper, we examine two such challenging problems.

Throughout this paper, we denote by **N** the set of positive integers and by **P** the set of prime numbers. Furthermore, given an integer $n \ge 2$, we shall write P(n) to denote its largest prime factor and $\omega(n)$ to denote the number of distinct prime factors of n.

2. Sum of the digits of prime numbers

Given a positive integer n, let s(n) stand for the sum of its (decimal) digits. Given a positive integer k which is not a multiple of 3, let $\rho(k)$ be the smallest prime number p such that s(p) = k. Below, is the table of values of $\rho(k)$ for $k = 2, 4, 5, 7, 8, \ldots, 83$.

Is it clear that $\rho(k)$ is a well defined function? Not at all! In fact, it is an open problem. Nevertheless, given any integer $k \ge 4$ which is not a multiple of 3, the set $A_k := \{p \in \mathbf{P} : s(p) = k\}$ is expected to be non empty and, in fact, is most likely infinite.

This research was supported in part by a grant from NSERC.

k	ho(k)		k	$\rho($	k)		k	$\rho($	k)
2	2		19		199		35	8	999
4	13		20		389		37	29	989
5	5		22	4	499		38	39	989
7	7		23		599		40	49	999
8	17		25	9	997		41	59	999
10	19		26	18	889		43	79	999
11	29		28	19	999		44	98	999
13	67		29	29	999		46	199	999
14	59		31	49	999		47	389	999
16	79		32	68	399		49	598	999
17	89		34	179	17989 50 5		599	999	
]				
	k	$\rho($	k)		k		$\rho(k)$		
	52	79	9999		68		59999999		
	53	98	9999		70		189997	999	
	55	299	8999		71		899999	999	
	56	299	9999		73		2899999999		
	58	499	9999		74		3899999	999	
	59	699	9899		76		6898999	999	
	61	898	9999		77		6998999	999	
	62	989	9999		79		7999999	999	
	64	1999	19999999		80		998999999		
	65	2999	29999999		82	2	2999899999		
	67	5989	9999		83	3	999998	999	

The size of A_2 is a different matter. Indeed, since one can easily show that any element of A_2 larger than 2 must be of the form $a_r := 10^{2r} + 1$ and since a_0 and a_1 are primes, and since, using a computer, one can establish that a_r is composite for $2 \le r \le 17$, it follows that any prime number $p \in A_2$, $p \ne 2$, 11, 101, must be larger than $2^{2^{18}}$. Therefore, it is possible that $\#A_2 = 3$.

On the other hand, one can easily show that $\#A_k \ge 1$ infinitely often. Indeed, assume the contrary, meaning that $\#A_k \ge 1$ only a finite number of times, and therefore assume that there exists a positive integer k_0 such that $\#A_k = 0$ for all $k \ge k_0$, thus implying that $s(p) < k_0$ for all primes p. Let $\ell = \lfloor k_0/9 \rfloor + 1$ and consider the sequence of integers

$$a_m := 10^{\ell} \cdot m + \underbrace{99\ldots9}_{\ell \text{ times}} \quad (m = 1, 2, \ldots).$$

It follows from Dirichlet's theorem that there exists a positive integer m_0 such that a_{m_0} is a prime number, in which case

$$s(a_{m_0}) = s(m_0) + 9\ell > s(m_0) + 9\frac{k_0}{9} \ge 1 + k_0 > k_0,$$

thus contradicting the fact that $s(p) < k_0$ for all primes p.

Finally, observe that $\rho(k)$ cannot be "too large" all the time; in fact, one can show that there exist infinitely many positive integers k such that $\rho(k) \leq 10^{2k}$. Indeed, assume that there are only finitely many positive integers k such that $\rho(k) \leq 10^{2k}$, in which case there exists an integer k_0 such that $\rho(k) > 10^{2k}$ for all $k \geq k_0$. This means in particular that, for all $k \geq k_0$, each prime p with s(p) = k is such that $p > 10^{2k}$. Therefore, in particular, if $p \leq 10^{2k_0}$ then $s(p) < k_0$. But this is impossible since the number of primes $p \leq 10^{2k_0}$ is much larger than $S(k_0)$, the number of primes p with no more than $2k_0$ digits and such that $s(p) < k_0$. Indeed, (1)

$$S(k_{0}) < \sum_{\substack{k=2\\k/3 \notin N}}^{k_{0}-1} \#\{p \in \mathbf{P} : s(p) = k \text{ with } p \text{ having no more than } 2k_{0} \text{ digits}\} < \\ < \sum_{k=2}^{k_{0}-1} \#\{n \in \mathbf{N} : s(n) = k \text{ with } n \text{ having no more than } 2k_{0} \text{ digits}\} < \\ < \sum_{k=2}^{k_{0}-1} p_{9}(k, 2k_{0}-1),$$

where $p_r(k, 2k_0 - 1)$ stands for the number of partitions of k into parts $\in [0, r]$, but with no more than $2k_0 - 1$ zeros amongst its parts. Since $p_9(k, 2k_0 - 1) < p_9(k) \cdot 2^{2k_0-1}$, where $p_r(k)$ is the number of partitions of k into parts $\in [1, r]$, then it follows from (1) that

(2)
$$S(k_0) < 4^{k_0} \sum_{k=2}^{k_0-1} p_9(k).$$

Using the general estimate

$$p_r(k) \sim \frac{k^{r-1}}{r!(r-1)!} \quad (k \to \infty)$$

(see Corollary 15.1 of Nathanson [4]), it follows from (2) that

(3)
$$S(k_0) \ll 4^{k_0} \sum_{k=2}^{k_0-1} \frac{k^8}{9!8!} \ll 4^{k_0} k_0^9,$$

a quantity which is clearly smaller that

$$\pi(10^{2k_0}) \approx \frac{10^{2k_0}}{2k_0 \log 10}$$

if k_0 is sufficiently large, thus proving our claim.

Here are some other open problems concerning $\rho(k)$:

1. It is easy to show that if $\rho(k)$ exists, then $\rho(k) \ge (a+1)10^b - 1$, where $b = [k/9] \ a = k - 9b$. In the above table, equality holds when k = 5, 7, 10, 11, 14, 16, 17, 19, 22, 23, 28, 29, 31, 35, 40, 41, 43, 46, 50, 52, 56, 58, 64, 65, 68, 71 and 79. Does it hold infinitely often? For that matter, is it true that

$$\lim_{k \to \infty} \frac{\rho(k)}{(a+1)10^b - 1} = 1?$$

2. Is it true that $\rho(k) \equiv 9 \pmod{10}$ for all k > 25? That $\rho(k) \equiv \equiv 99 \pmod{100}$ for all k > 38? That $\rho(k) \equiv 999 \pmod{1000}$ for all k > 59?

3. The sum of the prime factors of an integer

Let S(n) denote the sum of the prime factors of n taken with multiplicity. A number n is called a *Ruth-Aaron number* if S(n) = S(n+1). It is not known if there exist infinitely many such numbers, a problem which was first studied in 1974 by Nelson, Penney and Pomerance [5].

Similarly, let $\beta(n)$ denote the sum of the distinct prime factors of the integer $n \geq 2$. It is not known if the equation

(4)
$$\beta(n) = \beta(n+1)$$

has infinitely many solutions. Moreover, if one could show that there are infinitely many prime numbers p such that the three numbers

$$r = 6p - 1, \quad s = 10p - 1, \quad q = 15p - 4$$

are also prime numbers, then the corresponding number n = 4pq = rs - 1 would automatically be a solution of $\beta(n) = \beta(n+1)$ (= 16p-2), thus enabling one to construct an infinite family of solutions of (4). The existence of such an infinite sequence of 4-tuples of primes would follow from the *prime k-tuples conjecture* stated by L.E. Dickson [1] in 1904. This conjecture can be written as follows:

If a_1, a_2, \ldots, a_k and b_1, b_2, \ldots, b_k are integers such that $a_i > 0$ and $gcd(a_i, b_i) = 1$, and if for every prime number $p \leq k$, there exists a positive integer n such that none of the integers $a_i n + b_i$ is divisible by p, then there exists an infinite number of positive integers n such that each number $a_i n + b_i$ is prime.

The following table provides the values of B(x), the number of solutions $n \leq x$ of (4), for $x = 10^{j}$, j = 1, 2, ..., 9.

x	10^{1}	10^{2}	10^{3}	10^{4}	10^{5}	10^{6}	107	108	10^{9}
B(x)	2	5	10	20	41	140	495	1749	6651

Eventhough no interesting lower bound is known for B(x), upper bounds have been obtained. In 1978, Erdős and Pomerance [2] proved that, if C(x)stands for the number of Ruth-Aaron numbers $\leq x$, then

(5)
$$C(x) = O\left(\frac{x}{\log x}\right).$$

Very recently, Pomerance [6] improved this estimate by establishing that

(6)
$$C(x) = O\left(\frac{x(\log\log x)^4}{\log^2 x}\right).$$

Now, as Pomerance [6] observed, (5) and (6) also hold if C(x) is replaced by B(x).

Although very little is known about the solutions of (4), it is nevertheless interesting to consider an even more general problem, that is to find, given an arbitrary integer $k \ge 2$, the possible solutions of

(7)
$$\beta(n) = \beta(n+1) = \ldots = \beta(n+k-1).$$

In the particular case k = 3, the number n = 89460294 is a solution of (7) and a computer search reveals that there is no other solution $n < 10^{10}$. Nevertheless, given any integer $k \ge 2$, equations (7) may indeed have infinitely many solutions. In fact, Carl Pomerance (private communication) came up with a heuristic argument indicating that, given any arbitrary small $\varepsilon > 0$, the number $B_k(x)$ of solutions $n \leq x$ of (7) satisfies

$$B_k(x) \gg x^{1-\varepsilon}.$$

His argument goes as follows. Let $k \ge 2$ and choose an arbitrary small $\varepsilon > 0$. Let r be a fixed positive integer such that

$$\frac{k-1}{r} < \frac{\varepsilon}{2},$$

and consider the set $\{n \leq x : P(n) \leq n^{1/r}\}$, which is known to be of density $\rho(r)$, where ρ is the Dickman function (see for instance the book of Tenenbaum [7]). It is reasonable to assume that the set

$$D_k := \{n \le x : \max(P(n), P(n+1), \dots, P(n+k-1)) \le n^{1/r}\}$$

is also a set of positive density δ_k (with possibly $\delta_k \approx \rho(r)^k$). Since $\beta(n) \leq P(n)\omega(n) \leq P(n)\log n$ for each integer $n \geq 3$, then if $n \in D_k$, we have

$$\beta(n+i) \in [2, n^{1/r} \log(n+k-1)]$$

for i = 0, 1, ..., k - 1. It follows that the probability that (7) holds for a given integer $n \in [3, x]$ is equal to

$$\delta_k \left(n^{1/r} \log(n+k-1) \right)^{-(k-1)} = \delta_k \frac{1}{n^{(k-1)/r} \log^{(k-1)}(n+k-1)} > \delta_k \frac{1}{n^{\varepsilon}},$$

provided n is large enough so that $\log^{(k-1)}(n+k-1) < n^{\varepsilon/2}$. Therefore, if x is sufficiently large, we would have

$$B_k(x) > \delta_k \frac{x}{x^{\varepsilon}} \gg x^{1-\varepsilon}$$

as claimed.

Now consider the function $\beta^*(n) := \beta(n) - P(n)$. It is clearly much smaller than $\beta(n)$ since, denoting by $P_2(n)$ the second largest prime factor of n (or 0 if no such factor exists), we have

$$\beta^*(n) \le (\omega(n) - 1)P_2(n) \le (\log n)\sqrt{n}$$

Hence one might expect the corresponding equations

(8)
$$\beta^*(n) = \beta^*(n+1) = \ldots = \beta^*(n+k-1),$$

to have more solutions. In fact, we can show that (8) has infinitely many solutions n if one accepts the prime k-tuples conjecture.

Hence, we shall prove the following result.

Theorem 1. Let $k \geq 2$ be an integer. If one assumes that the prime k-tuples conjecture is true, then there exist infinitely many positive integers n such that

$$\beta^*(n-k+1) = \beta^*(n-k+2) = \ldots = \beta^*(n).$$

Proof. Let $k \ge 2$ and $A = \{p \in \mathbf{P} : p \le k\}$. Let E_1, E_2, \ldots, E_k be k sets of primes and R a positive integer satisfying the following conditions:

(a) $(E_i \setminus A) \cap (E_j \setminus A) = \emptyset$ if $i \neq j$;

(b) each prime $q \in A$ belongs to one and only one of the sets $E_i, E_{i+1}, \ldots, E_{i+q-1}$ for each integer $i \in [1, k-q+1]$;

(c) $\sum_{p \in E_1} p = \sum_{p \in E_2} p = \dots = \sum_{p \in E_k} p = R.$

For the moment, assume that such sets E_i 's with an appropriate integer R exist. Then let $Q_i = \prod_{p \in E_i} p$ for i = 1, 2, ..., k and consider the system of congruences

(9)
$$\begin{cases} n-k+1 \equiv 0 \pmod{Q_1}, \\ n-k+2 \equiv 0 \pmod{Q_2}, \\ \vdots \\ n \equiv 0 \pmod{Q_k}. \end{cases}$$

Using the Chinese Remainder Theorem, we are guaranteed the existence of a solution $n_0 < Q := Q_1 Q_2 \dots Q_k$ of (9), each other solution n of (9) being of the form

$$n = n_0 + xQ$$
, with $x = 0, 1, 2, 3, \dots$

Hence, if we set

$$g_i := \gcd\left(\frac{n_0 + k - i}{Q_i}, \frac{Q}{Q_i}\right) \quad (1 \le i \le k),$$

then, for each integer $i \in [1, k]$,

$$n - k + i = n_0 - k + i + xQ = Q_i g_i \left(\frac{n_0 - k + i}{Q_i g_i} + x\frac{Q}{Q_i g_i}\right) = Q_i g_i \cdot q_i(x),$$

where each polynomial $q_i(x)$, $1 \leq i \leq k$, is defined implicitly. If we can find a value of x, say r, for which the corresponding numbers $q_1(r), q_2(r), \ldots, q_k(r)$ are all prime numbers, with

(10)
$$q_i(r) > P(Q_i g_i) \quad (1 \le i \le k),$$

then our theorem will be proved, since in this case

$$\beta^*(n-k+i) = \beta^*(Q_i g_i q_i(r)) = \beta(Q_i g_i) = \beta(Q_i) = R \quad (1 \le i \le k).$$

Now

$$\operatorname{pgcd}\left(\frac{n_0 - k + i}{Q_i g_i}, \frac{Q}{Q_i g_i}\right) = 1 \quad (1 \le i \le k)$$

and, for each integer $i \in [1, k]$, no prime $p \in A$ divides each value of $q_i(x)$. Hence we may conclude, assuming the *prime k-tuples conjecture*, that there exists one positive integer r (as a matter of fact, infinitely many) such that the corresponding numbers $q_1(r), q_2(r), \ldots, q_k(r)$ are all primes, and moreover that they satisfy (10).

Hence, in order to complete the proof of Theorem 1, it remains to show that one can always find k sets of primes E_1, \ldots, E_k and in integer R satisfying conditions (a), (b) and (c).

First, observe that if we denote by $s_3(n)$ the number of representations of n as a sum of three primes, then using the known estimate

$$\sum_{q_1+q_2+q_3=n} (\log q_1)(\log q_2)(\log q_3) \sim \frac{n^2}{2} G(n) \quad (n \to \infty),$$

where

$$G(n) = \prod_{p|n} \left(1 - \frac{1}{(p-1)^2} \right) \prod_{p \notin n} \left(1 + \frac{1}{(p-1)^2} \right)$$

(see for instance Friedlander and Goldston [3]), one can easily establish that

$$s_3(n) \gg \frac{n^2}{\log^3 n} \quad (n \to \infty).$$

It follows from this estimate that, provided an odd integer n is large enough, the number of representations of n as a sum of three odd primes can be as large as one desires. Using this observation, we now show how one can contruct sets of primes E_1, E_2, \ldots, E_k and a corresponding odd integer R satisfying conditions (a), (b) and (c).

Let R be a large odd integer; how large it needs to be will become clear at the end of this argument. Let $F_i = \{r_i, s_i, t_i\}, i = 1, 2, ..., k$, be k sets each containing exactly three distinct primes whose sum is equal to R, and such that $F_i \cap F_j = \emptyset$ if $i \neq j$. Let $R_1, R_2, R_3, ...$ be odd integers defined implicity by the following relations:

$$\begin{array}{c} 2+R_{1}=R,\\ r_{1}+s_{1}+t_{1}=R,\\ 2+3+5+R_{2}=R,\\ r_{2}+s_{2}+t_{2}=R,\\ 2+R_{1}=R,\\ 3+7+R_{3}=R,\\ 2+R_{1}=R,\\ 5+11+R_{4}=R,\\ 2+3+13+R_{5}=R,\\ r_{3}+s_{3}+t_{3}=R, \end{array}$$

and so on, where:

- on each *i*-th row with *i* odd, we place the prime 2;
- on each *i*-th row with $i \ge 3$:
 - if there exists a prime p which has already appeared on the (i p)-th row,
 - * we place it on this *i*-th row;
 - * we also place the smallest prime which has not yet appeared on the previous lines, with the additional constraint that we place the two smallest if the left side of the equation is even;
 - if no such prime p exists, we place the next set F_j of primes;
- we end the process when the k-th line has been constructed.

We then define the sets E_i 's as follows (where the primes q_i 's are primes whose existence is guaranteed by the fact that each odd large integer can be written as the sum of three primes):

$$E_1 = \{2, q_1, q_2, q_3\}, \text{ where } q_1 + q_2 + q_3 = R_1,$$

 $E_2 = \{r_1, s_1, t_1\},$

$$\begin{split} E_3 &= \{2, 3, 5, q_4, q_5, q_6\}, & \text{where } q_4 + q_5 + q_6 = R_2, \\ E_4 &= \{r_2, s_2, t_2\}, \\ E_5 &= \{2, q_7, q_8, q_9\}, & \text{where } q_7 + q_8 + q_9 = R_3, \\ E_6 &= \{3, 7, q_{10}, q_{11}, q_{12}\}, & \text{where } q_{10} + q_{11} + q_{12} = R_4, \\ E_7 &= \{2, q_{13}, q_{14}, q_{15}\}, & \text{where } q_{13} + q_{14} + q_{15} = R_5, \\ E_8 &= \{5, 11, q_{16}, q_{17}, q_{18}\}, & \text{where } q_{16} + q_{17} + q_{18} = R_6, \\ E_9 &= \{2, 3, 13, q_{19}, q_{20}, q_{21}\}, & \text{where } q_{19} + q_{20} + q_{21} = R_7, \\ E_{10} &= \{r_3, s_3, t_3\}, \end{split}$$

and so on, up to E_k .

Since R can be taken arbitrarily large, we may assume that each of the primes in the F_i 's are larger than the q_i 's, which in turn may be assumed to be larger than the p's which we have been adding on some lines. By doing so, we ensure that conditions (a), (b) and (c) are satisfied, thus completing the proof of Theorem 1.

For each positive real number $\varepsilon < 1$, let

$$\beta_{\varepsilon}(n) := \sum_{\substack{p \mid n \\ p < n^{1-\varepsilon}}} p$$

Since, in the above proof, the prime k-tuples conjecture guarantees the existence of infinitely many integers x such that $q_i(x)$ is prime for $1 \le i \le k$, and since for each $i \in [1, k]$, the sequence of primes $q_i(x)$ tends to $+\infty$, it is clear that the following result also holds.

Theorem 2. Given an integer $k \ge 2$ and a positive real number $\varepsilon < < 1$. Then, assuming the prime k-tuples conjecture, there exist infinitely many positive integers n such that

$$\beta_{\varepsilon}(n-k+1) = \beta_{\varepsilon}(n-k+2) = \ldots = \beta_{\varepsilon}(n).$$

Using the idea of the proof of Theorem 1, we can also establish that, assuming the *prime k-tuples conjecture*, the quotients $\frac{\beta(n-k+j)}{\beta(n-k+i)}$, with $i, j \in [1, k]$ can be arbitrarily close to 1, infinitely many often. More precisely, we can prove the following result.

Theorem 3. Given an integer $k \geq 2$ and assuming the prime k-tuples conjecture,

(11)
$$\liminf_{n \to \infty} \frac{\max_{1 \le i \le k} \beta(n-k+i)}{\min_{1 \le i \le k} \beta(n-k+i)} = 1.$$

Proof. We only need to show that, given any real number $\delta > 0$, there exist infinitely many integers n such that

(12)
$$\frac{\max_{1 \le i \le k} \beta(n-k+i)}{\min_{1 \le i \le k} \beta(n-k+i)} \le 1+\delta.$$

Let N be a large positive integer and consider the system of congruences

$$\begin{cases} n-k+1 \equiv 0 \pmod{N-k+1}, \\ n-k+2 \equiv 0 \pmod{N-k+2}, \\ \vdots \\ n \equiv 0 \pmod{N}, \end{cases}$$

of which $n_0 = N$ is clearly the smallest positive solution, in which case all other solutions n are given by $n = n_0 + xQ$, x = 0, 1, 2, ..., where

$$Q := (N - k + 1)(N - k + 2) \dots N$$

We then have, for each integer $i \in [1, k]$,

$$n - k + i = n_0 - k + i + xQ = N - k + i + xQ =$$

= $(N - k + i)\left(1 + x\frac{Q}{N - k + i}\right) = (N - k + i) \cdot q_i(x),$

say. Assuming the prime k-tuples conjecture, we may conclude that there exists a positive integer r such that $q_1(r), \ldots, q_k(r)$ are all prime numbers. For the corresponding integer n, with $\beta(n-k+j) \geq \beta(n-k+i)$, we have

$$\begin{split} 1 &\leq \frac{\beta(n-k+j)}{\beta(n-k+i)} = \frac{\beta((N-k+j)q_{j}(r))}{\beta((N-k+i)q_{i}(r))} = \frac{\beta(N-k+j)+q_{j}(r)}{\beta(N-k+i)+q_{i}(r)} = \\ &= \frac{\beta(N-k+j)+(n-k+j)/(N-k+j)}{\beta(N-k+i)+(n-k+j)/(N-k+i)} \leq \\ &\leq \frac{\beta(N-k+j)+(n-k+j)/(N-k+j)}{(n-k+i)/(N-k+i)} = \\ &= \frac{\beta(N-k+j)\frac{N-k+j}{n-k+j}+1}{\frac{n-k+i}{n-k+j}\cdot\frac{N-k+j}{N-k+i}} \leq \\ &\leq 1+\delta, \end{split}$$

provided n is large enough, thus proving Theorem 3.

Computational data. The algorithm mentioned in the proof of Theorem 1 is not optimal; it only guarantees the existence of a set of primes E_i 's with an appropriate integer R. Much simpler configurations can be obtained by inspection, say for example for $1 \le k \le 5$. For k = 2, one can take R = 5 with $E_1 = \{5\}$ and $E_2 = \{2,3\}$ which yields as part of the solutions of (8) the numbers n = 65, 185, 365, 785 and 905. For k = 3, take R = 19 with $E_1 = \{19\}, E_2 = \{2,17\}$ and $E_3 = = \{3, 5, 11\}$ providing the solutions

n = 8161393, 18607213, 26068513, 64014553 and 67212253.

For k = 4, one can take R = 36 with $E_1 = \{2, 5, 29\}$, $E_2 = \{13, 23\}$, $E_3 = \{2, 3, 31\}$ and $E_4 = \{17, 19\}$, with solutions

73494447431170, 111699918460090, 112793884769890,

127098796611370 and 141955900971130.

For k = 5, one can take R = 50, $E_1 = \{13, 37\}$, $E_2 = \{2, 5, 43\}$, $E_3 = \{3, 47\}$, $E_4 = \{2, 7, 41\}$ and $E_5 = \{19, 31\}$, providing the solutions

981705863038517929, 1036978848810729409, 1052399282229876529,

1595959700642651929 and 2050448382796747609.

Acknowledgement. The author would like to thank Carl Pomerance for his helpful comments which helped to improve the quality of this paper.

References

- Dickson L.E., A new extension of Dirichlet's theorem on prime numbers, Messenger of Math., 33 (1904), 155-161.
- [2] Erdős P. and Pomerance C., On the largest prime factors of n and n + 1, Aequationes Mathematicae, 17 (1978), 311-321.
- [3] Friedlander J.B. and Goldston D.A., Sums of three or more primes, Trans. Amer. Math. Soc., 349 (1997), 287-310.
- [4] Nathanson M.B., Elementary methods in number theory, Springer, 2000.
- [5] Nelson C., Penney D.E. and Pomerance C., 714 and 715, J. Recreational Math., 7 (1974), 87-89.

- [6] Pomerance C., Ruth-Aaron numbers revisited, Bolyai Soc. Math. Stud. 11, Paul Erdős and his Mathematics I., Budapest, 2002, 567-579.
- [7] **Tenenbaum G.**, Introduction to analytic and probabilistic number theory, Cambridge University Press, 1995.

(Received June 4, 2003)

J.-M. De Koninck Département de mathématiques et de statistique Université Laval Québec G1K 7P4 Canada jmdk@mat.ulaval.ca