

## MOMENT FUNCTIONS ON STURM-LIOUVILLE HYPERGROUPS

Á. Orosz and L. Székelyhidi

(Debrecen, Hungary)

*Dedicated to Professor Imre Kátaí on the occasion of his 70th birthday*

**Abstract.** This paper presents some recent results concerning functional equations on Sturm–Liouville hypergroups. The general form of additive functions, exponentials and moment functions on these types of hypergroups is given.

### 1. Introduction

The concept of DJS-hypergroup (according to the initials of C.F. Dunkl, R.I. Jewett and R. Spector) can be introduced using different axiom systems. The way of introducing the concept here is due to R. Lasser (see e.g. [2], [6]). One begins with a locally compact Hausdorff space  $K$ , the space  $\mathcal{M}(K)$  of all finite complex regular measures on  $K$ , the space  $\mathcal{M}_c(K)$  of all finitely supported measures in  $\mathcal{M}(K)$ , the space  $\mathcal{M}^1(K)$  of all probability measures in  $\mathcal{M}(K)$ , and the space  $\mathcal{M}_c^1(K)$  of all compactly supported probability measures in  $\mathcal{M}(K)$ . The point mass concentrated at  $x$  is denoted by  $\delta_x$ . Suppose that we have the following:

- ( $H^*$ ) There is a continuous mapping  $(x, y) \mapsto \delta_x * \delta_y$  from  $K \times K$  into  $\mathcal{M}_c^1(K)$ , the latter being endowed with the weak\*-topology with respect to the space of compactly supported complex valued continuous functions on  $K$ . This mapping is called *convolution*.

- ( $H^\vee$ ) There is an involutive homeomorphism  $x \mapsto x^\vee$  from  $K$  to  $K$ . This mapping is called *involution*.
- ( $He$ ) There is a fixed element  $e$  in  $K$ . This element is called *identity*.

Identifying  $x$  by  $\delta_x$  the mapping in  $(H^*)$  has a unique extension to a continuous bilinear mapping from  $\mathcal{M}(K) \times \mathcal{M}(K)$  to  $\mathcal{M}(K)$ . The involution on  $K$  extends to an involution on  $\mathcal{M}(K)$ . Then a *DJS-hypergroup*, or simply *hypergroup*, is a quadruple  $(K, *, \vee, e)$  satisfying the following axioms: for any  $x, y, z$  in  $K$  we have

- (H1)  $\delta_x * (\delta_y * \delta_z) = (\delta_x * \delta_y) * \delta_z$ ;  
 (H2)  $(\delta_x * \delta_y)^\vee = \delta_{y^\vee} * \delta_{x^\vee}$ ;  
 (H3)  $\delta_x * \delta_e = \delta_e * \delta_x = \delta_x$ ;  
 (H4)  $e$  is in the support of  $\delta_x * \delta_{y^\vee}$  if and only if  $x = y$ ;  
 (H5) the mapping  $(x, y) \mapsto \text{supp}(\delta_x * \delta_y)$  from  $K \times K$  into the space of nonvoid compact subsets of  $K$  is continuous, the latter being endowed with the Michael-topology (see [2]).

If  $\delta_x * \delta_y = \delta_y * \delta_x$  holds for all  $x, y$  in  $K$ , then we call the hypergroup *commutative*. If  $x^\vee = x$  holds for all  $x$  in  $K$  then we call the hypergroup *Hermitian*. By (H2) any Hermitian hypergroup is commutative. For instance, if  $K = G$  is a locally compact Hausdorff-group,  $\delta_x * \delta_y = \delta_{xy}$  for all  $x, y$  in  $K$ ,  $x^\vee$  is the inverse of  $x$ , and  $e$  is the identity of  $G$ , then we obviously have a hypergroup  $(K, *, \vee, e)$ , which is commutative if and only if the group  $G$  is commutative. However, not every hypergroup originates in this way.

In any hypergroup  $K$  we identify  $x$  by  $\delta_x$  and we define the *right translation operator*  $T_y$  by the element  $y$  in  $K$  according to the formula

$$T_y f(x) = \int_K f d(\delta_x * \delta_y),$$

for any  $f$  integrable with respect to  $\delta_x * \delta_y$ . In particular,  $T_y$  is defined for any continuous complex valued function on  $K$ . Similarly, we can define *left translation operators* but at this moment we do not need any extra notation for them.

Sometimes one uses the suggestive notation

$$f(x * y) = \int_K f d(\delta_x * \delta_y),$$

for any  $x, y$  in  $K$ . However, we call the attention to the fact that actually  $f(x * y)$  has no meaning in itself, because  $x * y$  is in general not an element of  $K$ , hence  $f$  is not defined at  $x * y$ .

If  $K$  is a discrete topological space, then we call the hypergroup a *discrete hypergroup*. An important special class of discrete hypergroups are the *polynomial hypergroups* which are closely related to orthogonal polynomials. For the definition and a detailed study of polynomial hypergroups the reader should refer to [2].

Another important class of hypergroups is the class of *Sturm-Liouville hypergroups*. The definition and some basic properties of Sturm-Liouville hypergroups will be given in the following section.

In our former paper [7] we presented some recent results concerning functional equations on hypergroups. The aim was to give some idea for the treatment of classical functional equation problems in the hypergroup setting. We described the general form of additive functions, exponentials and moment functions of second order on polynomial hypergroups. Here we consider similar problems concerning classical functional equations on Sturm-Liouville hypergroups.

## 2. Sturm-Liouville hypergroups

Sturm-Liouville hypergroups represent another important class of hypergroups, which arise from Sturm-Liouville boundary value problems on the nonnegative reals. In order to build up the Sturm-Liouville operator basic to the construction of hypergroups one introduces the Sturm-Liouville functions. For further details see [2]. In what follows  $\mathbb{R}_0$  denotes the set of nonnegative real numbers.

The continuous function  $A : \mathbb{R}_0 \rightarrow \mathbb{R}$  is called a *Sturm-Liouville function*, if it is positive and continuously differentiable on the positive reals. Different assumptions on  $A$  can be found in [2] which lead to the desired Sturm-Liouville problem. For a given Sturm-Liouville function  $A$  one defines the *Sturm-Liouville operator*  $L_A$  by

$$L_A f = -f'' - \frac{A'}{A} f',$$

where  $f$  is a twice continuously differentiable real function on the positive reals. Using  $L_A$  one introduces the differential operator  $l$  by

$$l[u](x, y) = (L_A)_x u(x, y) - (L_A)_y u(x, y) =$$

$$= -\partial_1^2 u(x, y) - \frac{A'(x)}{A(x)} \partial_1 u(x, y) + \partial_2^2 u(x, y) + \frac{A'(y)}{A(y)} \partial_2 u(x, y),$$

where  $u$  is twice continuously differentiable for all positive reals  $x, y$ . Here  $(L_A)_x$  and  $(L_A)_y$  indicates that  $L_A$  operates on functions depending on  $x$  or  $y$ , respectively.

A hypergroup on  $\mathbb{R}_0$  is called a *Sturm-Liouville hypergroup* if there exists a Sturm-Liouville function  $A$  such that given any real-valued  $C^\infty$ -function  $f$  on  $\mathbb{R}_0$  the function  $u_f$  defined by

$$u_f(x, y) = \int_{\mathbb{R}_0} f d(\delta_x * \delta_y)$$

for all positive  $x, y$  is twice continuously differentiable and satisfies the partial differential equation

$$l[u_f] = 0$$

with  $\partial_2 u_f(x, 0) = 0$  for all positive  $x$ . Hence  $u_f$  is a solution of the Cauchy-problem

$$\partial_1^2 u(x, y) + \frac{A'(x)}{A(x)} \partial_1 u(x, y) = \partial_2^2 u(x, y) + \frac{A'(y)}{A(y)} \partial_2 u(x, y),$$

$$\partial_2 u(x, 0) = 0$$

for all positive  $x, y$ . From general properties of one-dimensional hypergroups given in [2] it follows that  $u_f(y, 0) = u_f(0, y) = f(y)$  and  $\partial_1 u_f(0, y) = 0$  hold, whenever  $y$  is a positive real number. In other words,  $u_f$  is the unique solution of the boundary value problem

$$(1) \quad \begin{aligned} \partial_1^2 u(x, y) + \frac{A'(x)}{A(x)} \partial_1 u(x, y) &= \partial_2^2 u(x, y) + \frac{A'(y)}{A(y)} \partial_2 u(x, y), \\ \partial_1 u(0, y) &= 0, \quad \partial_2 u(x, 0) = 0, \\ u(x, 0) &= f(x), \quad u(0, y) = f(y) \end{aligned}$$

for all positive  $x, y$ . As this boundary value problem uniquely defines  $u_f$  for any  $f$ , we may consider it the *boundary value problem defining the Sturm-Liouville hypergroup*.

If a Sturm-Liouville hypergroup structure is given on  $\mathbb{R}_0$  by the Sturm-Liouville function  $A$ , then we denote it by  $(\mathbb{R}_0, A)$ . If the Sturm-Liouville function  $A$  satisfies

$$(2) \quad \frac{A'(x)}{A(x)} = \frac{\alpha_0}{x} + \alpha_1(x)$$

for all  $x \neq 0$  in a neighborhood of 0 with  $\alpha_0 > 0$  such that  $\alpha_1$  is an odd  $C^\infty$ -function on  $\mathbb{R}$  and the function  $\frac{A'}{A}$  is nonnegative and decreasing, further  $A$  is increasing with  $\lim_{x \rightarrow +\infty} A(x) = +\infty$ , then  $A$  is called a *Chébli-Trimèche function* and the corresponding Sturm-Liouville hypergroup is called a *Chébli-Trimèche hypergroup*. Special cases are represented by the *Bessel-Kingman hypergroups* with  $A(0) = 0$  and

$$A(x) = x^\alpha$$

for all positive  $x$  and some  $\alpha > 0$ , and the *hyperbolic hypergroups*, where  $A(0) = 0$  and

$$A(x) = \sinh^a x$$

for all positive  $x$  and some  $a > 0$ .

If the Sturm-Liouville function  $A$  is twice continuously differentiable on the positive reals and satisfies (2), where  $\alpha_0 = 0$  and  $\alpha_1$  is continuously differentiable on the positive reals, then  $A$  is called a *Levitan function* and the corresponding Sturm-Liouville hypergroup is called a *Levitan hypergroup*. Special cases are represented by the *cosh hypergroup*, where

$$A(x) = \cosh^2 x$$

for all nonnegative  $x$ , and the *square hypergroup*, where

$$A(x) = (1 + x)^2$$

for all nonnegative  $x$  (see [8]). For more about these hypergroups and their applications see [2].

### 3. Exponentials, additive functions and moment functions on hypergroups

Let  $K$  be a hypergroup with convolution  $*$ , involution  $^\vee$ , and identity  $e$ . For any  $y$  in  $K$  let  $T_y$  denote the right translation operator on the space of all complex valued functions on  $K$  which are integrable with respect to  $\delta_x * \delta_y$  for any  $x, y$  in  $K$ . In particular, any continuous complex valued function belongs to this class.

The continuous complex valued function  $m$  on  $K$  is called an *exponential*, if it is not identically zero, and

$$T_y m(x) = m(x)m(y)$$

holds for all  $x, y$  in  $K$ . In other words  $m$  satisfies the functional equation

$$m(x * y) = m(x)m(y).$$

The continuous complex valued function  $a$  on  $K$  is called *additive*, if it satisfies

$$T_y a(x) = a(x) + a(y)$$

for all  $x, y$  in  $K$ . In other words this means that

$$a(x * y) = a(x) + a(y)$$

holds for any  $x, y$  in  $K$ . It is obvious that any linear combination of additive functions is additive again. However, in contrast to the case of groups, the product of exponentials is not necessarily an exponential.

The third important class of functions we want to study in this work is the class of moment functions. Moments of probability measures on a hypergroup can be introduced in terms of moment functions. The notion of moment functions has been formalized in [8] (see also [2]). For any nonnegative integer  $N$  the complex valued function  $f$  on  $K$  is called a *moment function of order  $N$* , if there are complex valued continuous functions  $f_k$  on  $K$  for  $k = 0, 1, \dots, N$  such that  $f_0 = 1$ ,  $f_N = f$ , and

$$(3) \quad f_k(x * y) = \sum_{j=0}^k \binom{k}{j} f_j(x) f_{k-j}(y)$$

holds for  $k = 0, 1, \dots, N$  and for all  $x, y$  in  $K$ . In this case we say that the functions  $f_k$  ( $k = 0, 1, \dots, N$ ) form a *moment sequence of order  $N$* . Hence moment functions of order 1 are exactly the additive functions. In [3] the general form of moment functions of order  $N = 1$  and  $N = 2$  have been determined in the case of polynomial hypergroups. We can generalize this concept by omitting the hypothesis  $f_0 = 1$  but still  $f_0$  is nonidentically zero. In this case  $f_0$  is an exponential function and we say that  $f_0$  *generates the generalized moment sequence of order  $N$*  and  $f_k$  is a *generalized moment function of order  $k$  with respect to  $f_0$*  ( $k = 0, 1, \dots, N$ ). For instance, generalized moment functions of order 1 with respect to the exponential  $f_0$  are solutions of the *sine functional equation*

$$f_1(x * y) = f_0(x)f_1(y) + f_0(y)f_1(x)$$

for any  $x, y$  in  $K$ .

The study of moment functions and moment sequences on hypergroups leads to the study of the above system of functional equations. We remark that

a similar system of functional equations on groupoids has been investigated and solved in [1].

#### 4. Exponentials and additive functions on Sturm-Liouville hypergroups

Let  $K = (\mathbb{R}_0, A)$  be a Sturm-Liouville hypergroup. Now we describe all exponentials defined on  $K$  (see also [2]).

**Theorem 4.1.** *Let  $K = (\mathbb{R}_0, A)$  be the Sturm-Liouville hypergroup corresponding to the Sturm-Liouville function  $A$ . Then the continuous function  $m : \mathbb{R}_0 \rightarrow \mathbb{C}$  is an exponential on  $K$  if and only if it is  $C^\infty$  and there exists a complex number  $\lambda$  such that*

$$(4) \quad m''(x) + \frac{A'(x)}{A(x)}m'(x) = \lambda m(x), \quad m(0) = 1, \quad m'(0) = 0$$

holds for any positive  $x$ .

**Proof.** First suppose that the function  $m : \mathbb{R}_0 \rightarrow \mathbb{C}$  is  $C^\infty$  on  $\mathbb{R}_0$  and it satisfies the given boundary value problem. Then the function

$$m(x * y) = \int_0^\infty m(t) d(\delta_x * \delta_y)(t)$$

and also the function  $(x, y) \rightarrow m(x)m(y)$  is a solution of the boundary value problem defining the hypergroup, hence they are equal and  $m$  is an exponential.

Conversely, suppose that  $m : \mathbb{R}_0 \rightarrow \mathbb{C}$  is an exponential on the hypergroup  $K$ . Then the function  $u_m(x, y) = m(x)m(y)$  is a solution of the boundary value problem defining the hypergroup, hence we obtain

$$\left( m''(x) + \frac{A'(x)}{A(x)}m'(x) \right) m(y) = \left( m''(y) + \frac{A'(y)}{A(y)}m'(y) \right) m(x)$$

holds for each positive  $x, y$ , and there exists a complex  $\lambda$  with

$$m''(x) + \frac{A'(x)}{A(x)}m'(x) = \lambda m(x)$$

for all positive  $x$ , consequently  $m$  is  $C^\infty$  on  $\mathbb{R}_0$ . The relations  $m(0) = 1$  and  $m'(0) = 0$  are immediate consequences of the fact that  $m$  is an exponential and the neutral element of the hypergroup is zero.

Hence any exponential function on a Sturm-Liouville hypergroup is an eigenfunction of the Sturm-Liouville operator corresponding to the given hypergroup. Each complex number is an eigenvalue and there is a one-to-one correspondence between complex numbers and exponentials. For any fixed complex  $\lambda$  we shall denote by  $x \mapsto \varphi(x, \lambda)$  the unique solution of the boundary value problem (4). Then the function  $\varphi : \mathbb{R}_0 \times \mathbb{C} \rightarrow \mathbb{C}$  represents a one-parameter family of exponentials of the Sturm-Liouville hypergroup  $K$ , which is called the *exponential family* of  $K$ . For instance, the complex number  $\lambda = 0$  corresponds to the eigenvalue problem

$$m''(x) + \frac{A'(x)}{A(x)}m'(x) = 0, \quad m(0) = 1, \quad m'(0) = 0,$$

which obviously has the unique solution  $m \equiv 1$ , hence  $\varphi(x, 0) = 1$  for each  $x$  in  $\mathbb{R}_0$ .

## 5. Additive functions on Sturm-Liouville hypergroups

Again let  $K = (\mathbb{R}_0, A)$  be a Sturm-Liouville hypergroup. Now we describe all additive functions defined on  $K$  (see also [2]).

**Theorem 5.1.** *Let  $K = (\mathbb{R}_0, A)$  be the Sturm-Liouville hypergroup corresponding to the Sturm-Liouville function  $A$ . Then the continuous function  $a : \mathbb{R}_0 \rightarrow \mathbb{C}$  is an additive function on  $K$  if and only if it is  $C^\infty$  and there exists a complex number  $\lambda$  such that*

$$(5) \quad a''(x) + \frac{A'(x)}{A(x)}a'(x) = \lambda, \quad a(0) = 0, \quad a'(0) = 0$$

*holds for any positive  $x$ .*

**Proof.** The proof is very similar to that of the previous theorem. First suppose that the function  $a : \mathbb{R}_0 \rightarrow \mathbb{C}$  is  $C^\infty$  and it satisfies the given boundary value problem. Then the function

$$a(x * y) = \int_0^\infty a(t) d(\delta_x * \delta_y)(t)$$



and also the function  $(x, y) \rightarrow a(x) + a(y)$  is a solution of the boundary value problem defining the hypergroup, hence they are equal and  $a$  is an additive function.

Conversely, suppose that  $a : \mathbb{R}_0 \rightarrow \mathbb{C}$  is an additive function on the given hypergroup  $K$ . Then the function  $u_a(x, y) = a(x) + a(y)$  is a solution of the boundary value problem defining the hypergroup, hence we obtain

$$a''(x) + \frac{A'(x)}{A(x)} a'(x) = a''(y) + \frac{A'(y)}{A(y)} a'(y)$$

holds for each positive  $x, y$ , and there exists a complex  $\lambda$  with

$$a''(x) + \frac{A'(x)}{A(x)} a'(x) = \lambda$$

for all positive  $x$ , consequently  $a$  is  $C^\infty$  on  $\mathbb{R}_0$ . The relations  $a(0) = 0$  and  $a'(0) = 0$  are immediate consequences of the fact that  $a$  is additive and the neutral element of the hypergroup is zero.

It is obvious that the unique solution  $a_\lambda$  of the boundary value problem (5) is  $\lambda a_1$ , where  $a_1$  is the unique solution of (5) with  $\lambda = 1$ . This means that all additive functions of a Sturm-Liouville hypergroup are constant multiples of a fixed nonzero additive function. We call  $a_1$  the *generating additive function* of the Sturm-Liouville hypergroup  $(\mathbb{R}_0, A)$ .

It turns out that the boundary value problem (5) can be solved explicitly. Namely, we have the following theorem (see [8]).

**Theorem 5.2.** *Let  $K = (\mathbb{R}_0, A)$  be the Sturm-Liouville hypergroup corresponding to the Sturm-Liouville function  $A$ . Then the generating additive function of the hypergroup  $K$  is given by*

$$(6) \quad a_1(x) = \int_0^x \int_0^y \frac{A(t)}{A(y)} dt dy$$

*for each nonnegative  $x$ . Hence any additive function of the hypergroup  $K$  is given by*

$$(7) \quad a_\lambda(x) = \lambda \int_0^x \int_0^y \frac{A(t)}{A(y)} dt dy$$

*for each nonnegative  $x$ .*

**Proof.** The proof is obvious using standard methods from the theory of linear differential equations. Another way of proving the statement is direct verification and using the uniqueness theorem.

As an illustration we compute the additive functions on the Bessel-Kingman hypergroup, which is a special Chébli-Trimèche hypergroup. Here  $A(x) = x^\alpha$  for all nonnegative  $x$  with some positive number  $\alpha$ . In this case we have

$$a_1(x) = \int_0^x \int_0^y \frac{t^\alpha}{y^\alpha} dt dy = \frac{x^2}{2(\alpha + 1)}$$

and

$$a_\lambda(x) = \lambda \int_0^x \int_0^y \frac{t^\alpha}{y^\alpha} dt dy = \frac{\lambda x^2}{2(\alpha + 1)}$$

for each nonnegative  $x$  and complex number  $\lambda$ .

Another example is given here for a special Levitan hypergroup, the square hypergroup, where  $A(x) = (1 + x)^2$  for all nonnegative  $x$ . From the above formulas we have

$$a_1(x) = \int_0^x \int_0^y \frac{(1+t)^2}{(1+y)^2} dt dy = \frac{x^3 + 3x^2}{6(x+1)}$$

and

$$a_\lambda(x) = \lambda \int_0^x \int_0^y \frac{(1+t)^2}{(1+y)^2} dt dy = \frac{\lambda(x^3 + 3x^2)}{6(x+1)}$$

for each nonnegative  $x$  and complex number  $\lambda$ .

## 6. Moment functions on Sturm-Liouville hypergroups

Let  $K = (\mathbb{R}_0, A)$  be a Sturm-Liouville hypergroup. In this section we describe all generalized moment functions defined on  $K$ . We remark that in [4] and [5] the general form of generalized moment functions on polynomial hypergroups is given.

**Theorem 6.1.** *Let  $K = (\mathbb{R}_0, A)$  be the Sturm-Liouville hypergroup corresponding to the Sturm-Liouville function  $A$  and let  $N$  be a positive integer.*

The continuous functions  $f_k : \mathbb{R}_0 \rightarrow \mathbb{C}$  ( $k = 0, 1, \dots, N$ ) form a sequence of generalized moment functions on the hypergroup  $K$  if and only if they are  $C^\infty$  and there are complex numbers  $c_k$  for  $k = 0, 1, \dots, N$  such that

$$(8) \quad f_0''(x) + \frac{A'(x)}{A(x)} f_0'(x) = c_0 f_0(x), \quad f_0(0) = 1, \quad f_0'(0) = 0$$

and

$$(9) \quad f_k''(x) + \frac{A'(x)}{A(x)} f_k'(x) = \sum_{j=0}^k \binom{k}{j} c_j f_{k-j}(x), \quad f_k(0) = 0, \quad f_k'(0) = 0$$

holds for each positive  $x$  and for  $k = 1, 2, \dots, N$ .

**Proof.** First we proof the sufficiency. If the functions  $f_k : \mathbb{R}_0 \rightarrow \mathbb{C}$  ( $k = 0, 1, \dots, N$ ) satisfy the conditions (8) and (9), then  $f_0$  is an exponential function and hence  $f_0(x * y) = f_0(x)f_0(y)$  holds for all nonnegative numbers  $x$  and  $y$ . We show that equation (3) holds for all  $k = 1, \dots, N$ , namely, the function

$$h(x, y) = \sum_{j=0}^k \binom{k}{j} f_j(x) f_{k-j}(y)$$

is a solution of the differential equation in (1). Indeed, the differential equation in (1) is equivalent to

$$\begin{aligned} \sum_{j=0}^k \binom{k}{j} f_j''(x) f_{k-j}(y) + \frac{A'(x)}{A(x)} \sum_{j=0}^k \binom{k}{j} f_j'(x) f_{k-j}(y) &= \\ &= \sum_{j=0}^k \binom{k}{j} f_j(x) f_{k-j}''(y) + \frac{A'(y)}{A(y)} \sum_{j=0}^k \binom{k}{j} f_j(x) f_{k-j}'(y), \end{aligned}$$

which is equivalent to

$$\begin{aligned} \sum_{j=0}^k \binom{k}{j} \left( f_j''(x) + \frac{A'(x)}{A(x)} f_j'(x) \right) f_{k-j}(y) &= \\ &= \sum_{j=0}^k \binom{k}{j} \left( f_{k-j}''(y) + \frac{A'(y)}{A(y)} f_{k-j}'(y) \right) f_j(x), \end{aligned}$$

that is, to

$$\begin{aligned} \sum_{j=0}^k \binom{k}{j} \left( \sum_{t=0}^j \binom{j}{t} c_t f_{j-t}(x) \right) f_{k-j}(y) &= \\ &= \sum_{j=0}^k \binom{k}{j} \left( \sum_{s=0}^{k-j} \binom{k-j}{s} c_s f_{k-j-s}(y) \right) f_j(x). \end{aligned}$$

But this equation holds true, since by choosing  $l = j + s$ , the right hand side is equal to

$$\sum_{l=0}^k \sum_{s=0}^l \binom{k}{l-s} \binom{k-(l-s)}{s} c_s f_{k-l}(y) f_{l-s}(x),$$

which is obviously equal to the left hand side. Moreover, the boundary value conditions in (1) are also satisfied, as

$$\partial_1 h(0, y) = \sum_{j=0}^k f'_j(0) f_{k-j}(y) = 0,$$

and

$$h(0, y) = \sum_{j=0}^k f_j(0) f_{k-j}(y) = f_k(y),$$

and similarly  $\partial_2 h(x, 0) = 0$ , and  $h(x, 0) = f_k(x)$ , hence  $h$  is the unique solution of the boundary value problem, which implies that  $h(x, y) = f(x * y)$ .

Conversely, suppose that the continuous functions  $f_k : \mathbb{R}_0 \rightarrow \mathbb{C}$  ( $k = 0, 1, \dots, N$ ) form a generalized moment sequence of order  $N$ . Then, by definition,  $f_0$  is an exponential and the conditions of (8) are satisfied. Now we proceed by induction and assume that (9) holds for  $f_0, f_1, \dots, f_k$ , with some positive integer  $k < N$ . We have

$$(10) \quad f_{k+1}(x * y) = \sum_{j=0}^{k+1} \binom{k+1}{j} f_j(x) f_{k+1-j}(y),$$

and by the definition of the hypergroup this implies that

$$\begin{aligned} \sum_{j=0}^{k+1} \binom{k+1}{j} f''_j(x) f_{k+1-j}(y) + \frac{A'(x)}{A(x)} \sum_{j=0}^{k+1} \binom{k+1}{j} f'_j(x) f_{k+1-j}(y) &= \\ &= \sum_{j=0}^{k+1} \binom{k+1}{j} f_j(x) f''_{k+1-j}(y) + \frac{A'(y)}{A(y)} \sum_{j=0}^{k+1} \binom{k+1}{j} f_j(x) f'_{k+1-j}(y). \end{aligned}$$

Rearranging the terms and using the induction hypothesis we have

$$\begin{aligned}
& \left( f''_{k+1}(x) + \frac{A'(x)}{A(x)} f'_{k+1}(x) \right) f_0(y) + \\
& \quad + \sum_{j=0}^k \binom{k+1}{j} \left( \sum_{t=0}^j \binom{j}{t} c_t f_{j-t}(x) \right) f_{k+1-j}(y) = \\
& = \left( f''_{k+1}(y) + \frac{A'(y)}{A(y)} f'_{k+1}(y) \right) f_0(x) + \\
& \quad + \sum_{j=1}^{k+1} \binom{k+1}{j} \left( \sum_{t=0}^{k+1-j} \binom{k+1-j}{t} c_t f_{k+1-j-t}(y) \right) f_j(x),
\end{aligned}$$

therefore

$$\begin{aligned}
& \left( f''_{k+1}(x) + \frac{A'(x)}{A(x)} f'_{k+1}(x) \right) f_0(y) + \sum_{j=0}^k \frac{k+1}{j} c_j f_0(x) f_{k+1-j}(y) + \\
& \quad + \sum_{j=1}^k \binom{k+1}{j} \left( \sum_{t=0}^{j-1} \binom{j}{t} c_t f_{j-t}(x) \right) f_{k+1-j}(y) = \\
& = \left( f''_{k+1}(y) + \frac{A'(y)}{A(y)} f'_{k+1}(y) \right) f_0(x) + \sum_{j=1}^{k+1} \binom{k+1}{j} c_{k+1-j} f_0(y) f_j(x) + \\
& \quad + \sum_{j=1}^k \binom{k+1}{j} \left( \sum_{t=0}^{k-j} \binom{k+1-j}{t} c_t f_{k+1-j-t}(y) \right) f_j(x).
\end{aligned}$$

It is easy to see, that the last terms on the two sides are equal:

$$\begin{aligned}
& \sum_{j=1}^k \binom{k+1}{j} \left( \sum_{t=0}^{j-1} \binom{j}{t} c_t f_{j-t}(x) \right) f_{k+1-j}(y) = \\
& = \sum_{t=0}^{k-1} \sum_{s=1}^{k-t} \binom{k+1}{s} \binom{k+1-s}{t} c_t f_{k+1-s-t}(x) f_s(y) = \\
& = \sum_{t=0}^{k-1} \sum_{s=1}^{k-t} \binom{k+1}{s+t} \binom{s+t}{t} c_t f_s(y) f_{k+1-s-t}(x) = \\
& = \sum_{j=1}^k \binom{k+1}{j} \left( \sum_{t=0}^{k-j} \binom{k+1-j}{t} c_t f_{k+1-j-t}(y) \right) f_j(x).
\end{aligned}$$

This means that

$$\begin{aligned} & \left( f''_{k+1}(x) + \frac{A'(x)}{A(x)} f'_{k+1}(x) - \sum_{j=1}^{k+1} \binom{k+1}{j} c_{k+1-j} f_j(x) \right) f_0(y) = \\ & = \left( f''_{k+1}(y) + \frac{A'(y)}{A(y)} f'_{k+1}(y) - \sum_{j=0}^k \binom{k+1}{j} c_j f_{k+1-j}(y) \right) f_0(x) \end{aligned}$$

holds for each positive  $x$  and  $y$ , hence there exists a complex number  $c_{k+1}$  such that

$$f''_{k+1}(x) + \frac{A'(x)}{A(x)} f'_{k+1}(x) - \sum_{j=1}^{k+1} \binom{k+1}{j} c_{k+1-j} f_j(x) = c_{k+1} f_0(x).$$

As a consequence of (10) we also have  $f_{k+1}(0) = 0$ , and due to

$$0 = \sum_{j=0}^{k+1} \binom{k+1}{j} f_j(x) f'_{k+1-j}(0) = f_0(x) f'_{k+1}(0)$$

we get that  $f'_{k+1}(0) = 0$ . Hence (9) holds for  $k+1$  and the theorem is proved by induction.

**Theorem 6.2.** *Let  $K = (\mathbb{R}_0, A)$  be the Sturm-Liouville hypergroup corresponding to the Sturm-Liouville function  $A$  with the exponential family  $\varphi$  and let  $N$  be a positive integer. The continuous functions  $f_k : \mathbb{R}_0 \rightarrow \mathbb{C}$  ( $k = 0, 1, \dots, N$ ) form a sequence of generalized moment functions of order  $N$  on the hypergroup  $K$  if and only if there are complex numbers  $c_0, c_1, \dots, c_N$  such that*

$$(13) \quad f_k(x) = \partial_t^k \varphi(x, f(t)) \Big|_{t=0}$$

holds for each  $x$  in  $\mathbb{R}_0$ , where  $\varphi$  is the exponential family of the Sturm-Liouville hypergroup  $K$  and

$$f(t) = \sum_{j=0}^N c_j \frac{t^j}{j!}$$

for each  $t$  in  $\mathbb{R}$ .

**Proof.** Let  $k$  be in  $\{0, 1, \dots, N\}$ , let  $\varphi$  be the exponential family of the hypergroup  $K$ , and let  $f$  be the function given in the theorem. If we take

$\lambda = f(t)$  in (4) and differentiate the equation  $k$ -times with respect to  $t$  we get that

$$(14) \quad \partial_x^2 \partial_t^k \varphi(x, f(t)) + \frac{A'(x)}{A(x)} \partial_x \partial_t^k \varphi(x, f(t)) = \sum_{j=0}^k \binom{k}{j} f^{(j)}(t) \partial_t^{k-j} \varphi(x, f(t)).$$

Taking  $t = 0$  we have for  $f_k(x) = \partial_t^k \varphi(x, f(t))|_{t=0}$  the following equation

$$f_k''(x) + \frac{A'(x)}{A(x)} f_k'(x) = \sum_{j=0}^k \binom{k}{j} c_j f_{k-j}(x),$$

furthermore  $f_0(0) = 1$ ,  $f_0'(0) = 0$ , and  $f_k(0) = 0$ ,  $f_k'(0) = 0$  in case of  $k \neq 0$ . This means that all the conditions of Theorem 6.1 are satisfied and  $f_0, f_1, \dots, f_N$  form a generalized moment sequence.

To prove the converse we assume that the functions  $f_0, f_1, \dots, f_N$  form a generalized moment sequence and we prove by induction. It is obvious that the statement is true for  $f_0$  and we suppose that  $f_j(x) = \partial_t^j \varphi(x, f(t))|_{t=0}$  for  $j = 0, 1, \dots, k$ , where  $k$  is in  $\{1, 2, \dots, N\}$ . We consider the function

$$g(x) = f_{k+1}(x) - \partial_t^{k+1} \varphi(x, f(t))|_{t=0}$$

for each positive  $x$ . Then the expression  $g''(x) + \frac{A'(x)}{A(x)} g'(x)$  is equal to

$$f_{k+1}''(x) + \frac{A'(x)}{A(x)} f_{k+1}'(x) - \partial_x^2 \partial_t^{k+1} \varphi(x, f(t))|_{t=0} - \frac{A'(x)}{A(x)} \partial_x \partial_t^{k+1} \varphi(x, f(t))|_{t=0}$$

and using Theorem 6.1 and (14) we get

$$\begin{aligned} c_0 f_{k+1}(x) + \sum_{j=1}^{k+1} \binom{k+1}{j} c_j \partial_t^{k+1-j} \varphi(x, f(t))|_{t=0} - \\ - \sum_{j=0}^{k+1} \binom{k+1}{j} c_j \partial_t^{k+1-j} \varphi(x, f(t))|_{t=0} = c_0 g(x). \end{aligned}$$

Consequently

$$g''(x) + \frac{A'(x)}{A(x)} g'(x) = c_0 g(x), \quad g(0) = 0, \quad g'(0) = 0,$$

hence  $g(x) \equiv 0$  and the proof is complete.

## References

- [1] **Aczél J.**, Functions of binomial type mapping groupoids into rings, *Math. Zeitschrift*, **154** (1977), 115-124.
- [2] **Bloom W.R. and Heyer, H.**, *Harmonic analysis of probability measures on hypergroups*, de Gruyter, Berlin, 1995.
- [3] **Gallardo L.**, Some methods to find moment functions on hypergroups, *Harmonic analysis and hypergroups*, eds. Anderson J.M., Litvinov G.L., Ross K.A., Singh A.I., Sunder V.S. and Wildberger N.J., de Gruyter Studies in Mathematics, de Gruyter, Birkhäuser, Boston, Basel, Berlin, 1998, 13-31.
- [4] **Orosz Á. and Székelyhidi L.**, Moment functions on polynomial hypergroups in several variables, *Publ. Math. Debrecen*, **65** (3-4) (2004), 429-438.
- [5] **Orosz Á. and Székelyhidi L.**, Moment functions on polynomial hypergroups, *Arch. Math.*, **85** (2005), 141-150.
- [6] **Ross K.A.**, Hypergroups and signed hypergroups, *Harmonic analysis on hypergroups*, eds. Anderson J.M., Litvinov G.L., Ross K.A., Singh A.I., Sunder V.S. and Wildberger N.J., de Gruyter Studies in Mathematics, de Gruyter, Birkhäuser, Boston, Basel, Berlin, 1998, 77-91.
- [7] **Székelyhidi L.**, Functional equations on hypergroups, *Functional equations, inequalities and applications*, ed. Rassias Th.M., Kluwer Academic Publishers, 2003, 167-181.
- [8] **Zeuner H.**, Moment functions and laws of large numbers on hypergroups, *Math. Zeitschrift*, **211** (1992), 369-407.

**Á. Orosz and L. Székelyhidi**

Institute of Mathematics

University of Debrecen

H-4010 Debrecen, P.O.B. 12

Hungary

[oagota@math.klte.hu](mailto:oagota@math.klte.hu)

[szekely@math.klte.hu](mailto:szekely@math.klte.hu)