GROUPS ADMITTING ONLY QUASI–PERIODIC FACTORIZATIONS

S. Szabó (Pécs, Hungary)

Dedicated to Imre Kátai on his 70th birthday

Abstract. We will show that if a finite abelian group is a direct product of two subgroups of relatively prime orders and one of the subgroups has prime order, then each normalized factorization of the group must be quasiperiodic.

1. Introduction

Let G be a finite abelian group and let A, B be a subsets of G. If each $g \in G$ can be written in the form

(1)
$$g = ab, \quad a \in A, \quad b \in B,$$

then the product AB is equal to G. If each $g \in G$ can be written uniquely in the form (1), then we say that the product AB is direct and we refer to the equation G = AB as a factorization of G.

If the identity element e belongs to the subset e A of G, then we say that A is a normalized subset of G. If both factors A and B of a factorization G = AB are normalized subsets we call the factorization a normalized factorization.

A subset A of G is defined to be periodic if there is a $g \in G \subset \{e\}$ such that Ag = A. Note that if g and h are periods of A, then so is gh provided $gh \neq e$. It follows then that the periods of A together with the identity element

Mathematics Subject Classification: 20K01 (05B45, 52C22, 68R05)

form a subgroup H of G. We will call H the subgroup of periods of A. The reader can verify that there is a subset A_1 of G such that the product A_1H is direct and is equal to A. If A is normalized then $H \subset A$ and A_1 can be chosen to be a subset of A. If one of the factors in the factorization G = AB is periodic, then we call the factorization periodic.

We say that a partition $B = B_1 \cup \cdots \cup B_n$ is a regular partition of the subset B of G if there is a periodic subset $C = \{c_1, \ldots, c_n\}$ of G such that $c_1 = e$, $|C| \ge 2$ and $AB_i = AB_1c_i$ for each i, $1 \le i \le n$. According to [2] we call a normalized factorization G = AB quasi-periodic if one of the factors has a regular partition.

In [3] one can find another definition of the quasi-periodic factorization. Namely, a partition $B = B_1 \cup \cdots \cup B_n$ of the subset B of G is defined to be regular if there is a subgroup $H = \{h_1, \ldots, h_n\}$ of G such that $h_1 = e$, $|H| \ge 2$ and $AB_i = AB_1h_i$ for each $i, 1 \le i \le n$. The normalized factorization G = ABis called quasi-periodic if at least one of the factors A, B has a regular partition. The two definitions are equivalent. A proof can be found in [7].

In 1950 G. Hajós [2] asked if each factorization of each finite abelian group is quasi-periodic. In 1974 A.D. Sands [4] showed that an earlier factorization construction of N.G. de Bruijn [1] is a counter-example for the Hajós question. It was verified in [8] that another factorization construction of N.G. de Bruijn is not quasi-periodic either. In [7] a large family of nonquasiperiodic factorizations was exhibited.

One might wonder if each normalized factorization of a finite abelian group G is quasi-periodic, then does it hold for each subgroup H of G. A.D. Sands [6] showed that for this question the answer is "no". Using Sands' method in this paper we will show that if a finite abelian group G is a direct product of its subgroups H and K such that |H|, |K| are relatively prime and |K| is a prime, then G admits only quasi-periodic normalized factorizations.

2. The result

Let G be a finite abelian group and let H be a subgroup of G. If f_1, \ldots, f_n is a complete set of representatives of cosets modulo H, then the cosets Hf_1, \ldots, Hf_n form a partition of G. Therefore for a given subset A of G the sets $A_1 = A \cap Hf_1, \ldots, A_n = A \cap Hf_n$ form a partition of A. We will use this observation in the proof of the next theorem. The other observation we will use is that G = AB is a factorization of G if and only if the sets Ab, $B \in B$ form a partition of G.

Theorem 1. Let p be a prime. Let G be a finite abelian group that is the direct product of its subgroups H and K. If p does not divide |H| and |K| = p, then each normalized factorization of G is quasi-periodic.

Proof. Let G = AB be a normalized factorization of G. Since the order of G is equal to the product of the orders of A and B we may assume that $p \not||A|$ and p||B|. Let $K = \langle f \rangle$. The factor A can be written in the form

$$A = A_0 \cup A_1 \cup \cdots \cup A_{p-1},$$

where $A_i = A \cap Hf^i$. Further there are subsets $D_0, \ldots, D_{p-1} \subset H$ such that $A_i = D_i f^i$. Thus

$$A = D_0 \cup D_1 f^1 \cup \dots \cup D_{p-1} f^{p-1}.$$

Similarly B can be written in the form

$$B = B_0 \cup B_1 \cup \cdots \cup B_{p-1},$$

where $B_i = B \cap Hf^i$. There are subsets $C_0, \ldots, C_{p-1} \subset H$ such that $B_i = C_i f^i$. Therefore

$$B = C_0 \cup C_1 f^1 \cup \dots \cup C_{p-1} f^{p-1}.$$

We claim that $AB_v = AB_0f^v$ holds for each $v, 1 \le v \le p-1$. Using the above notations the claim is equivalent to

$$(D_0 \cup D_1 f^1 \cup \dots \cup D_{p-1} f^{p-1})(C_v f^v) = (D_0 \cup D_1 f^1 \cup \dots \cup D_{p-1} f^{p-1})(C_0) f^v,$$

that is, equivalent to

$$D_0C_v \cup D_1C_vf^1 \cup \dots \cup D_{p-1}C_vf^{p-1} = D_0C_0 \cup D_1C_0f^1 \cup \dots \cup D_{p-1}C_0f^{p-1}.$$

In fact we will verify that $D_u C_v = D_u C_0$ holds for each $u, v, 0 \le u \le p-1, 1 \le v \le p-1$.

Choose an integer t(i) such that

$$t(i) \equiv 1 \pmod{|H|},$$
$$t(i) \equiv i \pmod{|K|}.$$

Such a t(i) exists for each $i, 0 \leq i \leq p-1$. By Proposition 3 of [5], in the factorization G = AB the factor A can be replaced by $A^{t(i)}$ to get the normalized factorization $G = A^{t(i)}B$. Note that

$$A^{t(i)} = D_0^{t(i)} \cup D_1^{t(i)} f^{t(i)} \cup \dots \cup D_{p-1}^{t(i)} f^{(p-1)t(i)} =$$

= $D_0 \cup D_1 f^i \cup \dots \cup D_{p-1} f^{(p-1)i}.$

The factorization

$$G = (D_0 \cup D_1 f^i \cup \dots \cup D_{p-1} f^{(p-1)i})B$$

provides that the sets

(2)
$$D_0B, D_1Bf^i, \dots, D_{p-1}Bf^{(p-1)i}$$

form a partition of G. The i = 0 and i = 1 special cases of (2) give that

(3)
$$D_1 B \cup D_2 B \cup \cdots \cup D_{p-1} B = D_1 B f \cup D_2 B f^2 \cup \cdots \cup D_{p-1} f^{p-1} B.$$

Choose an integer $u, 1 \le u \le p-1$. Note that

$$D_u B \cap D_w B f^w = \emptyset$$

for each $w,\ 1\leq w\leq p-1,\ w\neq u.$ In order to verify the claim assume on the contrary that

$$D_u B \cap D_w B f^w = \emptyset$$

for some u and w. Multiplying by f^{ui} we get

(4) $D_u B f^{ui} \cap D_w B f^{w+iu} = \emptyset.$

The congruence

$$wi \equiv w + iu \pmod{p}$$

is solvable for i and so (4) contradicts to (2). This contradiction proves our claim.

From (3) it follows that $D_u B = D_u B f^u$. Using this equation repeatedly we get that

$$D_u B = D_u B f = D_u B f^2 = \dots = D_u B f^{p-1}.$$

Choose an integer $v, 1 \le v \le p-1$. The equation $D_u B = D_u B f^{p-v}$ can be written in the form

$$D_u(C_0 \cup C_1 f^1 \cup \dots \cup C_{p-1} f^{p-1}) = D_u(C_0 f^{p-v} \cup C_1 f^{p-v+1} \cup \dots \cup C_{p-1} f^{p-v+p-1}),$$

that is, in the form

$$D_u C_0 \cup D_u C_1 f^1 \cup \dots \cup D_u C_{p-1} f^{p-1} =$$
$$D_u C_0 f^{p-v} \cup D_u C_1 f^{p-v+1} \cup \dots \cup D_u C_{p-1} f^{p-v+p-1}.$$

Intersecting by H we get $D_u C_0 = D_u C_v$. This holds for each $u, v, 1 \le u \le \le p-1, 1 \le v \le p-1$. (The u = 0 case is not covered.)

It remains to verify that $D_0C_v = D_0C_0$ holds for each v, $1 \le v \le p-1$. In fact we will show that $D_uC_0D_uC_v$ holds for each $u, v, 0 \le u \le p-2, 1 \le v \le p-1$. Multiplying the factorization G = AB by f we get the normalized factorization G = (Af)B. Here

$$Af = D_0 f \cup D_1 f^2 \cup \dots \cup D_{p-2} f^{p-1} \cup D_{p-1}.$$

(Now D_{p-1} will play the earlier role of D_0 .) Repeating the whole argument in this setting we get the required result.

References

- de Bruijn N.G., On the factorization of finite abelian groups, Indag. Math. Kon. Ned. Akad. Wetersch., 15 (1953), 258-264.
- [2] Hajós G., Sur le problème de factorisation des groupes cycliques, Acta Math. Acad. Sci. Hungar., 1 (1950), 189-195.
- [3] Fuchs L., Abelian groups, Akadémia Kiadó, Budapest 1958.
- [4] Sands A.D., On a conjecture of G. Hajós, *Glasgow Math. J.*, 15 (1974), 88-89.
- [5] Sands A.D., Replacement of factors by subgroups in the factorization of abelian groups, Bull. London Math. Soc., 32 (2000), 297-304.
- [6] Sands A.D., A note on quasi-periodic factorization (submitted)
- [7] Sands A.D. and Szabó S., Nonquasi-periodic factorizations for certain finite abelian groups, Int. J. of Algebra and Computation, 17 (2007), 837-849.
- [8] Szabó S., Quasi-periodic factorizations, J. of Algebra and its Applications, 6 (2006), 793-798.

S. Szabó

Institute of Mathematics and Informatics University of Pécs Ifjúság u. 6 H-7624 Pécs, Hungary sszabo7@hotmail.com