

SOME REMARKS ON A RESULT OF F. LUCA AND I. SHPARLINSKI

I. Kátai (Budapest, Hungary)

Abstract. Luca and Shparlinski proved that $n^{1/\omega(n)} \pmod{1}$ is uniformly distributed. Generalizations are proved here. $\omega(n)$ = number of prime divisors of n .

1. Introduction

Luca and Shparlinski investigated the sequences

$$n^{1/\omega(n)} \pmod{1}, \quad n^{1/\Omega(n)} \pmod{1}, \quad \left(\prod_{p|n} \right)^{1/\omega(n)} \pmod{1}$$

and proved that counted them up to $(n \leq) x$, the discrepancy is $\frac{1}{(\log x)^{1+o(1)}}$. Here and in the sequel $\omega(n)$ is the number of prime divisors, $\Omega(n)$ is the number of prime power divisors of n , p runs over the set of primes.

The main idea of the proof of the first sequence is to write the integers n as $n = mp$, where p is the largest prime divisor of n , then for fixed m count the distribution of $m^{\frac{1}{\omega(m)+1}} p^{\frac{1}{\omega(m)+1}} \pmod{1}$, where p runs over the set of primes p in the interval $P(m) < p < \frac{x}{m}$, where in general $P(n)$ = the largest prime divisor of n .

Mathematics Subject Classification: 11J71, 11K65

The European Union and the European Social Fund have provided financial support to the project under the grant agreement TÁMOP-4.2.1/B-09/1/KMR-2010-003.

The distribution of

$$p^{\frac{1}{\omega(m)+1}} \pmod{\left(\frac{1}{m^{1/\omega(m)+1}}\right)}$$

can be computed by using the short interval version of the theorem of Huxley [2] and Heath-Brown [3], according to

$$\pi(x+y) - \pi(x) = \frac{y}{\log x} \left(1 + O\left(\frac{(\log \log x)^4}{\log x}\right) \right).$$

Similar assertions with some weaker error term can be proved by other "arithmetical" sequences. These depend on the following obvious Lemma 1 and nontrivial theorems for short interval version of sums of multiplicative functions. Let $\pi_k(x) = \#\{n \leq x \mid \omega(n) = k\}$.

Lemma 1. *Let $\kappa(n)$ ($n = 1, 2, \dots$) be a sequence of real numbers, $0 < \kappa(n) < 1$. Assume that for $X > X_0$, the set of integers $n \in [X, 2X)$ can be classified into disjoint sets $\mathcal{I}_1, \mathcal{I}_2, \dots, \mathcal{I}_t$, $\mathcal{T}$ such that in the notation*

$$S_j(x) = \sum_{\substack{n \leq x \\ n \in \mathcal{I}_j}} 1, \quad T(x) = \sum_{\substack{n \leq x \\ n \in \mathcal{T}}} 1$$

the following conditions hold:

- (1) $\kappa(n) = \kappa_k$ is fixed for $n \in \mathcal{I}_k$, $k = 1, \dots, t$.
- (2) $T(2X) \ll X\delta(X)$, where $\delta(x)$ is a monotonically decreasing function defined in $1 < x < \infty$ such that $\delta(\sqrt{x}) \ll \delta(x)$,
- (3)

$$\begin{aligned} \max_{\substack{\frac{\delta(X)}{\kappa_k} x^{1-\kappa_k} \leq h \leq x}} \max_{x \in [X, 2X]} \left| \frac{S_k(x+h) - S_k(x)}{h} - \frac{S_k(2X) - S_k(X)}{X} \right| &\ll \\ &\ll \frac{S_k(2X)}{X} \cdot \delta(X). \end{aligned}$$

Then the discrepancy of the sequence $n^{\kappa(n)} \pmod{1}$ ($n = 1, \dots, [x]$) is $\mathcal{O}(\delta(x))$.

2. Short interval version of sums of some multiplicative function

By using the so called Hooley-Huxley contour for functions depending on Dirichlet L -function, Ramachandra [4] proved a general theorem:

Let S_1, S_2, S_3 be the sets of L -series, the derivatives, and the logarithms of L -series, respectively. $\log L(s, \chi)$ is defined by analytic continuation from the halfplane $\sigma = \operatorname{Re} s > 1$; for some complex z , we define

$$L(s, \chi)^z = \exp(z \log L(s, \chi)).$$

Let $P_1(s)$ be any finite product (with complex exponents) of functions of S_1 . Let $P_2(s)$ be any finite power product (with nonnegative integral exponents) of functions of S_2 . Let also $P_3(s)$ denote any finite power product with nonnegative exponents of S_3 . Let c_n be a sequence of complex numbers such that $|c_n| \ll n^\varepsilon$ for every $\varepsilon > 0$ and

$$\sum \frac{|c_n|}{n^\sigma} < \infty \quad \text{for } \sigma > 1/2.$$

Let $F_0(s) = \sum_n \frac{c_n}{n^s}$. Furthermore, let

$$F_1(s) = P_1(s) P_2(s) P_3(s) F_0(s) = \sum \frac{g_n}{n^s},$$

and

$$E(x) = \sum_{n \leq x} g_n.$$

Let $0 < r < 1/2$. We define the contour $C(r)$ by starting from the circle $\{s \mid |s-1| = r\}$, removing the point $1-r$, and proceeding on the remaining portion of the circle in the clockwise direction. Let $C_0 = C(r)$. Assume that r is so small that $F_1(s)$ has no singularities on the boundary and in interior of it, except, possibly, the place $s = 1$.

Let $N_\chi(\alpha, T)$ be the number of roots ϱ of $L(s, \chi)$ in the region $\operatorname{Re} \varrho \geq \alpha$, $|\operatorname{Im} \varrho| \leq T$. Assume that

$$N_\chi(\alpha, T) = \mathcal{O}\left(T^{B(1-\alpha)} (\log T)^2\right)$$

holds for all characters occurring in P_1, P_2 and P_3 .

Let $\varphi = 1 - 1/B + \varepsilon$.

Remark. According to Huxley's result, φ can be any constant greater than $\frac{7}{12}$.

Theorem of Ramachandra. *Let x be sufficiently large and $1 \leq h \leq x$. Let*

$$I(x, h) = \frac{1}{2\pi i} \int_0^h \left(\int_{C_0} F_1(s) (v+x)^{s-1} ds \right) dv.$$

Then

$$E(x+h) - E(x) = I(x, h) + \mathcal{O}_\varepsilon \left(h \cdot \exp \left(-(\log x)^{1/6} \right) + x^\varphi \right).$$

In [5] it was deduced from the above theorem that

$$\sum_{\substack{x \leq n \leq x+h \\ \omega(n)=k}} 1 = (1 + o(1)) \frac{h}{x} \pi_k(x)$$

uniformly for any $k \leq \log \log x + c_x \sqrt{\log \log x}$, where $c_x \rightarrow \infty$ sufficiently slowly, if $x^{\frac{7}{12}+\varepsilon} \leq h \leq x$. Sankaranarayanan and Srinivas [8] gave a version of Ramachandra's result in which the function $F_1(s)$ may depend on a parameter.

In our paper [7] we deduced the following theorem: Let $C_1 = C \left(\frac{1}{\log x} \right)$, and let L^-, L^+ be defined as the intervals of straightlines

$$L^- = \left[\left(1 - \frac{1}{r} \right) e^{-i\pi}, \left(1 - \frac{1}{\log x} \right) e^{-i\pi} \right],$$

$$L^+ = \left[\left(1 - \frac{1}{\log x} \right) e^{i\pi}, \left(1 - \frac{1}{r} \right) e^{i\pi} \right].$$

Let C^* be the contour going along L^- starting from $\left(1 - \frac{1}{r} \right) e^{-i\pi}$, then on C_1 , and finally, on L^+ .

Theorem (1 and 2, in [7]). *Assume that $F_1(s)$ satisfies the conditions stated in Ramachandra's theorem. Let $r > 0$ and $\varepsilon > 0$ be constants so that $\frac{7}{12} + \varepsilon < \frac{2}{3} - \frac{2r}{3}$, and that $U(s) := F_1(s) (s-1)^{-z}$ is analytic in the disc $|s-1| \leq r$. Consequently $U(s) = A_0 + A_1(s-1) + \dots + A_k(s-1)^k + (s-1)^{k+1} V(s)$ holds for every fixed k , where $V(s)$ is bounded in $|s-1| \leq r$.*

If $x^{\frac{7}{12}+\varepsilon} \leq h \leq x^{\frac{2}{3}-\frac{2\varepsilon}{3}}$, then

$$\begin{aligned}
 \frac{E(x+h) - E(x)}{h} &= \frac{1}{2\pi i} \int_{C^*} F_1(s) x^{s-1} ds + \mathcal{O}\left(\exp\left(-(\log x)^{1/6}\right)\right) = \\
 (2.1) \qquad &= \sum_{l=0}^k A_l \frac{\Gamma(l-z)}{(\log x)^{l-z+1}} \cdot \frac{(-1)^{l+1} \sin \pi}{\pi} + \\
 &\quad + \mathcal{O}\left(\frac{1}{(\log x)^{k+2-\operatorname{Re} z}}\right),
 \end{aligned}$$

whenever $\operatorname{Re} z \leq k+1$.

3. Short interval versions of the theorems of Sathe and Selberg

Let furthermore

$$\pi_l(x) := \#\{n \leq x \mid \omega(n) = l\}; \quad N_l(x) := \#\{n \leq x \mid \Omega(n) = l\}.$$

Further we shall use the abbreviation: $x_1 = \log x$, $x_2 = \log x_1$, etc. By using our theorem for

$$\begin{aligned}
 F_1(s) &= \sum \frac{z^{\omega(n)}}{n^s} = \zeta^z(s) h(s), \\
 h(s) &= \prod_p \left(1 - \frac{1}{p^s}\right)^z \left(1 - \frac{z}{p^s - 1}\right)
 \end{aligned}$$

($h(s)$ is regular in $\operatorname{Re} s > 1/2$ for arbitrary $z \in \mathbb{C}$), and for

$$\begin{aligned}
 F_1(s) &= \sum \frac{z^{\Omega(m)}}{m^s} = \zeta^z(s) h(s), \\
 h(s) &= \prod_p \left(1 - \frac{z}{p^s}\right)^{-1} \left(1 - \frac{1}{p^s}\right)^z
 \end{aligned}$$

($h(s)$ is regular in $\operatorname{Re} s > 1/2$, whenever $|z| \leq 2 - \delta$, $\sigma > 0$).

We deduced the following assertions which will be quoted now as Lemma 2 and 3.

Lemma 2. *Let $c_1 > 0$, $\varepsilon > 0$ be arbitrary positive constants, r be another constant so that $\frac{7}{12} + \varepsilon \leq \frac{2}{3} - \frac{2r}{3}$. Then, for $|z| \leq c_1$, $x^{\frac{7}{12} + \varepsilon} \leq h \leq x^{\frac{2}{3} - \frac{2r}{3}}$ we have*

$$h^{-1} \sum_{x \leq m \leq x+h} z^{\omega(m)} = \varphi(z) (\log x)^{z-1} + \mathcal{O}\left((\log x)^{\operatorname{Re} z - 2}\right),$$

where

$$\varphi(z) = \frac{1}{\Gamma(z+1)} \prod_p \left(1 - \frac{1}{p}\right)^z \left(1 + \frac{z}{p-1}\right).$$

Lemma 3. *Let $|z| \leq 2 - \delta$, $0 < \delta < 1$ be an arbitrary constant, ε, r be as in Lemma 2, $x^{\frac{7}{12} + \varepsilon} \leq h \leq x^{\frac{2}{3} - \frac{2r}{3}}$. Then*

$$h^{-1} \sum_{x \leq m \leq x+h} z^{\Omega(m)} = G(z) (\log x)^{z-1} + \mathcal{O}\left((\log x)^{\operatorname{Re} z - 2}\right),$$

$$G(z) = \frac{1}{\Gamma(z+1)} \prod_p \left(1 - \frac{z}{p}\right)^{-1} \left(1 - \frac{1}{p}\right)^z.$$

Lemma 4. *Let c be an arbitrary constant, $\varepsilon, r > 0$ be such constants for which $\frac{7}{12}\varepsilon < \frac{2}{3} - \frac{2r}{3}$ (i.e. $r < \frac{3}{4} - \frac{3}{2}\varepsilon$). Then, uniformly as $1 \leq k \leq \leq cx_2$, $x^{\frac{7}{12} + \varepsilon} \leq h \leq x^{\frac{2}{3} - \frac{2r}{3}}$, we have*

$$\pi_k(x+h) - \pi_k(x) = \frac{h}{x_1} \cdot \frac{x_2^{k-1}}{(k-1)!} \left\{ \varphi\left(\frac{k-1}{x_2}\right) + \mathcal{O}\left(\frac{k}{x_2^2}\right) \right\}.$$

It is known (see e.g. in J. Kubilius [9], or in G. Tenenbaum [10]) that under the conditions stated for $|z|$ in Lemma 2 and 3,

$$\frac{1}{x} \sum_{m \leq x} z^{\Omega(m)} = G(z) (\log x)^{z-1} + \mathcal{O}\left((\log x)^{\operatorname{Re} z - 2}\right),$$

$$\frac{1}{x} \sum_{m \leq x} z^{\omega(m)} = \varphi(z) (\log x)^{z-1} + \mathcal{O}\left((\log x)^{\operatorname{Re} z - 2}\right),$$

consequently

$$\left| \frac{1}{h} \sum_{x \leq m \leq x+h} z^{\Omega(m)} - \frac{1}{x} \sum_{m \leq x} z^{\Omega(m)} \right| \ll (\log x)^{\operatorname{Re} z - 2},$$

$$\left| \frac{1}{h} \sum_{x \leq m \leq x+h} z^{\omega(m)} - \frac{1}{x} \sum_{m \leq x} z^{\omega(m)} \right| \ll (\log x)^{\operatorname{Re} z - 2}$$

and by integrating with respect to z on the unit circle we obtain that

$$(3.1) \quad \max_k \left| \frac{\pi_k(x+h) - \pi_k(x)}{h} - \frac{\pi_k(x)}{x} \right| \ll \frac{1}{\log x},$$

$$(3.2) \quad \max_k \left| \frac{N_k(x+h) - N_k(x)}{h} - \frac{N_k(x)}{x} \right| \ll \frac{1}{\log x}.$$

(3.1), (3.2) is proved under the condition $h \leq x^{\frac{2}{3} - \frac{2r}{3}}$. By using the formula (17) and (19) in [10] Chapter II.6., page 205 we obtain that

$$(3.3) \quad \max_{x < y < 2x} \left| \frac{\pi_k(x)}{x} - \frac{\pi_k(y)}{y} \right| \ll \frac{1}{\log x}$$

uniformly as $k \leq Ax_2$, and that

$$(3.4) \quad \max_{x < y < 2x} \left| \frac{N_k(x)}{x} - \frac{N_k(y)}{y} \right| \ll \frac{1}{\log x},$$

uniformly as $k \leq (2 - \delta)x_2$. Hence we obtain easily

Lemma 5. *(3.1) is true uniformly as $x^{\frac{7}{12} + \varepsilon} \leq h \leq x$, $k \leq Ax_2$; (3.2) is true uniformly as $x^{\frac{7}{12} + \varepsilon} \leq h \leq x$, $k \leq (2 - \delta)x_2$.*

4. Main result

Theorem 4. *Let $Q(n) = a_0 n^m + a_1 n^{m-1} + \dots + a_m \in R[x]$ be a polynomial, $a_0 > 0$.*

(1) *Assume that for every $X \geq 2$ for $n \in [X, 2X]$ $\kappa(n)$ depends only on $\omega(n)$ if $\omega(n) \leq e \log \log X$.*

Assume furthermore that

$$(4.1) \quad \frac{5}{12m} - \varepsilon > \kappa(n) > \frac{1}{(\log X)^{1-\delta}} \quad \text{for } n \in [X, 2X]$$

satisfying $\omega(n) \leq e \log \log X$. Then the discrepancy of the sequence $Q(n)^{\kappa(n)} \pmod{1}$ ($n \leq x$) is $\mathcal{O}\left(\frac{x_2}{x_1}\right)$.

- (2) Assume that for every $X \geq 2$ for $n \in [X, 2X]$ $\kappa(n)$ depends only on $\Omega(n)$ if $\Omega(n) \leq \beta \log \log X$, where $1 < \beta < 2$. Assume furthermore that (4.1) holds if $\Omega(n) \leq \beta \log \log X$. Then the discrepancy of $Q(n)^{\kappa(n)} \pmod{1}$ ($n \leq x$) is $\mathcal{O}\left(\frac{1}{x_1} x_1^{\beta \log \frac{e}{\beta}}\right)$.

Remark. The proof is a straightforward consequence of (3.1),(3.2) since the sums

$$(4.2) \quad \begin{aligned} & \# \left\{ n | n \in \mathcal{P}_k, \quad A < Q^{\kappa(n)}(n) < A + \Delta \right\} = \\ & = \# \left\{ n | n \in \mathcal{P}_k, \quad Q^{-1} \left(A^{\frac{1}{\kappa(n)}} \right) < n < Q^{-1} \left((A + \Delta)^{\frac{1}{\kappa(n)}} \right) \right\} \end{aligned}$$

and similarly

$$(4.3) \quad \begin{aligned} & \# \left\{ n | n \in \mathcal{N}_k, \quad A < Q^{\kappa(n)}(n) < A + \Delta \right\} = \\ & = \# \left\{ n | n \in \mathcal{N}_k, \quad Q^{-1} \left(A^{\frac{1}{\kappa(n)}} \right) < n < Q^{-1} \left((A + \Delta)^{\frac{1}{\kappa(n)}} \right) \right\} \end{aligned}$$

can be estimated by error $\mathcal{O}\left(\frac{1}{\log x}\right)$, uniformly as $\frac{1}{\log x} \leq \Delta \leq 1$, uniformly as $k \leq ex_2$ in the case (4.2), and as $k \leq \beta x_2$ in the case (4.3).

Furthermore, we observe that

$$\# \{n \leq x | \omega(n) > ex_2\} \ll \frac{x}{x_1},$$

$$\# \{n \leq x | \Omega(n) > \beta x_2\} \ll \frac{x}{x_1} x_1^{\beta \log \frac{e}{\beta}}.$$

The theorem follows from these observations easily. We omit the details.

5. Further remark

Let $\tau(n)$ be the number of divisors of n . Let K run over the set square-full numbers, $\mathcal{M}_K$ be the set of square-free integers m which are coprime to K . Every integer $n \in \mathbb{N}$ can be uniquely written as $n = Km$, where K is the square-full part and m is the square-free part of n .

Let $\mathcal{M}_{K,l}$ be the set of those integers n the square-full part of which is K , the square-free part is m , and $\omega(m) = l$. Let

$$M_{K,l}(x) = \#\{n \leq x, n \in \mathcal{M}_{K,l}\}.$$

By using the method applied in [11] one can prove that

$$\left| \frac{M_{K,l}(x+h) - M_{K,l}(x)}{h} - \frac{M_{K,l}(x)}{x} \right| \ll \frac{1}{K(\log x)^{1/2}}$$

uniformly as

$$x^{\frac{7}{12}+\varepsilon} \leq h \leq x^{0.66}, \quad K \leq x_1^4, \quad |x_2 - l| \leq x_2^{3/4}.$$

Hence it follows

Theorem 2. *Let Q be as in Theorem (1). Then the sequence $Q(n)^{1/\tau(n)}$ is uniformly distributed (mod 1).*

We omit the proof.

References

- [1] **Luca F. and Shparlinski I.E.**, Uniformity of distribution modulo 1 of the geometric mean prime divisor, *Bol. Soc. Matem. Mexicana*, **12** (2006), 155-163.
- [2] **Huxley M.N.**, On the difference between consecutive primes, *Invent. Math.*, **15** (1972), 164-170.
- [3] **Heath-Brown D.R.**, The number of primes in a short interval, *J. Reine und Angewandte Mathematik*, **389** (1988), 22-63.
- [4] **Ramachandra K.**, Some problems on analytic number theory, *Acta Arithmetica*, **31** (1976), 313-324.
- [5] **Kátai I.**, A remark on a paper of Ramachandra, *Number Theory. Proc. Ootacamund*, ed. K. Alladi, Lecture Notes in Math. 1122, Springer Verlag, 1984, 147-152.
- [6] **Kátai I. and Subbarao M.V.**, Some remarks on a paper of Ramachandra, *Liet. matem. rink.*, **43** (4) (2003), 497-506.
- [7] **Kátai I. and Subbarao M.V.**, The distribution of integers with given number of prime factors in almost all short intervals, *The Riemann zeta function and related themes. Proc. of Int. Conf. held at National Institute*

- of Advanced Studies, Bangalore, 13-15 December, 2003*, Lecture Notes Series **2**, Ramanujan Mathematical Society, 2006, 115-120.
- [8] **Sankaranarayanan A. and Srinivas K.**, On the papers of Ramachandra and Kátai, *Acta Arith.*, **624** (1992), 373-382.
- [9] **Кубилюс Й.**, *Вероятностные методы в теории чисел*, Гос. изд. полит. и науч. лит. ЛитССР, Вильнюс, 1962. (**Kubilius J.P.**, *Probabilistic methods in the theory of numbers*, Amer. Math. Soc. Trans. of Math. Monographs **11**, Providence, RI, 1964.)
- [10] **Tenenbaum G.**, *Introduction to analytic and probabilistic number theory*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 1995.
- [11] **Kátai I. and Subbarao M.V.**, On the local distribution of the iterated divisor function, *Mathematica Pannonica*, **15** (1) (2004), 127-140.

(Received September 10, 2010)

I. Kátai

Department of Computer Algebra
Eötvös Loránd University
Pázmány Péter sét. 1/C
H-1117 Budapest, Hungary
`katai@compalg.inf.elte.hu`