

RESULTS ON CLASSES OF FUNCTIONAL EQUATIONS

TRIBUTE TO ANTAL JÁRAI

by János Aczél and Che Tat Ng

While *individual* noncomposite functional equations in several variables had been solved at least since d'Alembert 1747 [9] and Cauchy 1821 [8], results on broad *classes* of such equations began appearing in the 1950's and 1960's. On general *methods of solution* see e.g. Aczél [1] and for *uniqueness of solutions* Aczél [2, 3], Aczél and Hosszú [6], Miller [20], Ng [21, 22], followed by several others. – Opening up and cultivating the field of *regularization* is *mainly Jári's achievement*. By regularization we mean assuming weaker regularity conditions, say measurability, of the unknown function and proving differentiability of several orders, for whole classes of functional equations. Differentiability of the unknown function(s) in the functional equation often leads to differential equations that are easier to solve.

For example, in Aczél and Chung [5] it was shown that locally Lebesgue integrable solutions of the functional equation

$$\sum_{i=1}^n f_i(x + \lambda_i y) = \sum_{k=1}^m p_k(x) q_k(y)$$

holding for x, y on open real intervals, with appropriate independence between the functions, are in fact differentiable infinitely many times. The differentiable solutions are then extracted using induced differential equations. Jári [11] showed that Lebesgue measurability and ordinary linear independence are sufficient to lead to the same solutions.

Aczél [4] called attention to some unsolved problems in the area of functional equations. One concerned Hilbert's fifth problem. Jári [15] formulated a problem that falls within that general call for non-composite functional equations in multiple variables. Here we exhibit the intricate problem he formulated and the sequence of results that led to its solution, and make references to his comprehensive book Jári [16].

Problem. Let T and Z be open subsets of \mathbb{R}^s and \mathbb{R}^m , respectively, and let D be an open subset of $T \times T$. Let $f : T \rightarrow Z$, $g_i : D \rightarrow T$ ($i = 1, \dots, n$), and $h : D \times Z^n \rightarrow Z$ be functions. Suppose that

$$f(t) = h(t, y, f(g_1(t, y)), \dots, f(g_n(t, y))) \quad \text{for all } (t, y) \in D ;$$

h is analytic;

g_1, \dots, g_n are analytic and for each $t \in T$ there exists a y for which

$$(t, y) \in D \quad \text{and} \quad \frac{\partial g_i}{\partial y} \quad \text{has rank } s \quad \text{for each } i = 1, \dots, n.$$

Is it true that every solution f which is measurable, or has the Baire property, is also analytic?

He proposed some incremental steps which may be taken to address the problem:

- (I) Measurability implies continuity.
- (II) Almost open solutions are continuous.
- (III) Continuous solutions are locally Lipschitz.
- (IV) Locally Lipschitz solutions are continuously differentiable.
- (V) All p -times continuously differentiable solutions are $(p + 1)$ -times differentiable.
- (VI) Infinitely many times differentiable solutions are analytic.

In [19] Járai and Székelyhidi outlined the above steps and gave a survey on the advances made. Many historic attributions were made to contributors in the field. Ng [23] contains results concerning the functional equation

$$f(x) + g(y) = h(T(x, y))$$

with given T . It is shown that under suitable assumptions, local boundedness of f implies the continuity of g .

Járai published a sequence of papers obtaining impressive results about that problem. [12] contains results regarding (I), (II), (IV), (V), and partially about (III). Step (III) is obtained for one variable in [13] and is generalized in [14]. In [15] Járai obtained the following result on the problem formulated above.

Theorem. *Suppose that the conditions of the Problem are satisfied and suppose that f has locally essentially bounded variation. Then f is infinitely many times differentiable.*

[16] contains, in Section 1, a summary account about the problem. We include some of it (abbreviated).

Theorem. (i) *If h is continuous and the functions g_i are continuously differentiable then every solution f , which is Lebesgue measurable or has Baire property, is continuous.*

(ii) *If h and g_i are p times continuously differentiable, then every almost everywhere differentiable solution f is p times continuously differentiable.*

(iii) If h and g_i are $\max\{2, p\}$ times differentiable and there exists a compact subset C of T such that for each $t \in T$ there exists a $y \in T$ satisfying $g_i(t, y) \in C$, besides the other stated rank condition on g_i , then every solution f , which is Lebesgue measurable or has the Baire property, is p times continuously differentiable ($1 \leq p \leq \infty$; $i = 1, \dots, n$).

Járai has deep insights and knowledge in the field of real analysis. He used the theorems reported in Giusti [10] swiftly, made fine and technical adaptations when necessary to get the above strong results.

In his book [16] many regularization theorems by him and others are assembled in a well organized way. For the convenience of the readers he has given several examples to illustrate how his general results can be applied to known functional equations. He devised and proved a general transfer principle which makes it possible to apply theorems concerning problems having only one unknown function also for cases with several unknown functions. A good example amongst many is the following

Theorem. Let $\alpha \neq \beta$ be fixed real numbers, $f, g_1, g_2 :]0, 1[\rightarrow \mathbb{R}$. Suppose that the functional equation

$$\begin{aligned} & f(x) + (1-x)^\alpha g_1(u/(1-x)) + (1-x)^\beta g_2(u/(1-x)) \\ &= f(u) + (1-u)^\alpha g_1(x/(1-u)) + (1-u)^\beta g_2(x/(1-u)) \end{aligned}$$

is satisfied for all $x, u \in]0, 1[$ with $x + u \in]0, 1[$. If the functions f, g_1, g_2 are Lebesgue measurable then they are \mathcal{C}^∞ .

He offered readers some details which precede the applications of his regularization theorems. The functional equation has its source in the study of symmetric divergences and distance measures and the differentiable solutions have been reported by Sander [25]. A more elaborate example is their joint work in [18] connected to the Weierstrass sigma function (as in [7]). They extended the results of M. Bonk [7] on the functional equation

$$\chi(u+v)\phi(u-v) = \sum_{\nu=1}^k f_\nu(u)g_\nu(v)$$

and treated it under weaker regularity assumptions.

Section 16 of the book contains results on (VI), analyticity. Járai's results as well as those of Páles [24] are covered. In Járai, Ng and Zhang [17] a composite type functional equation is solved under different regularity assumptions. The uniqueness theorem of Ng [22] is applied to obtain continuous solutions in one case, and the differentiation steps are used to extract the differentiable solutions in another case.

Acknowledgement. We thank the referee for helpful comments.

References

- [1] **Aczél, J.**, Some general methods in the theory of functional equations in one variable. New applications of functional equations (Russian), *Uspekhi Mat. Nauk (N.S.)*, **11** (1956), no. 3(69), 3–68.
- [2] **Aczél, J.**, Ein Eindeutigkeitssatz in der Theorie der Funktionalgleichungen und einige seiner Anwendungen, *Acta Math. Acad. Sci. Hungar.*, **15** (1964), 355–362.
- [3] **Aczél, J.**, On strict monotonicity of continuous solutions of certain types of functional equations, *Canad. Math. Bull.*, **9** (1966), 229–232.
- [4] **Aczél, J.**, Some unsolved problems in the theory of functional equations II., *Aequationes Math.*, **26** (1984), 255–260.
- [5] **Aczél, J. and J.K. Chung**, Integrable solutions of functional equations of a general type, *Studia Sci. Math. Hungar.*, **17** (1982), 51–87.
- [6] **Aczél, J. and M. Hosszú**, Further uniqueness theorems for functional equations, *Acta Math. Acad. Sci. Hungar.*, **16** (1965), 51–55.
- [7] **Bonk, M.**, 1994, The addition theorem of Weierstrass’s sigma function, *Math. Ann.*, **298** (1994), 591–610.
- [8] **Cauchy, A.L.**, *Cours d’analyse de l’École Polytechnique, I.*, Oeuvres, Sér. 2, vol. 3, Paris, 1867.
- [9] **d’Alembert, J.**, Recherches sur la courbe que forme une corde tendue mise en vibration, I, II., *Hist. Acad. Berlin*, 1747, 214–249.
- [10] **Giusti, E.**, *Minimal Surfaces and Functions of Bounded Variation*, Birkhäuser Verlag, Boston–Basel–Stuttgart, 1984.
- [11] **Járai, A.**, A remark to a paper of J. Aczél and J.K. Chung, *Studia Sci. Math. Hungar.*, **19** (1984), 273–274.
- [12] **Járai, A.**, On regular solutions of functional equations, *Aequationes Math.*, **30** (1986), 21–54.
- [13] **Járai, A.**, On continuous solutions of functional equations, *Publ. Math. Debrecen.*, **44** (1994), 115–122.
- [14] **Járai, A.**, On Lipschitz property of solutions of functional equations, *Aequationes Math.*, **47** (1994), 69–78.
- [15] **Járai, A.**, Solutions of functional equations having bounded variation, *Aequationes Math.*, **61** (2001), 205–211.
- [16] **Járai, A.**, *Regularity Properties of Functional Equations in Several Variables*, Springer, New York, 2005.
- [17] **Járai, A., C.T. Ng and W. Zhang**, A functional equation involving three means, *Rocznik Nauk.-Dydakt. Prace Mat.*, **17** (2000), 117–123.

- [18] **Járai, A. and W. Sander**, On the characterization of Weierstrass's sigma function, In: *Functional Equations – Results and Advances* (eds.: Z. Daróczy and Z. Páles), Kluwer, Dordrecht-Boston-London, 2002, pp. 29–79.
- [19] **Járai, A. and L. Székelyhidi**, Regularization and general methods in the theory of functional equations, *Aequationes Math.*, **52** (1996), 10–29.
- [20] **Miller, J.B.**, Aczél's uniqueness theorem and cellular internity, *Aequationes Math.*, **5** (1970), 319–325.
- [21] **Ng, C.T.**, Uniqueness theorems for a general class of functional equations, *J. Austral. Math. Soc.*, **11** (1970), 362–366.
- [22] **Ng, C.T.**, On the functional equation $f(x) + \sum_{i=1}^n g_i(y_i) = h(T(x, y_1, y_2, \dots, y_n))$, *Ann. Polon. Math.*, **27** (1973), 329–336.
- [23] **Ng, C.T.**, Local boundedness and continuity for a functional equation on topological spaces, *Proc. Amer. Math. Soc.*, **39** (1973), 525–529.
- [24] **Páles, Zs.**, On reduction of linear two variable functional equations to differential equations, *Aequationes Math.*, **43** (1992), 236–247.
- [25] **Sander, W.**, A generalization of a theorem of S. Picard, *Proc. Amer. Math. Soc.*, **73** (1979), 281–282.

J. Aczél and C.T. Ng

Department of Pure Mathematics

University of Waterloo

Waterloo, Ontario

Canada N2L 3G1

jdaczel@uwaterloo.ca

ctng@uwaterloo.ca

