A NOTE ON DYADIC HARDY SPACES

P. Simon (Budapest, Hungary)

Dedicated to the 60th birthday of Professor Antal Járai

Abstract. The usual L^p -norms are trivially invariant with respect to multiplication by *Walsh* functions. The analogous question will be investigated in the dyadic Hardy space **H**. We introduce an invariant subspace **H**_{*} of **H** in this sense and show some properties of **H**_{*}. For example a function in **H**_{*} will be constructed the *Walsh–Fourier* series of which diverges in L^1 -norm.

1. Introduction

Let $w_n \ (n \in \mathbf{N})$ be the Walsh-Paley system defined on the interval [0, 1). It is well-known that $w_n = \prod_{k=0}^{\infty} r_k^{n_k}$, where r_k is the k-th Rademacher function $(k \in \mathbf{N})$ and $n = \sum_{k=0}^{\infty} n_k 2^k \ (n_k = 0 \text{ or } 1 \text{ for all } k$'s) is the dyadic representation of n. If $n = \sum_{k=0}^{\infty} n_k 2^k$, $m = \sum_{k=0}^{\infty} m_k 2^k \in \mathbf{N}$ then $w_n w_m = w_{n \oplus m}$, where the operation \oplus is defined by

$$n \oplus m := \sum_{k=0}^{\infty} |n_k - m_k| 2^k.$$

Thus it is clear that

 $2^n \oplus m = 2^n + m \quad (n \in \mathbf{N}, \ m = 0, \dots, 2^n - 1),$

The Project is supported by the European Union and co-financed by the European Social Fund (grant agreement no. TAMOP 4.2.1/B-09/1/KMR-2010-0003). https://doi.org/10.71352/ac.35.255

i.e. $r_n w_m = w_{2^n} w_m = w_{2^n+m}$. (For more details we refer to the book [1].) For $1 \leq p \leq \infty$ let $L^p := L^p[0,1)$ and let $\|.\|_p$ denote the usual *Lebesgue* space and norm. If $f \in L^1$, $n \in \mathbb{N}$ then let $S_n f$ be the *n*-th *Walsh–Fourier* partial sum of f, i.e. $S_n f = f * D_n$, where $D_n := \sum_{k=0}^{n-1} w_k$ and * stands for dyadic convolution. We remark that $r_n D_{2^n} = D_{2^{n+1}} - D_{2^n}$ $(n \in \mathbb{N})$. The next famous property of D_{2^n} 's plays an important role in the Walsh analysis:

(1)
$$D_{2^n}(x) = \begin{cases} 2^n & (0 \le x < 2^{-n}) \\ 0 & (2^{-n} \le x < 1). \end{cases}$$

Therefore

$$S_{2^n}f(x) = 2^n \int_{I_n(x)} f$$
 $(x \in [0,1))$

Here $x \in I_n(x) := [j2^{-n}, (j+1)2^{-n})$ with a proper integer $j(x) = j = 0, \ldots, 2^n - 1$. Set $I_n := I_n(0)$.

We recall that

(2)
$$\sup_{n} \frac{\|D_n\|_1}{\log n} < \infty.$$

The dyadic maximal function f^* of $f \in L^1$ is defined as follows:

$$f^* := \sup_n |S_{2^n}f|.$$

Then for all p > 1 we have $||f||_p \le ||f^*||_p \le C_p ||f||_p$. (Here and later C_p, C will denote positive constants depending at most on p, although not always the same in different occurrences.) The so-called dyadic *Hardy* space $\mathbf{H} := \mathbf{H}[0, 1)$ is defined by means of the maximal function as follows:

$$\mathbf{H} := \{ f \in L^1 : \|f\| := \|f^*\|_1 < \infty \}.$$

The atomic structure of **H** is very useful in many investigations. Namely, we call a function $a \in L^{\infty}$ (dyadic) atom if $\int_{0}^{1} a = 0$ and there exists a dyadic interval $I_{n}(z)$ $(n \in \mathbf{N}, z \in [0, 1))$ such that a(x) = 0 $(x \in [0, 1) \setminus I_{n}(z))$ and $||a||_{\infty} \leq 2^{n}$. Let supp $a := I_{n}(z)$. The characterization of **H** by means of atoms reads as follows:

$$f \in \mathbf{H} \iff f = \sum_{k=0}^{\infty} \alpha_k a_k,$$

where all a_k 's are atoms and the coefficients α_k 's have the next property: $\sum_{k=0}^{\infty} |\alpha_k| < \infty$. Furthermore,

$$||f|| \sim \inf \sum_{k=0}^{\infty} |\alpha_k|,$$

where the infimum is taken over all atomic representations $\sum_{k=0}^{\infty} \alpha_k a_k$ of f. (For the martingale theoretic background we refer to [4].)

For example the functions $r_n D_{2^n}$ $(n \in \mathbf{N})$ are trivially atoms by (1). Thus

(3)
$$f := \sum_{n=0}^{\infty} \alpha_n r_{\nu_n} D_{2^{\nu_n}}$$

belongs to **H** if $\sum_{k=0}^{\infty} |\alpha_n| < \infty$ and the indices $\nu_0 < \nu_1 < \ldots$ are choosen arbitrarily. Moreover, $||f||_1 \le ||f|| \le \sum_{n=0}^{\infty} |\alpha_n|$.

It is not hard to see that the partial sums $S_{2^n}a$ $(n \in \mathbf{N})$ remain atoms if $a \in L^{\infty}$ is an atom. Indeed, if supp $a = I_N(z)$ $(N \in \mathbf{N}, z \in [0, 1))$ and $x \in [0, 1) \setminus I_N(z)$ then for all $n \in \mathbf{N}$ the intervals $I_n(x)$ and $I_N(z)$ are disjoint or $I_n(x) \cap I_N(z) = I_N(z)$. Thus

$$|S_{2^n}a(x)| = \left|2^n \int_{I_n(x)} a\right| = \left|2^n \int_{I_n(x)\cap I_N(z)} a\right| \le \left|2^n \int_{I_N(z)} a\right| = \left|2^n \int_0^1 a\right| = 0,$$

thus $S_{2^n}a(x) = 0$. Furthermore, $||S_{2^n}a||_{\infty} \leq ||a||_{\infty} \leq 2^N$, i.e. supp $S_{2^n}a = I_N(z)$ and $\int_0^1 S_{2^n}a = \int_0^1 a = 0$.

Therefore if $f = \sum_{k=0}^{\infty} \alpha_k a_k$ is an atomic representation of $f \in \mathbf{H}$ then $S_{2^n}f = \sum_{k=0}^{\infty} \alpha_k S_{2^n} a$ $(n \in \mathbf{N})$ is an atomic representation of $S_{2^n}f$. This means that $\|S_{2^n}f\| \leq \sum_{k=0}^{\infty} |\alpha_k|$, i.e. $\|S_{2^n}f\| \leq \|f\|$. (The last inequality follows also from the obvious estimation $(S_{2^n}f)^* \leq f^*$.)

We remark that **H** can be defined also in another way. To this end let $f \in L^1$ and

$$Qf := \left(\sum_{n=-1}^{\infty} (\delta_n f)^2\right)^{1/2}$$

be its quadratic variation, where $\delta_{-1}f := \int_0^1 f$, $\delta_n f := S_{2^{n+1}}f - S_{2^n}f = f * (r_n D_{2^n})$ $(n \in \mathbf{N})$. Then

$$||f|| \sim ||Qf||_1$$
, ill. $||f||_p \sim ||Qf||_p$ $(1$

If $f \in L^1$, $n \in \mathbf{N}$ and $k = 0, \ldots, 2^n - 1$, then w_k is constant on $I_n(x)$ ($x \in (0,1)$), consequently $w_k(x) \int_{I_n(x)} f = \int_{I_n(x)} (fw_k)$. This means that $w_k S_{2^n} f = S_{2^n}(fw_k)$. Furthermore, if $2^n \leq k \in \mathbf{N}$ is arbitrary then let us write $k = \sum_{j=0}^N k_j 2^j$ (with some $\mathbf{N} \ni N \ge n$). It is clear that

$$\delta_j(w_k S_{2^n} f) = \begin{cases} 0 & (j \neq N) \\ w_k S_{2^n} f & (j = N) \end{cases} \quad (j \in \mathbf{N}).$$

From this it follows that $Q(w_k S_{2^n} f) = |S_{2^n} f|$, i.e. for all $k \in \mathbf{N}$ we have

(4)
$$\|w_k S_{2^n} f\| = \|S_{2^n}(fw_k)\| \quad (k < 2^n) \quad \text{and} \\ \|w_k S_{2^n} f\| \le C \|S_{2^n} f\|_1 \quad (k \ge 2^n).$$

The Walsh–Paley system doesn't form a basis in L^1 . Moreover, there exists $f \in \mathbf{H}$ such that

$$\sup_{n} \|S_n f\|_1 = \infty.$$

However (see [3]), if $f \in \mathbf{H}$ then

$$\frac{1}{\log n} \sum_{k=1}^{n} \frac{\|S_k f\|_1}{k} \to \|f\| \qquad (n \to \infty),$$

or equivalently

$$\frac{1}{\log n} \sum_{k=1}^{n} \frac{\|f - S_k f\|_1}{k} \to 0 \qquad (n \to \infty).$$

For the sake of the completeness and in order to demonstrate the usefulness of the atomic structure we sketch some examples. Namely we take the function given by (3). If $l_n = 0, 1, \ldots, 2^{\nu_n} - 1$ $(n \in \mathbf{N})$ then

(*)
$$\|S_{2^{\nu_n}+l_n}f - S_{2^{\nu_n}}f\|_1 = |\alpha_n| \|D_{l_n}\|_1$$

It is well-known that $k_n \in \{0, 1, \dots, 2^{\nu_n} - 1\}$ can be choosen so that

$$||D_{k_n}||_1 \ge C\nu_n \qquad (n \in \mathbf{N})$$

holds. Then we get

$$||S_{2^{\nu_n}+k_n}f - S_{2^{\nu_n}}f||_1 \ge C|\alpha_n|\nu_n \qquad (n \in \mathbf{N}).$$

If $\sup_n |\alpha_n|\nu_n = \infty$ then $||S_{2^n}f||_1 \leq \sum_{k=0}^{\infty} |\alpha_k| < \infty$ implies $\sup_n ||S_nf||_1 = \infty$. It is obvious that $\alpha_n := 2^{-n}, \nu_n := 2^{n^2}$ $(n \in \mathbf{N})$ are suitable. (We remark that $\inf_n |\alpha_n|\nu_n > 0$ is trivially sufficient for the $||.||_1$ divergence of the Walsh-Fourier series of f.)

If $f \in \mathbf{H}$ is given by (3) then $||S_n f - f||_1 \to 0$ $(n \to \infty)$ if and only if $\nu_n \alpha_n \to 0$ $(n \to \infty)$. Indeed, if $l_n := k_n$'s are as above then $C\nu_n |\alpha_n| \leq |\alpha_n| ||D_{k_n}||_1$ and (*) proves necessity. It is known that $||S_{2^n}g - g||_1 \to 0$ $(n \to \infty)$ for all $g \in L^1$. Therefore (see (2)) $||D_{l_n}||_1 \leq C \log l_n \leq C\nu_n$ and $\nu_n \alpha_n \to 0$ $(n \to \infty)$ together with (*) imply the $||.||_1$ convergence of the series (3). Finally, we cite an example $f \in L^1 \setminus H$ such that $||S_n f - f||_1 \to 0 \quad (n \to \infty)$. To this end we take a special function $f := \sum_{n=0}^{\infty} \alpha_n r_n D_{2^n}$ in (3) such that the coefficients α_n form a null-sequence of bounded variation, i.e. $\sum_{n=0}^{\infty} |\alpha_n - \alpha_{n+1}| < \infty$. It is well-known that this assumption on the coefficients implies the $||.||_1$ -convergence of the series in question. Indeed, for all $n, m \in \mathbb{N}$, n < mit follows by (1) that

$$\left\|\sum_{k=n}^{m} \alpha_{k} r_{k} D_{2^{k}}\right\|_{1} = \left\|\sum_{k=n}^{m} \alpha_{k} (D_{2^{k+1}} - D_{2^{k}})\right\|_{1} = \\ = \left\|\sum_{k=n+1}^{m} (\alpha_{k-1} - \alpha_{k}) D_{2^{k}} + \alpha_{m} D_{2^{m}} - \alpha_{n} D_{2^{n}}\right\|_{1} \leq \\ \leq \sum_{k=n+1}^{m} |\alpha_{k-1} - \alpha_{k}| \|D_{2^{k}}\|_{1} + |\alpha_{m}| \|D_{2^{m}}\|_{1} + |\alpha_{n}| \|D_{2^{n}}\|_{1} = \\ = \sum_{k=n+1}^{m} |\alpha_{k-1} - \alpha_{k}| + |\alpha_{m}| + |\alpha_{n}| \rightarrow 0 \quad (n, m \rightarrow \infty).$$

Therefore $f \in L^1$. Furthermore, if $2^{-k-1} \le x < 2^{-k}$ $(k \in \mathbf{N})$ then

$$Qf(x) = \sqrt{\sum_{n=0}^{\infty} \alpha_n^2 D_{2^n}^2(x)} = \sqrt{\sum_{n=0}^k \alpha_n^2 2^{2n}} \ge |\alpha_k| 2^k,$$

and

$$\|Qf\|_1 \ge \sum_{k=0}^{\infty} \int_{2^{-k-1}}^{2^{-k}} Qf \ge \sum_{k=0}^{\infty} \int_{2^{-k-1}}^{2^{-k}} |\alpha_k| 2^k = \frac{1}{2} \sum_{k=0}^{\infty} |\alpha_k|.$$

This means that $||f|| = \infty$ if $\sum_{k=0}^{\infty} |\alpha_k| = \infty$. Now, we prove the $||.||_1$ convergence of the sequence $S_n f$. To this end let $1 \le n \in \mathbb{N}$ and $m_n = 0, ..., 2^n - 1$. Then by (2) we have

$$||S_{2^n + m_n} f - S_{2^n} f||_1 = ||\alpha_n r_n D_{m_n}||_1 = |\alpha_n| ||D_{m_n}||_1 \le C |\alpha_n| \log m_n \le C n |\alpha_n|.$$

Hence $n\alpha_n \to 0$ $(n \to \infty)$ is implies to $||S_{2^n+m_n}f - S_{2^n}f||_1 \to 0$ $(n \to \infty)$. Since $||S_{2^n}f - f||_1 \to 0$ $(n \to \infty)$ we get $||S_nf - f||_1 \to 0$ $(n \to \infty)$. A simple calculation shows that the sequence

$$\alpha_n := \frac{1}{(n+2)\log(n+2)} \qquad (n \in \mathbf{N})$$

satisfies all of the conditions above. By means of similar observations it can be proved that the assumption $\sum_{n=0}^{\infty} |\alpha_n| < \infty$ in (3) is necessary to $f \in \mathbf{H}$ in the general case as well.

2. Results

It is clear that for all $f \in L^p$ $(1 \le p \le \infty)$ and $n \in \mathbf{N}$ we have $fw_n \in L^p$ and $\|fw_n\|_p = \|f\|_p$. The situation in the case of **H** is more complicated. For example if we take the atoms $f_n := r_n D_{2^n} \in \mathbf{H}$ $(n \in \mathbf{N})$ then $\|f_n\| = 1$ and

$$||r_n f_n|| = ||D_{2^n}|| = ||D_{2^n}^*||_1 = \left||\max_{k \le n} D_{2^k}||_1\right|$$

where by (1)

$$\max_{k \le n} D_{2^k}(x) = \begin{cases} 2^k & (2^{-k-1} \le x < 2^{-k}, \ k = 0, \dots, n-1) \\ 2^n & (0 \le x < 2^{-n}). \end{cases}$$

From this it follows immediately that $||D_{2^n}|| = \frac{n+2}{2}$, i.e.

$$||r_n f_n|| = ||w_{2^n} f_n|| = \frac{n+2}{2} ||f_n||.$$

First we prove that an analogous relation holds in general.

Theorem 1. Let $k \in \mathbf{N}$. Then there exists a constant C_k such that for all $f \in \mathbf{H}$ the product fw_k belongs to \mathbf{H} and $||fw_k|| \leq C_k ||f||$.

Our example above shows that $C_{2^n} \geq \frac{n+2}{2}$ $(n \in \mathbf{N})$, i.e. $\sup_k C_k = \infty$. Since all *Walsh* functions are final products of *Rademacher* functions, we need to prove Theorem 1 only for $k = 2^n$ $(n \in \mathbf{N})$.

In this case let $f = \sum_{k=0}^{\infty} \alpha_k a_k$ be an atomic representation of $f \in \mathbf{H}$. Then

$$\|fw_{2^{n}}\| = \|fr_{n}\| = \|(fr_{n})^{*}\|_{1} \le \left\|\sum_{k=0}^{\infty} |\alpha_{k}|(a_{k}r_{n})^{*}\right\|_{1} \le \\ \le \sum_{k=0}^{\infty} |\alpha_{k}|\|(a_{k}r_{n})^{*}\|_{1} = \sum_{k=0}^{\infty} |\alpha_{k}|\|a_{k}r_{n}\|.$$

If we can show that

$$A_n := \sup_a \|ar_n\| < \infty$$

(where the supremum is taken over all atoms a), then

$$||(fr_n)^*||_1 \le A_n \sum_{k=0}^{\infty} |\alpha_k|,$$

i.e. $||fr_n|| \le A_n ||f||$.

Proof of the inequality (**). Let a be an atom, $k \in \mathbb{N}, x \in [0, 1)$. In the case k > n the *n*-th *Rademacher* function r_n is constant on the interval $I_k(x)$ and thus

$$S_{2^{k}}(ar_{n})(x) = 2^{k} \int_{I_{k}(x)} ar_{n} = 2^{k} r_{n}(x) \int_{I_{k}(x)} a$$

Therefore

$$(ar_n)^* = \sup_k |S_{2^k}(ar_n)| \le \max_{k \le n} |S_{2^k}(ar_n)| + \sup_{k > n} |S_{2^k}a| \le \le \max_{k \le n} |S_{2^k}(ar_n)| + \sup_k |S_{2^k}a| = \max_{k \le n} |S_{2^k}(ar_n)| + a^* =: (ar_n)^{**} + a^*.$$

From this it follows that

$$\begin{aligned} \|ar_n\| &= \|(ar_n)^*\|_1 \le \|(ar_n)^{**}\|_1 + \|a^*\|_1 = \\ &= \|(ar_n)^{**}\|_1 + \|a\| \le \|(ar_n)^{**}\|_1 + 1. \end{aligned}$$

This means that it is enough to show only

$$\sup_{a} \|(ar_n)^{**}\|_1 < \infty$$

(where the supremum is taken over all atoms a).

To this end let a be an atom. For the sake of simplicity we assume that supp $a = I_N$ (with some $N \in \mathbf{N}$). Then

$$||(ar_n)^{**}||_1 = \int_{I_N} (ar_n)^{**} + \int_{2^{-N}}^1 (ar_n)^{**} =: J_1(a) + J_2(a).$$

Hence by means of the *Cauchy* inequality and the properties of atoms it follows that

$$J_1(a) \le \left(\int_{I_N} ((ar_n)^{**})^2 \right)^{1/2} \cdot 2^{-N/2} \le 2^{-N/2} ||(ar_n)^{**}||_2 \le C_2 2^{-N/2} ||ar_n||_2 \le C_2 2^{-N/2} ||a||_{\infty} 2^{-N/2} \le C_2.$$

We will show that

$$\sup_{a} J_2(a) < \infty.$$

Indeed, if a is the atom as above and n < N, then $ar_n = a$, i.e.

$$J_2(a) \le \|(ar_n)^{**}\|_1 = \|\max_{k \le n} |S_{2^k}a|\|_1 \le \|a^*\|_1 = \|a\| \le 1.$$

Thus it can be assumed that $N \leq n$. Let k = 0, ..., n and $2^{-N} \leq x < 1$. Then

$$S_{2^k}(ar_n)(x) = 2^k \int_{I_k(x)} ar_n = 2^k \int_{I_k(x)\cap I_N} ar_n$$

where $I_k(x) \cap I_N \neq \emptyset$ exactly if $k \leq N-1$ and $x < 2^{-k}$ (in this case $I_k(x) = I_k$ and $I_k(x) \cap I_N = I_N$). This means that with the notation $k_0(x) := \max\{k = 0, ..., N-1 : x < 2^{-k}\}$ we get

$$(ar_n)^{**}(x) = \max_{k \le k_0(x)} |S_{2^k}(ar_n)(x)| = \max_{k \le k_0(x)} 2^k \left| \int_{I_N} ar_n \right| \le \\ \le \max_{k \le k_0(x)} 2^k ||a||_1 \le 2^{k_0(x)} \le \frac{1}{x}.$$

Summarizing the above facts it follows that

$$J_2(a) = \int_{2^{-N}}^{1} (ar_n)^{**} \le \int_{2^{-N}}^{1} \frac{dx}{x} \le C \log_2 2^N = CN \le Cn,$$

which proves Theorem 1. \blacksquare

Therefore it can be assumed that $\frac{n+2}{2} \leq C_{2^n} \leq C(n+1)$ $(n \in \mathbf{N})$. Furthermore, if $n = \sum_{j=0}^{\infty} n_j 2^j$ is the dyadic representation of $n \in \mathbf{N}$, then

$$||fw_n|| \le ||f|| \prod_{j=0}^{\infty} C_{2^j}^{n_j} \le C^{|n|}[n]||f|| \qquad (f \in \mathbf{H}),$$

where $|n| := \sum_{j=0}^{\infty} n_j$, and $[n] := \prod_{j=0}^{\infty} (j+1)^{n_j}$, and the above estimation cannot be improved. For example $|2^k| = 1$ and $[2^k] = k+1$ $(k \in \mathbf{N})$.

Theorem 1 involves the next concept: if $f \in \mathbf{H}$ then let

$$||f||_* := \sup_n ||fw_n||.$$

It follows immediately that $\|.\|_*$ is a norm, $\|.\| \leq \|.\|_*$ but (see the above remarks) $\|.\|_*, \|.\|$ are not equivalent. Moreover, it is not hard to construct $f \in \mathbf{H}$ such that $\|f\|_* = \infty$. Indeed, we take the function given in (3). Then for all $k \in \mathbf{N}$ we get

$$\|fr_{\nu_k}\| \ge |\alpha_k| \|D_{2^{\nu_k}}\| - \left\|\sum_{k \ne n=0}^{\infty} \alpha_n r_{\nu_k} r_{\nu_n} D_{2^{\nu_n}}\right\|.$$

It is clear that all products $r_{\nu_k}r_{\nu_n}D_{2^{\nu_n}}$ $(k \neq n \in \mathbf{N})$ are atoms, which implies

$$\left\|\sum_{k\neq n=0}^{\infty} \alpha_n r_{\nu_k} r_{\nu_n} D_{2^{\nu_n}}\right\| \le \sum_{n=0}^{\infty} |\alpha_n| = q < \infty.$$

Then

$$||f||_* \ge ||fr_{\nu_k}|| \ge |\alpha_k| ||D_{2^{\nu_k}}|| - q = |\alpha_k| \frac{\nu_k + 2}{2} - q \to \infty \qquad (k \to \infty)$$

follows by means of a suitable choice of parameters.

F. Schipp (see [2]) introduced the following norms

$$\|f\|_{*p} := \|\sup_{n} Q(fw_{n})\|_{p} , \|f\|^{*p} := \left\|\sup_{m,n} |S_{2^{m}}(fw_{n})|\right\|_{p}$$
$$(f \in L^{1}, \ 1 \le p < \infty),$$

and proved the non-trivial equivalence $||f||_{*p} \sim ||f||_p$ (1 . It is clear $that these norms are shift invariant, i.e. for all <math>n \in \mathbb{N}$ the equalities $||fw_n||_{*p} =$ $= ||f||_{*p}, ||fw_n||^{*p} = ||f||^{*p}$ hold. Furthermore, the inequality $||.||_* \leq ||.||^{*1}$ follows immediately. Moreover, for all $k \in \mathbb{N}$ we get

$$||fw_k|| \le C ||Q(fw_k)||_1 \le C ||\sup_n Q(fw_n)||_1 = C ||f||_{*1},$$

i.e. $||f||_* \leq C||f||_{*1}$ holds, too. Schipp proved for $F := \sum_{n=0}^{\infty} 2^{-n/2} r_{2^n} D_{2^{2^n}}$ that $F \in \mathbf{H}$ but $||F||_{*1} = \infty$. (This example is a special case of (3).) Our example above along with $||.|| \leq ||.||_* \leq ||.||^{*1}$ shows also the existence of $f \in \mathbf{H}$ such that $||f||^{*1} = \infty$. The question wheter the norm $||.||_{*1}$ and the norm $||.||^{*1}$ are equivalent or not remains open.

Let us introduce the space \mathbf{H}_* as follows:

$$\mathbf{H}_* := \{ f \in H : \|f\|_* < \infty \}.$$

Then \mathbf{H}_* is a proper subspace of \mathbf{H} . For all $n, k \in \mathbf{N}$ it is clear that $1 = ||w_n|| = ||w_k \oplus n|| = ||w_k w_n||$, i.e. $||w_n||_* = 1$. Thus $w_n \in \mathbf{H}_*$ and therefore every Walsh polynomial (finite linear combination of Walsh functions) belongs to \mathbf{H}_* . Furthermore, if $f \in \mathbf{H}_*$ then

$$||fw_n||_* = \sup_k ||fw_nw_k|| = \sup_k ||fw_{n\oplus k}|| = \sup_j ||fw_j|| = ||f||_*$$

In other words the norm $\|.\|_*$ is also invariant with respect to multiplication by *Walsh* functions.

Above we remarked that there exists $f \in \mathbf{H}$ such that its *Walsh–Fourier* series diverges in $\|.\|_1$ norm. We show that this result can be sharpened. Namely, the next theorem holds:

Theorem 2. There exists $f \in \mathbf{H}_*$ with $\|.\|_1$ -divergent Walsh-Fourier series.

Proof. We take the function $f := \sum_{n=0}^{\infty} \alpha_n r_{\nu_n} D_{2^{\nu_n}}$ from (3). It was shown above (see (*)) that $q := \sum_{n=0}^{\infty} |\alpha_n| < \infty$ and $\inf_n |\alpha_n|\nu_n > 0$ imply the $\|.\|_1$ divergence of the Walsh-Fourier series of f.

To the proof of $f \in H_*$ let $k = \sum_{j=0}^{\infty} k_j 2^j$ be the dyadic representation of $k \in \mathbf{N}$. Then $w_k = \prod_{j=0}^{\infty} r_j^{k_j}$. Taking into account that

$$w_k r_s D_{2^s} = \prod_{j=s}^{\infty} r_j^{k_j} r_s D_{2^s} \qquad (s \in \mathbf{N})$$

is obviously an atom, provided $k_s = 0$ or $k_s = 1$, but there is $j \ge s + 1$ such that $k_j = 1$. Let \mathbf{N}_s be the set of such k's. Then $k \in \mathbf{N}^s := \mathbf{N} \setminus \mathbf{N}_s$ iff $k = 2^s + \sum_{j=0}^{s-1} k_j 2^s$, i.e. $\mathbf{N}^s = \mathbf{N} \cap [2^s, 2^{s+1})$. In this case $w_k r_s D_{2^s} = D_{2^s}$.

If $k \notin \bigcup_{n=0}^{\infty} \mathbf{N}^{\nu_n}$, then

$$fw_k = \sum_{n=0}^{\infty} \alpha_n w_k r_{\nu_n} D_{2^{\nu_n}}$$

is an atomic representation of fw_k and so $||fw_k|| \le \sum_{n=0}^{\infty} |\alpha_n| = q$.

If $k \in \bigcup_{n=0}^{\infty} \mathbf{N}^{\nu_n}$, then there is a unique $m \in \mathbf{N}$ such that $k \in \mathbf{N}^{\nu_m}$:

$$fw_k = \alpha_m D_{2^{\nu_m}} + \sum_{m \neq n=0}^{\infty} \alpha_n w_k r_{\nu_n} D_{2^{\nu_n}} =: \alpha_m D_{2^{\nu_m}} + f_0.$$

The above observations lead to $||f_0|| \leq \sum_{n=0}^{\infty} |\alpha_n| = q < \infty$ and

$$||fw_k|| \le |\alpha_m| ||D_{2^{\nu_m}}|| + ||f_0|| \le C |\alpha_m|\nu_m + q.$$

We see that the assumption $\sup_n |\alpha_n| \nu_n < \infty$ is sufficient to

$$\sup_{k} \|fw_{k}\| \le C \sup_{n} |\alpha_{n}|\nu_{n} + q < \infty.$$

In this case $f \in H_*$. For example if $\alpha_n := 2^{-n}, \nu_n := 2^n$ $(n \in \mathbb{N})$, then the function $f = \sum_{n=0}^{\infty} 2^{-n} r_{2^{2^n}} D_{2^{2^n}}$ proves Theorem 2.

If $f \in \mathbf{H}$ then $Qf \in L^1$, i.e. $Qf = \left(\sum_{k=-1}^{\infty} (\delta_k f)^2\right)^{1/2} < \infty$ a.e. Thus $\left(\sum_{k=n}^{\infty} (\delta_k f)^2\right)^{1/2} \to 0 \quad (n \to \infty)$ a.e. and we get by *Lebesgue*'s theorem that

$$\|f - S_{2^n} f\| \le C \|Q(f - S_{2^n})\|_1 = C \left\| \left(\sum_{k=n}^{\infty} (\delta_k f)^2 \right)^{1/2} \right\|_1 \to 0 \qquad (n \to \infty).$$

However, this last convergence property doesn't hold true if the norm $\|.\|$ will be replaced by $\|.\|_*$. Indeed, taking the function $f \in \mathbf{H}_*$ from the proof of Theorem 2 we get analogously that

$$\|f - S_{2^{\nu_n}} f\|_* = \left\| \sum_{k=n}^{\infty} \alpha_k r_{\nu_k} D_{2^{\nu_k}} \right\|_* \ge C \inf_{k \ge n} |\alpha_k| \nu_k - q \qquad (n \in \mathbf{N}).$$

Let $\alpha_k := 2^{-k}, \nu_k := 2^{k+s}$ $(k \in \mathbf{N})$, where $s \in \mathbf{N}$ is defined by $2^s C > 2$. Then $q = \sum_{k=0}^{\infty} |\alpha_k| = 2$ and $||f - S_{2^{\nu_n}} f||_* \ge 2^s C - 2$ $(n \in \mathbf{N})$, i.e. $||f - S_{2^n} f||_*$ doesn't tend to zero if $n \to \infty$.

We recall that $||S_{2^n}f||_1 \leq ||f||_1$ $(f \in L^1)$, $||S_{2^n}f|| \leq ||f||$ $(f \in \mathbf{H}, n \in \mathbf{N})$. Applying (4) it is not hard to prove that an analogous inequality holds if we replace the norm ||.|| by $||.||_*$. Indeed,

$$||S_{2^{n}}f||_{*} = \sup_{k} ||w_{k}S_{2^{n}}f|| = \max\left\{\sup_{k<2^{n}} ||w_{k}S_{2^{n}}f||, \sup_{k\geq2^{n}} ||w_{k}S_{2^{n}}f||\right\} \le \\ \le \max\left\{\sup_{k<2^{n}} ||fw_{k}||, C||S_{2^{n}}f||_{1}\right\} \le \max\left\{\sup_{k} ||fw_{k}||, C||f||_{1}\right\} \le C||f||_{*}.$$

Hence if $f \in L^1$ then

$$||f||_* = \sup_n ||(fw_n)^*||_1 = \sup_n ||\sup_m |S_{2^m}(fw_n)||_1$$

Let p > 1 and $f \in L^p$. Then for arbitrary $n \in \mathbf{N}$ we can write

$$||fw_n|| = ||(fw_n)^*||_1 \le ||(fw_n)^*||_p \le C_p ||fw_n||_p = C_p ||f||_p,$$

i.e. $||f||_* \leq C_p ||f||_p$. Thus $L^p \subset H_*$. In other words $\bigcup_{p>1} L^p \subset \mathbf{H}_*$. We will show that the next statement holds:

Theorem 3. $H_* \setminus \left(\bigcup_{p>1} L^p\right) \neq \emptyset.$

Proof. Let $1 and take the function <math>f = \sum_{n=0}^{\infty} 2^{-n} r_{2^{2^n}} D_{2^{2^n}} =:$ =: $\sum_{n=0}^{\infty} \alpha_n r_{\nu_n} D_{2^{\nu_n}}$ as in the proof of Theorem 2. Then $f \in H_*$. On the other hand

$$\begin{split} \|f\|_{p}^{p} &\geq C_{p} \|Qf\|_{p}^{p} \geq C_{p} \left\| \sqrt{\sum_{n=0}^{\infty} \alpha_{n}^{2} D_{2^{\nu_{n}}}^{2}} \right\|_{p}^{p} \geq C_{p} \sum_{k=0}^{\infty} \int_{2^{-\nu_{k}}}^{2^{-\nu_{k}}} \left(\sum_{n=0}^{k} \alpha_{n}^{2} D_{2^{\nu_{n}}}^{2} \right)^{p/2} = \\ &= C_{p} \sum_{k=0}^{\infty} 2^{-\nu_{k}} \left(\sum_{n=0}^{k} \alpha_{n}^{2} 2^{2\nu_{n}} \right)^{p/2} \geq C_{p} \sum_{k=0}^{\infty} \alpha_{k}^{p} 2^{(p-1)\nu_{k}} = \infty. \blacksquare \end{split}$$

References

- Schipp, F., W.R. Wade, P. Simon and J. Pál, Walsh Series: An Introduction to Dyadic Harmonic Analysis, Akadémiai Kiadó, Budapest-Adam Hilger, Bristol-New York, 1990.
- [2] Schipp, F., On Paley-type inequality, Acta Sci. Math. (Szeged), 45 (1983), 357–364.
- [3] Simon, P., Strong convergence of certain means with respect to the Walsh–Fourier series, *Acta Math. Hungar.*, **49** (1987), 425–431.
- [4] Weisz, F., Martingale Hardy Spaces and their Applications in Fourier Analysis, Springer, Berlin-Heidelberg-New York, 1994.

P. Simon

Department of Numerical Analysis Eötvös Loránd University Pázmány P. sétány 1/C. H-1117 Budapest, Hungary simon@ludens.elte.hu