

ON SOME CLASSES OF PARTIAL DIFFERENCE EQUATIONS

Eszter Gselmann (Debrecen, Hungary)

*Dedicated to Professors Zoltán Daróczy and Imre Kátai
on the occasion of their 75th birthday*

Communicated by Antal Járai

(Received January 22, 2013; accepted April 09, 2013)

Abstract. In one of his work, appeared in 1969, John A. Baker initiated the systematic investigation of some partial difference equations. The main purpose of this paper is to continue and to extend these investigations. Firstly, we present how such type of equations can be classified into elliptic, parabolic and hyperbolic subclasses, respectively. After that, we show solution methods in the elliptic class. Here we will deal in details with the discrete version of the following partial differential equations: Laplace's equation, Poisson equation and the (in)homogeneous biharmonic equation.

1. Introduction

The aim of this paper is to give a systematic description of certain type of partial difference equations. Throughout this note \mathbb{N} , \mathbb{Z} , \mathbb{Q} and \mathbb{R} denote the sets of the natural, integer, rational and real numbers, respectively.

Key words and phrases: Partial difference equation; Laplace's equation, Poisson equation, biharmonic equation, mollification theorem, Harnack's principle.

2010 Mathematics Subject Classification: 39A14, 39B52.

This research has been supported by the Hungarian Scientific Research Fund (OTKA) Grant NK 81402 and by the TÁMOP 4.2.4.A/2-11-1-2012-0001 (Nemzeti Kiválóság Program - Hazai hallgatói, illetve kutatói személyi támogatást biztosító rendszer kidolgozása s működtetése konvergencia program) project implemented through the EU and Hungary co-financed by the European Social Fund.

<https://doi.org/10.71352/ac.40.285>

Let $\Omega \subset \mathbb{R}^n$ be a set and $u: \Omega \rightarrow \mathbb{R}$ be a function, then

$$\Delta_h u(x_1, \dots, x_n) = u(x_1, \dots, x_{i-1}, x_i + h, x_{i+1}, x_n) - u(x_1, \dots, x_n)$$

(x_i)

is called the *first order partial difference* of the function u with respect to the variable x_i . The higher order partial differences are defined recursively, that is, if $k \in \mathbb{N}$ and $k \geq 2$ and $i = 1, \dots, n$, then

$$\Delta_h^k u(\mathbf{x}) = \Delta_h \left(\Delta_h^{k-1} u(\mathbf{x}) \right)$$

$(x_i) \quad (x_i)$

assuming that $\mathbf{x} + k h e_i \in \Omega$ is valid for all $\mathbf{x} \in \Omega$, where e_i denotes the i^{th} standard base vector of \mathbb{R}^n . Further, if $k_1, \dots, k_n \in \mathbb{N} \cup \{0\}$ and $\sum_{i=1}^n k_i = N$, then

$$\Delta_h^N u(\mathbf{x}) = \Delta_h^{k_1} \dots \Delta_h^{k_n} u(\mathbf{x}).$$

$(x_1^{k_1}, \dots, x_n^{k_n}) \quad (x_1) \quad (x_n)$

In view of this notions on a *partial difference equation* we mean a functional equation that has the form

$$(1) \quad F \left(\Delta_h^N u(x_1, \dots, x_n), \dots, u(x_1, \dots, x_n), x_1, \dots, x_n \right) = 0,$$

$(x_1^{k_1}, \dots, x_n^{k_n})$

where F is a given function, u is the unknown function which has to be determined and $k_1, \dots, k_n \in \mathbb{N} \cup \{0\}$ are such that $\sum_{i=1}^n k_i = N$.

Equation (1) is called an N^{th} -order equation if it contains both $\Delta^N u$ and u . For the sake of simplicity, here we remark that other properties (e.g. linearity, quasilinearity, semilinearity etc.) of such type of equations can be defined analogously as in the theory of partial differential equations. Concerning partial difference equations we refer to the monograph of Cheng [2].

2. Classification of partial difference equations

Let $\Omega \subset \mathbb{R}^n$, $m \in \mathbb{N}$ and $\gamma_i: \Omega \rightarrow \mathbb{R}$ given functions and $\rho_{j,i} \in \mathbb{R}$, $j = 1, \dots, n$, $i = 1, \dots, m$ and let us consider the operator \mathcal{D} defined by

$$\mathcal{C}(\Omega) \ni u(x_1, \dots, x_n) \xrightarrow{\mathcal{D}} \sum_{i=1}^m \gamma_i(x_1, \dots, x_n) \cdot u(x_1 + \rho_{1,i} h, \dots, x_n + \rho_{n,i} h).$$

Here the domain Ω is assumed to be such that $\mathbf{x} + \rho_i h \in \Omega$ for all $i = 1, \dots, m$ and $\mathbf{x} = (x_1, \dots, x_n) \in \Omega$, where $\rho_i = (\rho_{1,i}, \dots, \rho_{n,i})$.

In this case let

$$A_{k,l}(\mathbf{x}) = A_{k,l}(x_1, \dots, x_n) \\ = \sum_{i=1}^m \rho_{k,i} \rho_{l,i} \gamma_i(x_1, \dots, x_n) \quad (k, l = 1, \dots, n, \mathbf{x} \in \Omega).$$

Further, let us consider the (symmetric) $n \times n$ matrix valued function Q defined by

$$Q(\mathbf{x}) = \begin{pmatrix} A_{11}(\mathbf{x}) & \dots & A_{1n}(\mathbf{x}) \\ \vdots & \ddots & \vdots \\ A_{n1}(\mathbf{x}) & \dots & A_{nn}(\mathbf{x}) \end{pmatrix} \quad (\mathbf{x} \in \Omega).$$

We say that the operator \mathcal{D} defined above is

- *elliptic* at the point $\mathbf{x} \in \Omega$, if $Q(\mathbf{x}) \in \mathcal{M}_{n \times n}(\mathbb{R})$ is a definite matrix;
- *hyperbolic* at the point $\mathbf{x} \in \Omega$, if $Q(\mathbf{x}) \in \mathcal{M}_{n \times n}(\mathbb{R})$ is an indefinite matrix;
- *parabolic* at the point $\mathbf{x} \in \Omega$, if $Q(\mathbf{x}) \in \mathcal{M}_{n \times n}(\mathbb{R})$ is a semidefinite matrix;

The systematic investigation of equations of the form

$$\mathcal{D}(u)(\mathbf{x}) + u(\mathbf{x}) = \Phi(x)$$

goes back to Baker [1], McKiernan [5], and Świątak [8]. The aim of this paper is to continue and to extend the results of John A. Baker. Additionally, not only the solutions will be derived but also we will get regularity theorems such as 'continuity implies infinitely many times differentiability'. Such type of results can be found e.g. in Járαι [4].

Finally, we remark that for linear, homogeneous, constant coefficient partial difference equations a method based on spectral synthesis was developed by László Székelyhidi in [9].

3. Elliptic partial difference equations

In this section we will investigate elliptic partial difference equations. In fact, as we shall see, they are nothing but the 'discrete' versions of elliptic partial differential equations. Here we remark that the investigations concerning

hyperbolic and parabolic partial difference equations will appear as a continuation of this work.

Among the most important of all partial differential equations are undoubtedly *Laplace's equation*

$$\Delta u(\mathbf{x}) = \sum_{i=1}^n \frac{\partial^2 u(\mathbf{x})}{\partial x_i^2} = 0 \quad (\mathbf{x} \in \Omega)$$

and the *Poisson equation*

$$-\Delta u(\mathbf{x}) = f(\mathbf{x}) \quad (\mathbf{x} \in \Omega).$$

In both equations the unknown function is $u: \overline{\Omega} \rightarrow \mathbb{R}$, where in general $\Omega \subset \mathbb{R}^n$ is assumed to be an open set. Further, in the second equation the function $f: \Omega \rightarrow \mathbb{R}$ is also given.

3.1. Auxiliary statements

In this subsection we list some preliminary tools that will be used subsequently, see also Evans [3] and Rudin [7].

Theorem 3.1 (Harnack's principle). *If the sequence functions $u_n: \Omega \rightarrow \mathbb{R}$ ($n \in \mathbb{R}$) is harmonic in the domain $\Omega \subseteq \mathbb{R}^n$ and*

$$u_1(x) \leq u_2(x) \leq \dots$$

at every point of Ω , then $\lim_{n \rightarrow \infty} u_n(x)$ either is infinite at every point of the domain or it is finite at every point of the domain, in both cases uniformly in each closed subdomain of Ω . In the latter case, the function

$$u(x) = \lim_{n \rightarrow \infty} u_n(x) \quad (x \in \Omega)$$

is harmonic in the domain Ω .

Remark 3.1. Harnack's principle can be generalized to monotone sequences of solutions of elliptic equations, as well.

Definition 3.1. We say that the function $\varphi \in C^\infty(\mathbb{R}^n)$ is a *mollifier*, if

(i) it is compactly supported

(ii)

$$\int_{\mathbb{R}^n} \varphi(x) dx = 1$$

(iii)

$$\lim_{\varepsilon \rightarrow 0+} \frac{1}{\varepsilon^n} \varphi\left(\frac{x}{\varepsilon}\right) = \delta(x) \quad (x \in \mathbb{R}^n),$$

where δ denotes the Dirac delta function and the limit is understood in the Schwartz space.

If the function φ is a mollifier and for all $x \in \mathbb{R}^n$ $\varphi(x) \geq 0$ holds, then φ is said to be *nonnegative mollifier*.

In case φ is a mollifier and there exists a function $\sigma \in \mathcal{C}^\infty(\mathbb{R})$ such that

$$\varphi(x) = \sigma(\|x\|) \quad (x \in \mathbb{R}^n),$$

then φ is termed to be a *(radially) symmetric mollifier*.

It is easy to see that if

$$\varphi(s) = \begin{cases} \exp\left(-\frac{1}{1-s}\right), & \text{if } s < 1 \\ 0, & \text{otherwise} \end{cases}$$

and

$$\omega(x) = \frac{\varphi(\|x\|^2)}{\int_{\mathbb{R}^n} \varphi(\|x\|^2) dx} \quad (x \in \mathbb{R}^n),$$

then the functions $\omega_\varepsilon: \mathbb{R}^n \rightarrow \mathbb{R}$ defined by

$$\omega_\varepsilon(x) = \frac{1}{\varepsilon^n} \omega\left(\frac{x}{\varepsilon}\right) \quad (x \in \mathbb{R}^n)$$

are nonnegative, symmetric mollifiers for all $\varepsilon > 0$.

Theorem 3.2 (Mollification Theorem). *Let $f \in \mathbb{L}^1(\mathbb{R}^n)$, then*

$$(i) \quad \lim_{\varepsilon \rightarrow 0+} \|f * \omega_\varepsilon - f\|_{\mathbb{L}^1} = 0$$

(ii) *for all $\varepsilon > 0$*

$$\text{supp}(f * \omega_\varepsilon) \subset \text{supp}(f) + \overline{B}(0, \varepsilon)$$

holds, where $\overline{B}(0, \varepsilon) = \{x \in \mathbb{R}^n \mid \|x\| \leq \varepsilon\}$.

(iii) *every sequence $(\varepsilon_n)_{n \in \mathbb{N}}$ of positive real numbers admits a subsequence $(\varepsilon_{n_k})_{k \in \mathbb{N}}$ such that $f * \omega_{\varepsilon_{n_k}}$ converges to f almost everywhere*

(iv) *if additionally, $f \in \mathcal{C}(\Omega)$, where $\Omega \subset \mathbb{R}^n$ is an open set, then*

$$\lim_{k \rightarrow \infty} f * \omega_{\varepsilon_{n_k}}(x) = f(x) \quad (x \in \Omega)$$

and the convergence is uniform on any compact subset of Ω

(v) for all $\varepsilon > 0$ we have $f * \omega_\varepsilon \in \mathcal{C}^\infty(\mathbb{R}^n)$, further, if $\alpha \in \mathbb{N}_0^n$ is a multi index, then

$$\partial^\alpha (f * \omega_\varepsilon) = f * (\partial^\alpha \omega_\varepsilon).$$

3.2. The discrete Laplace's equation

In this subsection we will investigate the *discrete Laplace's equation*, that is,

$$(\mathcal{L}) \quad \sum_{i=1}^n \Delta_h^2 u(\mathbf{x}) = \sum_{i=1}^n [u(\mathbf{x} + 2e_i h) - 2u(\mathbf{x} + e_i h) + u(\mathbf{x})] = 0.$$

Let us observe that in this case, the corresponding coefficient matrix $Q(\mathbf{x})$ is the $n \times n$ identity matrix at every point $\mathbf{x} \in \mathbb{R}^n$. Thus equation (\mathcal{L}) can be considered as the most representative example for the class of elliptic partial difference equations.

Theorem 3.3. *Let $u: \mathbb{R}^n \rightarrow \mathbb{R}$ be a continuous function and assume that u fulfills the discrete Laplace's equation on \mathbb{R}^n . Then the function u is harmonic on \mathbb{R}^n .*

Proof. Assume that for all $\varepsilon > 0$ the functions $\varphi_\varepsilon \in \mathcal{C}^\infty(\mathbb{R}^n)$ are nonnegative, symmetric mollifiers and let

$$u_\varepsilon(\mathbf{z}) = (u * \varphi_\varepsilon)(\mathbf{z}) = \int_{\mathbb{R}^n} u(\mathbf{x}) \varphi_\varepsilon(\mathbf{z} - \mathbf{x}) d\mathbf{x} \quad (\mathbf{z} \in \mathbb{R}^n).$$

Due to linearity of equation (\mathcal{L}) and because of the properties of the convolution,

$$\sum_{i=1}^n \Delta_h^2 u_\varepsilon(\mathbf{z}) = 0 \quad (\mathbf{z} \in \mathbb{R}^n).$$

In other words, this means that for all $\varepsilon > 0$ the functions u_ε also fulfill equation (\mathcal{L}) . Additionally, due to the Mollification Theorem, we also have that

$$\lim_{\varepsilon \rightarrow 0+} \|u_\varepsilon - u\|_{L^1} = 0.$$

Now let $(\varepsilon_n)_{n \in \mathbb{N}}$ be a null sequence for which the sequence of functions $(u_{\varepsilon_n})_{n \in \mathbb{N}}$ is monotone. From this we get that the limit $\lim_{n \rightarrow \infty} u_{\varepsilon_n}(\mathbf{z})$ exists for all $\mathbf{z} \in \mathbb{R}^n$ and the limit function is a harmonic function, too.

This means that the function u is a harmonic function.

Remark 3.2. In view of the previous theorem we immediately get that if a continuous function satisfies equation (\mathcal{L}) , then it is automatically in the class $\mathcal{C}^\infty(\mathbb{R}^n)$.

It is obvious however, that equation (\mathcal{L}) has nowhere continuous solutions, as well. To see this, let $\alpha: \mathbb{R} \rightarrow \mathbb{R}$ be a nowhere continuous additive function (see Corollary 5.2.2 in [6, page 130.]) and

$$u(\mathbf{x}) = u(x_1, \dots, x_n) = \prod_{i=1}^n \alpha(x_i) \quad (\mathbf{x} \in \mathbb{R}^n).$$

Finally, we remark that the method we used in the proof of Theorem 3.3 is also appropriate to solve more general elliptic equations such as

$$\frac{1}{h^2} \sum_{i=1}^m \gamma_i u(\mathbf{x} + \rho_i h) + u(\mathbf{x}) = \Phi(\mathbf{x}).$$

Assume namely that is equation is elliptic. Then the coefficient matrix (which will be independent of \mathbf{x} in this case), is

$$Q = \begin{pmatrix} A_{11} & \dots & A_{1n} \\ \vdots & \ddots & \vdots \\ A_{n1} & \dots & A_{nn} \end{pmatrix}$$

a definite matrix. Here,

$$A_{k,l} = \sum_{i=1}^m \gamma_i \rho_{k,i} \rho_{l,i} \quad (k, l = 1, \dots, n).$$

Thus applying the method based on the Mollification Theorem, we obtain that

$$\sum_{k,l=1}^n A_{k,l} \frac{\partial^2 u_\varepsilon(\mathbf{z})}{\partial x_k \partial x_l} + u_\varepsilon(\mathbf{z}) = \Phi_\varepsilon(\mathbf{z})$$

holds for all $\mathbf{z} \in \mathbb{R}^n$ and for all $\varepsilon > 0$, where

$$u_\varepsilon(\mathbf{z}) = (u * \varphi_\varepsilon)(\mathbf{z}) \quad \text{and} \quad \Phi_\varepsilon(\mathbf{z}) = (\Phi * \varphi_\varepsilon)(\mathbf{z}) \quad (\mathbf{z} \in \mathbb{R}^n).$$

Since the matrix Q is definite, this latter partial differential equation is of elliptic type. Thus, after a suitable change of the variables, it can be transformed into its canonical form. Using the theory of elliptic partial differential equations, the solution of the original partial difference equation can be obtained easily, with the help of Harnack's principle.

3.3. The discrete Poisson equation

Now we consider the *discrete Poisson equation*, that is,

$$(\mathcal{P}) \quad \frac{1}{h^2} \sum_{i=1}^n \Delta_h^2 u(\mathbf{x}) = f(\mathbf{x}),$$

where $f: \mathbb{R}^n \rightarrow \mathbb{R}$ is a given function.

Theorem 3.4. *Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be a continuous, compactly supported function and assume that $u: \mathbb{R}^n \rightarrow \mathbb{R}$ is also a continuous function which fulfills the discrete Poisson equation on \mathbb{R}^n . Then*

$$u(\mathbf{x}) = \int_{\mathbb{R}^n} \Phi(\mathbf{x} - \mathbf{y}) f(\mathbf{y}) d\mathbf{y} + C \quad (\mathbf{x} \in \mathbb{R}^n),$$

where C is a real constant and Φ stands for the fundamental solution of the Laplace's equation

$$\Phi(\mathbf{x}) = \frac{1}{2\pi} \ln(\|\mathbf{x}\|) \quad (\mathbf{x} \in \mathbb{R}^n)$$

if $n = 2$ and

$$\Phi(\mathbf{x}) = -\frac{1}{(n-2)s_n} \|\mathbf{x}\|^{-n+2} \quad (\mathbf{x} \in \mathbb{R}^n)$$

if $n \geq 3$, where s_n denotes the area of the n -dimensional unit sphere.

Proof. Similarly as in the previous theorem, assume that for all $\varepsilon > 0$ the functions $\varphi_\varepsilon \in \mathcal{C}^\infty(\mathbb{R}^n)$ are nonnegative, symmetric mollifiers and let

$$u_\varepsilon(\mathbf{z}) = \int_{\mathbb{R}^n} u(\mathbf{z}) \varphi_\varepsilon(\mathbf{z} - \mathbf{x}) d\mathbf{x} \quad (\mathbf{z} \in \mathbb{R}^n).$$

Due to the linearity of equation (\mathcal{P}) and because of the properties of the convolution, we obtain that

$$\sum_{i=1}^n \Delta_h^2 u_\varepsilon(\mathbf{z}) = h^2 f_\varepsilon(\mathbf{x}) \quad (\mathbf{z} \in \mathbb{R}^n),$$

where

$$f_\varepsilon(\mathbf{z}) = (f * \varphi_\varepsilon)(\mathbf{z}) \quad (\mathbf{z} \in \mathbb{R}^n).$$

In other words, this means that for all $\varepsilon > 0$ the functions u_ε fulfill a difference equation similar to equation (\mathcal{L}) . Additionally, due to the Mollification Theorem, we also have that

$$\lim_{\varepsilon \rightarrow 0^+} \|u_\varepsilon - u\|_{\mathbb{L}^1} = 0.$$

Since for all $\varepsilon > 0$, we have $u_\varepsilon, f_\varepsilon \in \mathcal{C}^\infty(\mathbb{R}^n)$, by differentiating twice with respect to h and putting $h = 0$, we obtain that

$$\Delta u_\varepsilon(\mathbf{z}) = f_\varepsilon(\mathbf{z}) \quad (\mathbf{z} \in \mathbb{R}^n),$$

yielding that for all $\varepsilon > 0$ the functions u_ε fulfill a Poisson equation. Applying a version of Harnack's principle concerning elliptic equations finally we get that the function u is a solution to the partial differential equation

$$\Delta u(\mathbf{z}) = f(\mathbf{z}) \quad (\mathbf{z} \in \mathbb{R}^n).$$

The statement of the theorem follows now from Theorem 1. of Evans [3, p. 22.].

3.4. The discrete biharmonic equation

Finally, we consider a fourth order partial difference equation, namely, the *discrete biharmonic equation*

$$(B) \quad \frac{1}{h^4} \sum_{i=1}^n \Delta_h^4 u(\mathbf{x}) + \frac{1}{h^4} \sum_{\substack{i,j=1 \\ i \neq j}}^n \Delta_h^4 u(\mathbf{x}) = 0.$$

The proof of the following result is similar to those in the previous subsections, hence we omit it.

Theorem 3.5. *Let $u: \mathbb{R}^n \rightarrow \mathbb{R}$ be a continuous function and assume that u fulfills the discrete biharmonic equation on \mathbb{R}^n . Then the function u is biharmonic on \mathbb{R}^n , that is,*

$$\Delta \Delta u(\mathbf{x}) = \sum_{i=1}^n \frac{\partial^4 u(\mathbf{x})}{\partial x_i^4} + \sum_{\substack{i,j=1 \\ i \neq j}}^n \frac{\partial^4 u(\mathbf{x})}{\partial x_i^2 \partial x_j^2} = 0 \quad (\mathbf{x} \in \mathbb{R}^n).$$

Remark 3.3. Similarly, as in case of the discrete Poisson equation, the solutions of the inhomogeneous biharmonic equation

$$\frac{1}{h^4} \sum_{i=1}^n \Delta_h^4 u(\mathbf{x}) + \frac{1}{h^4} \sum_{i,j=1}^n \Delta_h^4 u(\mathbf{x}) = \Phi(\mathbf{x})$$

can be derived in the same way as in the proof of Theorem 3.4.

References

- [1] **Baker, J.A.**, An analogue of the wave equation and certain related functional equations. *Canad. Math. Bull.*, **12** (1969), 837–846.
- [2] **Cheng, S.S.**, *Partial Difference Equations*, volume 3 of *Advances in Discrete Mathematics and Applications*. Taylor & Francis, London, 2003.
- [3] **Evans, L.C.**, *Partial Differential Equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.
- [4] **Járai, A.**, Continuity implies differentiability for solutions of functional equations—even with few variables. *Acta Sci. Math. (Szeged)*, **67(3-4)** (2001), 719–734.
- [5] **McKiernan, M.A.**, Boundedness on a set of positive measure and the mean value property characterizes polynomials on a space V^n . *Aequationes Math.*, **4** (1970), 31–36.
- [6] **Kuczma, M.**, *An Introduction to the Theory of Functional Equations and Inequalities. Cauchy's Equation and Jensen's Inequality.*, Second edition, Birkhäuser Verlag, Basel, 2009.
- [7] **Rudin, W.**, *Real and Complex Analysis*. McGraw-Hill Book Co., New York, third edition, 1987.
- [8] **Świątak, H.**, The regularity of the locally integrable and continuous solutions of nonlinear functional equations. *Trans. Amer. Math. Soc.*, **221(1)** (1976), 97–118.
- [9] **Székelyhidi, L.**, *Discrete Spectral Synthesis and its Applications*, Springer Monographs in Mathematics. Springer, Dordrecht, 2006.

Eszter Gselmann

Institute of Mathematics

University of Debrecen

Egyetem tér 1.

Debrecen

Hungary

gselmann@science.unideb.hu