

ON THE MOMENTS OF ROOTS OF LAGUERRE-POLYNOMIALS AND THE MARCHENKO–PASTUR LAW

Miklós Kornyik and György Michaletzky

(Budapest, Hungary)

Dedicated to remembering Antal Iványi

Communicated by Ferenc Schipp

(Received June 14, 2017; accepted August 12, 2017)

Abstract. In this paper we compute the leading terms in the sum of the k^{th} power of the roots of $L_p^{(\alpha)}$, the Laguerre-polynomial of degree p with parameter α . The connection between the Laguerre-polynomials and the Marchenko–Pastur distribution is expressed by the fact, among others, that the limiting distribution of the empirical distribution of the normalized roots of the Laguerre-polynomials is given by the Marchenko–Pastur distribution. We give a direct proof of this statement based on the recursion satisfied by the Laguerre-polynomials. At the same time, our main result gives that the leading term in p and $(\alpha + p)$ of the sum of the k^{th} power of the roots of $L_p^{(\alpha)}$ coincides with the k^{th} moment of the Marchenko–Pastur law. We also mention the fact that the expectation of the characteristic polynomial of a XX^T type random covariance matrix, where X is a $p \times n$ random matrix with iid elements, is $\ell_p^{(n-p)}$, i.e. the monic version of the p^{th} Laguerre polynomial with parameter $n - p$.

1. Introduction

In theory of orthogonal polynomials the limit of the empirical distribution of their roots is a much studied matter. In this paper we are going to study

Key words and phrases: Random covariance matrix, Laguerre polynomials, Marchenko–Pastur law

2010 Mathematics Subject Classification: 15A52, 60B20, 33C45.

<https://doi.org/10.71352/ac.46.137>

the limit distribution of the roots of Laguerre polynomials $L_p^{(\alpha_p)}$, where

$$(1.1) \quad L_p^{(\alpha)}(x) = \sum_{j=0}^p (-1)^j \binom{\alpha+p}{p-j} \frac{x^j}{j!} \quad \alpha \in \mathbb{R}$$

assuming that $\alpha_p/p \rightarrow c > -1$. For $\alpha > -1$ these polynomials are known to be orthogonal with respect to the measure $x^\alpha e^{-x} \mathbf{1}_{[0,\infty)} dx$, from which one can conclude that all the roots are distinct and lie in \mathbb{R}_+ . For $\alpha \in [-p+1, -1] \cap \mathbb{Z}$ one has that

$$L_p^{(\alpha)}(x) = x^{-\alpha} L_{p+\alpha}^{(-\alpha)}(x)$$

and hence one can make the conclusion that for such α values the polynomial $L_p^{(\alpha)}$ has $p+\alpha$ distinct positive roots and 0 is also a root with multiplicity $-\alpha$.

In Section 2 we show that the normalized generating function of the moments of the normalized roots of $L_p^{(\alpha_p)}$ satisfies the same quadratic fixed point equation in the limit as the generating function of the moments of the Marchenko–Pastur distribution.

In Section 3 we will explicitly show that the coefficient of the highest order term (viewed as a polynomial in p) of the k^{th} power of the roots of $L_p^{(\alpha)}$ coincides with the k^{th} moment of the corresponding Marchenko–Pastur distribution.

2. Convergence of the empirical distribution

Let us consider the roots of the Laguerre-polynomial $L_p^{(\alpha)}$ denoted by $\xi_{p,1}^{(\alpha)}, \dots, \xi_{p,p}^{(\alpha)}$. Let $M_p^{(\alpha)}(k)$ denote the sum of their k -th power. That is $M_p^{(\alpha)}(k) = \sum_{i=1}^p (\xi_{p,i}^{(\alpha)})^k$. Finally, $\mathcal{M}_p^{(\alpha)}$ denotes the power series determined by these coefficients, i.e.

$$(2.1) \quad \mathcal{M}_p^{(\alpha)}(z) = p + \sum_{k=1}^{\infty} M_p^{(\alpha)}(k) z^k.$$

Note that in case α is a negative integer in the interval $[-p+1, -1]$ zero is also a root of $L_p^{(\alpha)}$, which explains why the case $k=0$, i.e. the zeroth moment, had to be dealt with separately in (2.1). It is known that

$$\mathcal{M}_p^{(\alpha)}(z) = \frac{1}{z} \frac{(\ell_p^{(\alpha)})'(1/z)}{\ell_p^{(\alpha)}(1/z)} = -z \frac{(\tilde{\ell}_p^{(\alpha)})'(z)}{\tilde{\ell}_p^{(\alpha)}(z)} + p,$$

where $\ell_p^{(\alpha)}(x) = (-1)^p p! L_p^{(\alpha)}(x)$ is the monic version of $L_p^{(\alpha)}$, and for any polynomial of degree p we denote by $\widehat{\ell}(z) = z^p \ell(1/z)$ the so-called conjugate polynomial.

Theorem 2.1. *Let us assume that $\alpha = \alpha_p$ and $\frac{\alpha_p}{p} \rightarrow c \in (-1, \infty)$, as $p \rightarrow \infty$. Then the empirical distribution determined by the normalized roots (where p^{-1} is the normalization factor) of the Laguerre-polynomial $L_p^{(\alpha_p)}$ converges weakly to the Marchenko–Pastur distribution, given as*

$$(2.2) \quad \mu_c(A) = \begin{cases} -c\delta_0(A) + \nu_c(A), & \text{if } -1 < c < 0 \text{ and } \alpha_p \in \{-p+1, \dots, -1\} \\ & \text{for all } p, \\ \nu_c(A), & \text{if } c \geq 0, \end{cases}$$

for $A \in \mathcal{B}(\mathbb{R})$, where δ_0 denotes the Dirac-delta measure at 0, while the measure ν_c is absolutely continuous with density

$$d\nu_c(x) = \frac{\sqrt{(x_+ - x)(x - x_-)}}{2\pi x} \mathbf{1}_{[x_-, x_+]}(x) dx,$$

where $x_{\pm} = [\sqrt{c+1} \pm 1]^2$.

Remark 2.1. A more general version of this theorem – allowing for $c < -1$ – was proved by Martínez-González et al. in [3] using complex analysis and differential equations, but the proof presented here is based on elementary calculations using only the recursion equations satisfied by the Laguerre-polynomials.

Remark 2.2. Laguerre polynomials show a deep connection with random matrix theory in the following ways:

1. Forrester and Gamburd proved in [1] that the expectation of the characteristic polynomial of the random matrix XX^T is given by $\ell_p^{(n-p)}(z)$, i.e. $E \det(x \cdot I - XX^T) = \ell_p^{(n-p)}(x)$, where X is a $p \times n$ random matrix with independent, identically distributed entries with zero expectation and variance 1.
2. If X is a $p \times n$ random matrix in the same sense as above, then the weak limit of the empirical measure of the eigenvalues is a much studied question of random matrix theory, although it is usually normalized by n , which in our case means a normalization by $\alpha + p$. A well-known theory – proved by Marchenko and Pastur in [2] – states that the weak limit of the empirical measure of the eigenvalues of $\frac{1}{n}XX^T$ is given by $\tilde{\mu}_a$ as $\frac{p}{n} \rightarrow a > 0$, where $\tilde{\mu}_a$ is defined below. In the case of the present paper μ_c is the weak limit of the empirical measure of the eigenvalues of $\frac{1}{p}XX^T$.

3. The matrix theoretical Marchenko–Pastur distribution with parameter $a > 0$ is given by

$$\tilde{\mu}_a(A) = \begin{cases} (1 - \frac{1}{a}) \delta_0 + \tilde{\nu}_a(A), & \text{if } a \in (0, 1) \\ \tilde{\nu}_a(A) & \text{if } a \geq 1 \end{cases} \quad A \in \mathcal{B}(\mathbb{R}).$$

with $\tilde{\nu}_a$ being absolutely continuous with density

$$d\tilde{\nu}_a(x) = \frac{\sqrt{(x - \tilde{x}_-)(\tilde{x}_+ - x)}}{2\pi ax} \mathbf{1}_{[\tilde{x}_-, \tilde{x}_+]}(x) dx,$$

where $\tilde{x}_{\pm} = (1 \pm \sqrt{a})^2$. As mentioned before this version of the Marchenko–Pastur arises when the zeros of $\ell_p^{(\alpha_p)}(z)$ are normalized by a factor of $(p + \alpha_p)^{-1}$. The connection between $d\mu_c$ and $d\tilde{\mu}_a$ is the following:

$$(2.3) \quad a = \frac{1}{c + 1},$$

$$(2.4) \quad \mu_c = \tilde{\mu}_a \circ g^{-1},$$

where $g(x) = (c + 1)x$ for $x \in \mathbb{R}$. On the other hand it is known that the moments of $\tilde{\mu}_a$ are given by

$$(2.5) \quad \int x^k d\tilde{\mu}_a(x) = \sum_{j=1}^k \frac{1}{k} \binom{k}{j} \binom{k}{j-1} a^{j-1},$$

hence the moments of μ_c can be calculated as

$$\int x^k d\mu_c = \sum_{j=1}^k \frac{1}{k} \binom{k}{j} \binom{k}{j-1} (c + 1)^{k-j+1}.$$

Proof of the Theorem 2.1. Let $\ell_p^{(\alpha)}(z) := (-1)^p p! L_p^{(\alpha)}(z)$ denote the monic version of $L_p^{(\alpha)}(z)$ then

$$(2.6) \quad \ell_p^{(\alpha)}(z) = \sum_{j=0}^p (-1)^j \frac{(p)_j (\alpha + p)_j}{j!} z^{p-j}$$

with $(\beta)_k = \beta(\beta - 1) \cdots (\beta - k + 1)$ for $k > 0$. Note that if β is a positive integer and $k > \beta$ then $(\beta)_k = 0$. Thus it follows from (2.6) that for $\alpha \in \{-p + 1, -p + 2, \dots, -2, -1\}$

$$(2.7) \quad \ell_p^{(\alpha)}(z) = z^{-\alpha} \ell_{p+\alpha}^{(-\alpha)}(z).$$

This means that in this case zero is a root of $\ell_p(x)^{(\alpha)}(z)$ with multiplicity $-\alpha$ and the other $p + \alpha$ roots given by the Laguerre-polynomial $L_{p+\alpha}^{(-\alpha)}(z)$ of degree $p + \alpha$.

Case 1. Let us first consider the case when $\alpha_p \geq 0$ for all p , which also implies $\lim \alpha_p/p = c \geq 0$.

The recursion of the Laguerre-polynomials for arbitrary parameter $\alpha > -1$ is

$$(2.8) \quad a_p L_{p+1}^{(\alpha)}(z) = (b_p - z) L_p^{(\alpha)}(z) - c_p L_{p-1}^{(\alpha)}(z),$$

where $a_p = p + 1$, $b_p = 2p + \alpha + 1$ and $c_p = p + \alpha$ and also

$$(2.9) \quad p L_p^{(\alpha)}(z) = (p + \alpha) L_{p-1}^{(\alpha)}(z) - z L_{p-1}^{(\alpha+1)}(z).$$

These polynomials are known to be orthogonal with respect to the measure $z^\alpha e^{-z} \mathbf{1}_{[0, \infty)}(z) dz$, which implies that all the roots of $L_p^{(\alpha)}(x)$ lie in the interval $[0, \infty)$ and hence the sum of the k^{th} power of its roots is positive. Furthermore

$$(2.10) \quad \frac{d}{dz} L_p^{(\alpha_p)}(z) = -L_{p-1}^{(\alpha_p+1)}(z)$$

implying, after proper algebraic transformations, that

$$(2.11) \quad \frac{d}{dz} \widehat{\ell}_p^{(\alpha)}(z) = -(\alpha + p) p \widehat{\ell}_{p-1}^{(\alpha)}(z),$$

where $\widehat{\ell}_p^{(\alpha)}(z) = z^p \ell_p^{(\alpha)}(z^{-1})$. Applying this for $\alpha = \alpha_p$ we obtain that

$$(2.12) \quad \frac{1}{p} \mathcal{M}_p^{(\alpha_p)}\left(\frac{z}{p}\right) = \frac{\alpha_p + p}{p} \frac{z \widehat{\ell}_{p-1}^{(\alpha_p)}(z/p)}{\widehat{\ell}_p^{(\alpha_p)}(z/p)} + 1.$$

Also from recursion (2.8) we get

$$(2.13) \quad \widehat{\ell}_{p+1}^{(\alpha)}(z) = [1 - (\alpha + 2p + 1)z] \widehat{\ell}_p^{(\alpha)}(z) - z^2(p + \alpha) p \widehat{\ell}_{p-1}^{(\alpha)}(z).$$

Since the largest zero of $L_p^{(\alpha)}$ is no greater than $4p + 2\alpha + 3$ (see [4]) we obtain that $\widehat{\ell}_p^{(\alpha)}(z) > 0$, if $0 \leq z < \frac{1}{4p+2\alpha+3}$.

In this case one has that

$$\frac{\widehat{\ell}_{p-1}^{(\alpha)}(z)}{\widehat{\ell}_p^{(\alpha)}(z)} \leq \frac{1 - (\alpha + 2p + 1)z}{z^2 p (\alpha + p)} \leq \frac{1}{(p + \alpha) p z^2}.$$

Using the computations above we get that

$$\begin{aligned}
 \frac{d}{dz} \frac{\widehat{\ell}_{p-1}^{(\alpha)}(z)}{\widehat{\ell}_p^{(\alpha)}(z)} &= \frac{(\widehat{\ell}_{p-1}^{(\alpha)}(z))' \widehat{\ell}_p^{(\alpha)}(z) - (\widehat{\ell}_p^{(\alpha)}(z))' \widehat{\ell}_{p-1}^{(\alpha)}(z)}{(\widehat{\ell}_p^{(\alpha)}(z))^2} = \\
 (2.14) \quad &= (\alpha + p)p \left[\left(\frac{\widehat{\ell}_{p-1}^{(\alpha)}(z)}{\widehat{\ell}_p^{(\alpha)}(z)} \right)^2 - \frac{\widehat{\ell}_{p-2}^{(\alpha)}(z)}{\widehat{\ell}_p^{(\alpha)}(z)} \right] + (\alpha + 2p - 1) \frac{\widehat{\ell}_{p-2}^{(\alpha)}(z)}{\widehat{\ell}_p^{(\alpha)}(z)}.
 \end{aligned}$$

Since in the present case $\mathcal{M}_p^{(\alpha)}(z)$ is a convex, monotonically increasing function for $z \geq 0$, and furthermore $\mathcal{M}_p^{(\alpha)}(0) = 1$, one has

$$(2.15) \quad z \frac{d}{dz} \frac{\widehat{\ell}_{p-1}^{(\alpha)}(z)}{\widehat{\ell}_p^{(\alpha)}(z)} \leq \int_0^{2z} \frac{d}{dt} \frac{\widehat{\ell}_{p-1}^{(\alpha)}(t)}{\widehat{\ell}_p^{(\alpha)}(t)} dt = \frac{\widehat{\ell}_{p-1}^{(\alpha)}(2z)}{\widehat{\ell}_p^{(\alpha)}(2z)} - 1 \leq \frac{1}{4(\alpha + p)pz^2}$$

and so according to (2.14) and to (2.15) we have

$$\begin{aligned}
 (2.16) \quad \left| \left(\frac{\widehat{\ell}_{p-1}^{(\alpha)}(z/p)}{\widehat{\ell}_p^{(\alpha)}(z/p)} \right)^2 - \frac{\widehat{\ell}_{p-2}^{(\alpha)}(z/p)}{\widehat{\ell}_p^{(\alpha)}(z/p)} \right| &\leq \frac{p^3}{4(\alpha + p)^2 p^2 z^3} + \frac{\alpha + 2p - 1}{(\alpha + p)^2 p^2} \times \\
 &\times \frac{p^4}{(p - 1)(\alpha + p - 1)z^4}.
 \end{aligned}$$

Let $f_p^{(\alpha)}(z) := \frac{\widehat{\ell}_{p-1}^{(\alpha)}(z/p)}{\widehat{\ell}_p^{(\alpha)}(z/p)}$, for $p \geq 1$. According to (2.16) we have

$$f_p^{(\alpha)}(z) f_{p-1}^{(\alpha)} \left(z \frac{p-1}{p} \right) - (f_p^{(\alpha)})^2(z) \rightarrow 0$$

if $p \rightarrow \infty$ and $\alpha = \alpha_p \geq 0$, especially $f_p^{(\alpha_p)}(z) f_{p-1}^{(\alpha_p)}((p-1)z/p) - (f_p^{(\alpha_p)})^2(z) \rightarrow 0$ if $\frac{\alpha_p}{p} \rightarrow c$ as $p \rightarrow \infty$.

Applying (2.13) to $\widehat{\ell}_p^{(\alpha)}$ one has

$$\begin{aligned}
 1 &= \left(1 - z \frac{\alpha + 2p - 1}{p} \right) f_p^{(\alpha)}(z) - \\
 (2.17) \quad &- z^2 \frac{(p-1)(\alpha + p - 1)}{p^2} f_p^{(\alpha)}(z) f_{p-1}^{(\alpha)} \left(z \frac{p-1}{p} \right),
 \end{aligned}$$

hence we get that the accumulation points of $(f_p^{(\alpha_p)}(z))_{p \in \mathbb{N}}$ as $\alpha_p/p \rightarrow c$ satisfy the following equation in ξ

$$(2.18) \quad 1 = [1 - (c + 2)z] \xi - (c + 1)z^2 \xi^2.$$

The solutions of this equation are

$$\xi_{\pm} = \frac{1 - (c+2)z \pm \sqrt{[1 - (c+2)z]^2 - 4(c+1)z^2}}{2(c+1)z^2}.$$

Let us introduce the notation

$$f_{c-}(z) := \frac{1 - (c+2)z - \sqrt{[1 - (c+2)z]^2 - 4(c+1)z^2}}{2(c+1)z^2}.$$

In order to find the appropriate root let us look at the map $\xi \rightarrow \eta_c(\xi, z)$ for a fixed z defined by

$$1 = [1 - (c+2)z] \eta_c(\xi, z) - (c+1)z^2 \xi \eta_c(\xi, z)$$

and hence

$$\eta_c(\xi, z) = \frac{1}{1 - (c+2)z - (c+1)z^2 \xi}.$$

Note that the fixed points of this mapping are the solutions of (2.18).

In parallel with this for any fixed $\alpha \geq 0$ and $p \geq 1$ consider the following equation in ξ :

$$(2.19) \quad 1 = \left[1 - z \frac{\alpha + 2p - 1}{p} \right] \xi - z^2 \frac{(p-1)(\alpha + p - 1)}{p^2} \xi^2.$$

Denote by $\zeta_p^{(\alpha)}$ the largest nonnegative z value, for which both roots of this second-order equation are non-negative, i.e.

$$\zeta_p^{(\alpha)} = \sup \left\{ z \mid z \frac{\alpha + 2p - 1}{p} \leq 1, \right. \\ \left. 4z^2 \frac{(p-1)(\alpha + p - 1)}{p^2} \leq \left(1 - z \frac{\alpha + 2p - 1}{p} \right)^2 \right\}.$$

Short calculation shows that $\zeta_p^{(\alpha)} = \left(a_p^{(\alpha)} + 2\sqrt{b_p^{(\alpha)}} \right)^{-1}$, where $a_p^{(\alpha)} = \frac{\alpha + 2p - 1}{p}$ and $b_p^{(\alpha)} = \frac{(p-1)(\alpha + p - 1)}{p^2}$.

Now for $0 \leq z < \zeta_p^{(\alpha)}$ define the map $\xi \rightarrow \eta_p^{(\alpha)}(\xi, z)$ as the solution to

$$1 = \left[1 - z \frac{\alpha + 2p - 1}{p} \right] \eta_p^{(\alpha)}(\xi, z) - z^2 \frac{(p-1)(\alpha + p - 1)}{p^2} \eta_p^{(\alpha)}(\xi, z) \xi.$$

Thus

$$\eta_p^{(\alpha)}(\xi, z) = \frac{1}{1 - z \frac{\alpha + 2p - 1}{p} - z^2 \frac{(p-1)(\alpha + p - 1)}{p^2} \xi}.$$

Observe that $\eta_p^{(\alpha_p)}(\xi, z) \xrightarrow{p \rightarrow \infty} \eta_c(\xi, z) \quad \forall (\xi, z) \in \mathbb{R}^2$.

For the small positive values of ξ the functions $\eta_c(\xi, z)$ and $\eta_p^{(\alpha)}(\xi, z)$ are increasing. Let us denote by $g_{p-}^{(\alpha)}(z)$ the smaller fixed point of the mapping $\eta_p^{(\alpha)}(\xi, z)$ and observe that for $0 \leq z < \zeta_p^{(\alpha)}$ the inequality $\eta_p^{(\alpha)}(0, z) > 0$ holds true, thus for $0 \leq \xi < g_{p-}^{(\alpha)}(z)$ we have that

$$\xi < \eta_p^{(\alpha)}(\xi, z) \leq g_{p-}^{(\alpha)}(z).$$

We are going to prove by induction on p that for any fixed $\alpha \geq 0$ and $0 \leq z < \zeta_p^{(\alpha)}$ the inequality

$$(2.20) \quad f_p^{(\alpha)}(z) \leq g_{p-}^{(\alpha)}(z)$$

holds true. It is easy to check that for $p = 1$ we have that $\zeta_1^{(\alpha)} = \frac{1}{\alpha+1}$ and

$$g_{1-}^{(\alpha)}(z) = f_1^{(\alpha)}(z) = \frac{1}{1 - z(\alpha + 1)}.$$

On the other hand straightforward calculation gives that if $0 \leq z < \zeta_p^{(\alpha)}$ then $z \frac{p-1}{p} < \zeta_{p-1}^{(\alpha)}$ thus using the induction hypothesis for $p-1$ we obtain that

$$(2.21) \quad f_{p-1}^{(\alpha)}\left(z \frac{p-1}{p}\right) \leq g_{(p-1)-}^{(\alpha)}\left(z \frac{p-1}{p}\right).$$

The latter one is the smaller fixed point of the mapping

$$\eta_{p-1}^{(\alpha)}\left(\cdot, z \frac{p-1}{p}\right) : \xi \rightarrow \frac{1}{1 - z \frac{p-1}{p} \frac{\alpha+2p-3}{p-1} - z^2 \frac{(p-1)^2}{p^2} \frac{(p-2)(\alpha+p-2)}{(p-1)^2} \xi}.$$

On the other hand

$$\begin{aligned} \eta_{p-1}^{(\alpha)}\left(\xi, z \frac{p-1}{p}\right) &= \frac{1}{1 - z \frac{p-1}{p} \frac{\alpha+2p-3}{p-1} - z^2 \frac{(p-1)^2}{p^2} \frac{(p-2)(\alpha+p-2)}{(p-1)^2} \xi} = \\ &= \frac{1}{1 - z \frac{\alpha+2p-3}{p} - z^2 \frac{(p-2)(\alpha+p-2)}{p^2} \xi} \leq \\ &\leq \frac{1}{1 - z \frac{\alpha+2p-1}{p} - z^2 \frac{(p-1)(\alpha+p-1)}{p^2} \xi} = \eta_p^{(\alpha)}(\xi, z), \end{aligned}$$

proving that

$$(2.22) \quad g_{(p-1)-}^{(\alpha)}\left(z \frac{p-1}{p}\right) \leq g_{p-}^{(\alpha)}(z).$$

But equation (2.17) implies that for $\xi = f_{p-1}^{(\alpha)} \left(z^{\frac{p-1}{p}} \right)$

$$\eta_p^{(\alpha)}(\xi, z) = f_p^{(\alpha)}(z).$$

Comparing (2.21) and (2.22) we obtain that for $0 \leq z < \zeta_p^{(\alpha)}$

$$f_p^{(\alpha)}(z) \leq g_{p-}^{(\alpha)}(z)$$

proving the induction step.

Since $a_p^{(\alpha_p)} \rightarrow c + 2$, $b_p^{(\alpha_p)} \rightarrow c + 1$ and so $\zeta_p^{(\alpha_p)} \rightarrow \frac{1}{(\sqrt{c+1}+1)^2}$, if $\alpha_p/p \rightarrow c$ as $p \rightarrow \infty$ the following implication holds for large enough p :

$$\left[0, \frac{1}{2(\sqrt{c+1}+1)^2} \right) \subset \left[0, \zeta_p^{(\alpha_p)} \right).$$

Hence for $0 \leq z < \frac{1}{2(\sqrt{c+1}+1)^2}$ we have that $g_{p-}^{(\alpha_p)}(z) \xrightarrow{p \rightarrow \infty} f_{c-}(z)$ as $p \rightarrow \infty$, thus inequality (2.20) implies that

$$\lim_p f_p^{(\alpha_p)}(z) = f_{c-}(z).$$

Now let $\mathfrak{M}_c(z) = \lim_p \frac{1}{p} \mathcal{M}_p^{(\alpha_p)} \left(\frac{z}{p} \right)$. According to (2.12) we have that

$$\mathfrak{M}_c(z) = (c+1)z f_{c-}(z) + 1$$

from which one has

$$\begin{aligned} (2.23) \quad \mathfrak{M}_c(z) &= \frac{1 - cz - \sqrt{[1 - (c+2)z]^2 - 4(c+1)z^2}}{2z} = \\ &= \frac{1 - cz - \sqrt{(1 - cz)^2 - 4z}}{2z}. \end{aligned}$$

Case 2. Consider now the case when $\alpha_p \in \{-p+1, \dots, -1\}$ for all p in such a way that $\lim \alpha_p/p = c$ exists and $c > -1$. Obviously this implies $c \leq 0$.

In this case the recursion (2.8) is still valid, but orthogonality (with respect to $z^{\alpha_p} e^{-z} \mathbf{1}_{[0, \infty)}(z) dz$) cannot be assured. According to (2.6) one has that

$$\ell_p^{(\alpha_p)}(z) = z^{-\alpha} \ell_{p+\alpha_p}^{(-\alpha_p)}(z)$$

and so

$$\begin{aligned} \widehat{\ell}_p^{(\alpha_p)}(z) &= z^p \ell_p^{(\alpha_p)}(1/z) = z^p z^{\alpha_p} \ell_{p+\alpha_p}^{(-\alpha_p)}(1/z) = z^{p+\alpha_p} \ell_{p+\alpha_p}^{(-\alpha_p)}(1/z) = \\ &= \widehat{\ell}_{p+\alpha_p}^{(-\alpha_p)}(z) \end{aligned}$$

hence

$$\mathcal{M}_p^{(\alpha_p)}(z) = -z \frac{\frac{d}{dz} \widehat{\ell}_{p+\alpha_p}^{(-\alpha_p)}(z)}{\widehat{\ell}_{p+\alpha_p}^{(-\alpha_p)}(z)} + p = \mathcal{M}_{p+\alpha_p}^{(-\alpha_p)}(z) - \alpha_p,$$

therefore we immediately get that $\mathcal{M}_p^{(\alpha_p)}(z)$ is monotonically increasing convex function if $z \geq 0$ and

$$\frac{1}{p} \mathcal{M}_p^{(\alpha_p)}\left(\frac{z}{p}\right) = \frac{p+\alpha_p}{p} \frac{1}{p+\alpha_p} \mathcal{M}_{p+\alpha_p}^{(-\alpha_p)}\left(\frac{p+\alpha_p}{p} \frac{z}{p+\alpha_p}\right) - \frac{\alpha_p}{p}.$$

Due to $\alpha_p/p \rightarrow c$ we have $-\frac{\alpha_p}{p+\alpha_p} \rightarrow -\frac{c}{c+1}$ and

$$\frac{1}{p+\alpha_p} \mathcal{M}_{p+\alpha_p}^{(-\alpha_p)}\left(\frac{z}{p+\alpha_p}\right) \rightarrow \mathfrak{M}_{-\frac{c}{c+1}}(z).$$

Since $f_{p+\alpha_p}^{(-\alpha_p)}$ is a sequence with uniformly bounded derivatives (according to (2.15)) one has that for $0 \leq z < [2(\sqrt{c+1}+1)^2]^{-1}$

$$\mathfrak{M}_c(z) = (c+1) \mathfrak{M}_{-\frac{c}{c+1}}((c+1)z - c)$$

implying that

$$\mathfrak{M}_c(z) = \frac{1 - cz - \sqrt{(1 - cz)^2 - 4z}}{2z}.$$

Since $\mathfrak{M}_c(z)$ coincides with the generating function of the moments of μ_c , i.e.

$$\mathfrak{M}_c(z) = \sum_{k \geq 0} \int x^k d\mu_c(x) \cdot z^k, \quad \text{for } z \in [0, \frac{1}{2}(\sqrt{c+1}+1)^{-2}]$$

and μ_c is fully determined by its moments we have that the weak limit of the empirical measure of the normalized zeros of $\ell_p^{(\alpha_p)}(z)$ is μ_c .

Theorem 2.1 is hereby proved. ■

Corollary 2.1. *Theorem 2.1 also implies the convergence of the moments of the empirical distribution of the normalized roots of $\ell_p^{(\alpha_p)}$. In other words if $m_p^{(\alpha_p)}$ denotes the empirical distribution of the normalized roots of $\ell_p^{(\alpha_p)}$, then*

$$\int x^k dm_p^{(\alpha_p)}(x) \xrightarrow{p \rightarrow \infty} \int x^k d\mu_c(x) = \sum_{j=1}^k \frac{1}{k} \binom{k}{j} \binom{k}{j-1} (c+1)^{k-j+1} \quad \forall k \geq 0$$

when $\frac{\alpha_p}{p} \rightarrow c$.

3. The sum of the k^{th} power of the roots of $L_p^{(\alpha)}$

The following theorem shows that the connection between the root distribution of the Laguerre-polynomials and the Marchenko–Pastur distribution is not only an asymptotic connection but in a ”dominating way” it holds for large enough p values, as well.

Theorem 3.1. *Let $p \in \mathbb{N}$, $M_p^{(\alpha)}(k) := \sum_{j=1}^p \xi_{p,j}^k$, where $0 \leq \xi_{p,1}^{(\alpha)} < \xi_{2,p}^{(\alpha)} < \dots < \xi_{p,p}^{(\alpha)} < \infty$ denotes the roots of $L_p^{(\alpha)}$. Then for $\alpha \in \mathbb{R}$, $\alpha + p > k - 1$ one has*

$$M_p^{(\alpha)}(k) = \sum_{j=1}^k \frac{1}{k} \binom{k}{j} \binom{k}{j-1} p^j (\alpha + p)^{k-j+1} + f(\alpha + p, p),$$

where f is a polynomial in two variables with $\deg f \leq k$.

In case $\alpha + p \leq k - 1$ one has that the coefficient of the dominating term in $M_p^{(\alpha)}(k)$ is less than or equal to the quantity above.

Proof. Let us consider the Newton identities $\sum_{j=0}^{k-1} M_n^{(\alpha)}(k-j) a_{p-j} = -k a_{n-k}$, where a_{p-j} denotes the corresponding coefficient of $\ell_p^{(\alpha)}(x)$. It is known that $a_j = (-1)^{p+j} p! \binom{\alpha+p}{p-j} \frac{1}{j!}$ (see e.g. [4]), hence

$$a_{p-j} = (-1)^j \frac{(\alpha + p)_j (p)_j}{j!}.$$

Writing the Newton identities in matrix form we obtain that

$$(3.1) \quad \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ a_{p-1} & 1 & 0 & \dots & 0 \\ a_{p-2} & a_{p-1} & 1 & \dots & 0 \\ & & & \ddots & \\ a_{p-(k-1)} & a_{p-(k-2)} & a_{p-(k-3)} & \dots & 1 \end{bmatrix} \begin{bmatrix} M_p^{(\alpha)}(1) \\ M_p^{(\alpha)}(2) \\ M_p^{(\alpha)}(3) \\ \vdots \\ M_p^{(\alpha)}(k) \end{bmatrix} = \begin{bmatrix} -a_{p-1} \\ -2a_{p-2} \\ -3a_{p-3} \\ \vdots \\ -ka_{p-k} \end{bmatrix}.$$

Thus

$$M_p^{(\alpha)}(k) = \det \begin{bmatrix} 1 & 0 & 0 & \dots & -a_{p-1} \\ a_{p-1} & 1 & 0 & \dots & -2a_{p-2} \\ a_{p-2} & a_{p-1} & 1 & \dots & -3a_{p-3} \\ & & & \ddots & \\ a_{p-(k-1)} & a_{p-(k-2)} & a_{p-(k-3)} & \dots & -ka_{p-k} \end{bmatrix}$$

according to Cramer's rule and the fact that the determinant of the matrix in (3.1) is 1. In general, let us introduce the following notation:

$$A(k, l) := \det \begin{bmatrix} 1 & 0 & \dots & (\alpha + p)_l \cdot p \\ a_{p-1} & 1 & \dots & -2 \frac{(\alpha + p)_{l+1} \cdot (p)_2}{2} \\ & & \ddots & \\ a_{p-(k-1)} & a_{p-(k-2)} & \dots & -k \frac{(-1)^k (\alpha + p)_{l-k+1} \cdot (p)_k}{k!} \end{bmatrix}$$

for $k \geq 2, l \geq 1$ and $A(1, l) = (p + \alpha)_l p$ for $l \geq 1$. With this notation $A(k, 1) = M_p^{(\alpha)}(k)$ and it can be proved by induction that for $k \geq 2$

$$(3.2) \quad A(k, l) = \sum_{r=1}^l p(\alpha + p - r)_{l-r} A(k-1, r) + A(k-1, l+1),$$

In fact, for $k \geq 3$ let us subtract $p(\alpha + p)_l$ times the first column of the matrix in the definition of $A(k, l)$ from the last of the same. The j^{th} element of the last column obtained this way can be written as

$$\begin{aligned} & -(-1)^j \frac{(\alpha + p)_{l+j-1} (p)_j}{(j-1)!} - (-1)^{j-1} \frac{(\alpha + p)_l p (\alpha + p)_{j-1} (p)_{j-1}}{(j-1)!} = \\ & = -(-1)^{j-1} \frac{(\alpha + p)_{j-1} (p)_{j-1}}{(j-1)!} [(\alpha + p - j + 1)_l (p - j + 1) - (\alpha + p)_l p] = \\ & = (-1)^{j-1} \frac{(\alpha + p)_{j-1} (p)_{j-1}}{(j-1)!} (j-1) \left(\sum_{r=1}^l (\alpha + p - j + 1)_r (\alpha + p - r)_{l-r} p + 1 \right) \end{aligned}$$

due to

$$\prod_{i=1}^m c_i - \prod_{i=1}^m d_i = \sum_{h=1}^m \prod_{1 \leq e < h} c_e (c_h - d_h) \prod_{m \geq e > h} d_e,$$

with $m = l + 1$, $c_i = (\alpha + p - j - i + 2)$, $d_i = (\alpha + p - i + 1)$ for $1 \leq i \leq l$ and $c_{l+1} = (p - j + 1)$, $d_{l+1} = p$. This proves the recursion in (3.2) for $k \geq 3$. On the other hand

$$\begin{aligned} A(2, l) &= \det \begin{bmatrix} 1 & (\alpha + p)_l p \\ -(\alpha + p)p & -(\alpha + p)_{l+1} (p)_2 \end{bmatrix} = (\alpha + p)_l p (\alpha + p - l + lp) = \\ &= \sum_{r=1}^l p(\alpha + p - r)_{l-r} (\alpha + p)_r p + (\alpha + p)_{l+1} p \end{aligned}$$

proving (3.2) for $k = 2$.

Note that in case $k \leq p + \alpha$ we have $(\alpha + p)_l > 0$ for $0 \leq l \leq k$, and $(p + \alpha)_l = (p + \alpha)^l + O((p + \alpha)^{l-1})$ hence the multiplier in the sum in (3.2) does not change the (positive) coefficient - nor its sign - of the highest order terms of $A(k - 1, r)$.

We are going to prove that viewing $A(k, l)$ as a polynomial of the variables p and $p + \alpha$ one has $\deg A(k, l) = k + l$. The proof goes by induction on k . For $k = 1$ and l arbitrary this is an immediate consequence of its definition. In fact - assuming the induction hypothesis for $k - 1$ and l arbitrary - we have that

$$\begin{aligned} \deg p(p + \alpha)_{l-r} A(k - 1, r) &= k - 1 + r + l - r + 1 = k + l \\ &\quad \text{for } 1 \leq r \leq l \leq k \leq p + \alpha, \text{ and} \\ \deg A(k - 1, l + 1) &= k + l, \end{aligned}$$

and using that there is no cancellation in the highest degree terms we obtain that $\deg A(k, l) = k + l$, hence we immediately get that $\deg_p M_p^{(\alpha)}(k) = k + 1$.

Computing the leading coefficient in p and $\alpha + p$ of $A(k, 1)$ leads to the following graph theoretical question: Let $G = ((\mathbb{Z}_{\geq 0})^2, \vec{E})$ be the following graph: there is a directed arrow from (a_1, b_1) pointing to (a_2, b_2) if and only if $a_2 = a_1 + 1$ and $b_2 \geq b_1 - 1$. We shall also use the word edge instead of arrow in case we are not interested in its direction. We will call an edge $(a, b_1) \rightarrow (a + 1, b_2)$ an upward edge if $b_2 \geq b_1$, if $b_2 = b_1 - 1$ we will refer to it as a downward edge. The height of an edge $((a, b_1), (a + 1, b_2))$ is going to be defined as $b_2 - b_1$, total height of a set of edges is the sum of their heights.

Let us call a path ending in (k, l) for $k \geq 1, l \geq 1$ legal if it starts in the origin and after that it stays strictly above the line $y = 0$. Since $\deg_p A(k, l) = k + l$, it can be written as $A(k, l) = \sum_{j=0}^{k+l} a_j^{(k,l)} p^j (\alpha + p)^{k+l-j} + \text{L.O.T.}$, for some $a_j^{(k,l)}$, $j = 0, \dots, k + l$, where L.O.T. means lower order terms. But the recursion (3.2) implies that the degree of p in $A(k, l)$ cannot be larger than k and it is at least 1, for any $l \geq 1$, thus $A(k, l) = \sum_{j=1}^k a_j^{(k,l)} p^j (\alpha + p)^{k+l-j} + \text{L.O.T.}$ Using the recursion (3.2) again we obtain that

$$a_j^{(k,l)} = \sum_{h=1}^l a_{j-1}^{(k-1,h)} + a_j^{(k-1,l+1)}.$$

Our claim is that $a_j^{(k,l)}$ is equal to the number of legal paths $b_j^{(k,l)}$ ending in (k, l) with exactly j upward edges.

For $k = 1, l \geq 1$ we have that $A(1, l) = p(\alpha + p)_l$ thus the highest order term is $p(\alpha + p)^l$ and so $a_1^{(1,l)} = 1$, while $a_j^{(1,l)} = 0$ for $j \neq 1$ obviously coinciding with the values $b_j^{(1,l)}$, $j \geq 0, l \geq 1$ since in this case the path consists of one single upward edge.

For the induction step $k - 1 \mapsto k$ consider the following: Each of the legal paths ending in (k, l) has to go through exactly one of the points $(k - 1, r)$

$1 \leq r \leq l + 1$. A path with exactly j upward edges going through the points $(k - 1, r)$ for $1 \leq r \leq l$ should have $j - 1$ upward edges before these points, while a path going through $(k - 1, l + 1)$ has j upward edges before this point. Therefore the number of legal paths ending in (k, l) and having j upward edges is the sum of the number of legal paths ending in $(k - 1, r)$ with $1 \leq r \leq l$ with $j - 1$ upward edges plus the number of legal paths ending in $(k - 1, l + 1)$ with j upward edges. In other words:

$$b_j^{(k,l)} = \sum_{r=1}^l b_{j-1}^{(k-1,r)} + b_j^{(k-1,l+1)}.$$

Thus the number of legal paths satisfies the same recursion as the coefficients in the sequence $A(k, l)$. Since for $k = 1$ they are equal the induction argument gives that $a_j^{(k,l)} = b_j^{(k,l)}$ for $j = 1, \dots, k$, $k \geq 1$, $l \geq 1$.

Now let us turn our attention to computing the coefficients of the highest order term of $M_p^{(\alpha)}(k) = A(k, 1) = \sum_{j=1}^k a_j^{(k,1)} p^j (\alpha + p)^{k-j+1} + \text{L.O.T.}$ As we proved before the coefficient $a_j^{(k,1)}$ is given by the number of legal paths ending in $(k, 1)$ with j upward edges. In this case there are $k - j$ downward edges with total height $-(k - j)$ hence the total height of the upward edges is $k - j + 1$. Since the length of the legal path from the origin to $(k, 1)$ is k there are $\binom{k}{j}$ possibilities to choose the positions of the j upward edges. On the other hand the total height of the upward edges is $k - j + 1$, and there are $\binom{k}{j-1}$ ways writing it as a sum of j non-negative numbers when the sequence of the summands matters. Choosing these numbers as the heights of the upward edges we obtain a path from the origin to $(k, 1)$ which is not necessarily legal, since they can cross the line $y = 0$. For such a given path let (x, y) denote the node of the path with the largest first coordinate such that its second coordinate is not greater than the second coordinate of any other node of the path (i.e. the latest "global minimum" of the path). By placing this node with the tail of the path in the origin this new path is a legal path ending in $(k - x, 1 + y)$. Taking the first part of the original path (connecting the origin with (x, y)) and gluing it to $(k - x, 1 + y)$ we will get a legal path ending in $(k, 1)$. We will say that two paths are equivalent if the cut-and-glue process described above results in the same legal path. The equivalence class of a path consists of its periodic horizontal translations, so in each equivalence class there are k paths. Since the cut-and-glue process gives the same legal path for each equivalence class, thus the number of legal paths ending in $(k, 1)$ having j upward edges is given by $\frac{1}{k} \binom{k}{j} \binom{k}{j-1}$, hence

$$M_p^{(\alpha)}(k) = \sum_{j=1}^k \frac{1}{k} \binom{k}{j} \binom{k}{j-1} p^j (\alpha + p)^{k-j+1} + \text{L.O.T}$$

and so Theorem 3.1 is proved. ■

Remark 3.1. If $\alpha/p = c$ with $c \in (-1, \infty)$ and $k < \alpha + p + 1$ then

$$(3.3) \quad \sum_{l=1}^p (\xi_{p,l}^{(\alpha)})^k = \sum_{j=1}^k \frac{1}{k} \binom{k}{j} \binom{k}{j-1} (c+1)^{k-j+1} p^{k+1} + f(\alpha + p, p)$$

hence we immediately get that

$$\int x^k dm_p^{(\alpha_p)}(x) \xrightarrow{p \rightarrow \infty} \int x^k d\mu_c(x)$$

if $\frac{\alpha_p}{p} \rightarrow c$ for all $k \geq 0$.

We also emphasize that even in the case when $\alpha < 0$ is not an integer thus $L_p^{(\alpha)}(z)$ has complex roots with nonzero imaginary part, the limit relation above holds true. But since now the measure is not concentrated on the real line this property is not enough for the identification of the the limit measure.

References

- [1] **Forrester, P. J. and A. Gamburd**, Counting formulas associated with some random matrix averages, *Journal of Combinatorial Theory, Series A*, **113(6)** (2006), 934–951.
- [2] **Marchenko, V. A. and L. A. Pastur**, Distribution of eigenvalues for some sets of random matrices, *Matematicheskii Sbornik*, **114(4)** (1967), 507–536.
- [3] **Martínez-Finkelshtein, A., P. Martínez-González and R. Orive**, On asymptotic zero distribution of Laguerre and generalized Bessel polynomials with varying parameters, *Journal of Computational and Applied Mathematics*, **113(1)** (2001), 477–487.
- [4] **Szegő, G.**, *Orthogonal Polynomials*, American Mathematical Society, Colloquium Publications, Voluma XXIII, 1939.

M. Kornyik and Gy. Michaletzky

Department of Probability Theory and Statistics

Eötvös Loránd University

1117 Budapest

Pázmány Péter sétány 1/A

Hungary

koma@cs.elte.hu

michaletzky@caesar.elte.hu

