

SEGRE'S UPPER BOUND FOR THE REGULARITY INDEX OF $2n + 2$ NON-DEGENERATE DOUBLE POINTS IN \mathbb{P}^n

Tran Nam Sinh and Phan Van Thien

(Hue City, Viet Nam)

Communicated by Bui Minh Phong

(Received May 20, 2017; accepted August 6, 2017)

Abstract. We prove the Segre's upper bound for the regularity index of $2n + 2$ non-degenerate double points that do not exist $n+1$ points lying on a $(n - 2)$ -plane in \mathbb{P}^n .

1. Introduction

Let P_1, \dots, P_s be a set of distinct points in a projective space with n -dimension $\mathbb{P}^n := \mathbb{P}_k^n$, with k as an algebraically closed field. Let \wp_1, \dots, \wp_s be the homogeneous prime ideals of the polynomial ring $R := k[x_0, \dots, x_n]$ corresponding to the points P_1, \dots, P_s . Let m_1, \dots, m_s be positive integers and $I = \wp_1^{m_1} \cap \dots \cap \wp_s^{m_s}$. Denote $Z = m_1 P_1 + \dots + m_s P_s$ the zero-scheme defined by I , and we call Z a set of s fat points in \mathbb{P}^n .

The homogeneous coordinate ring of Z is

$$A = R/(\wp_1^{m_1} \cap \dots \cap \wp_s^{m_s}).$$

The ring $A = \bigoplus_{t \geq 0} A_t$ is a one-dimension Cohen-Macaulay k -graded algebra whose multiplicity is $e(A) = \sum_{i=1}^s \binom{m_i+n-1}{n}$. The Hilbert function $H_A(t) =$

Key words and phrases: Fat points, regularity index, zero-scheme.

2010 Mathematics Subject Classification: Primary 14C20, Secondary 13D40.

<https://doi.org/10.71352/ac.46.327>

$= \dim_k A_t$ increases strictly until it reaches the multiplicity $e(A)$, at which it stabilizes. The regularity index of Z is defined to be the least integer t such that $H_A(t) = e(A)$, and we denote it by $\text{reg}(Z)$ (or $\text{reg}(A)$).

In 1961, Segre (see [10]) showed the upper bound for regularity index of generic fat points $Z = m_1P_1 + \cdots + m_sP_s$ in \mathbb{P}^2 :

$$\text{reg}(Z) \leq \max \left\{ m_1 + m_2 - 1, \left\lceil \frac{m_1 + \cdots + m_s}{2} \right\rceil \right\}$$

with $m_1 \geq \cdots \geq m_s$.

For arbitrary fat points $Z = m_1P_1 + \cdots + m_sP_s$ in \mathbb{P}^2 , in 1969 Fulton (see [9]) gave the following upper bound:

$$\text{reg}(Z) \leq m_1 + \cdots + m_s - 1.$$

This bound was later extended to arbitrary fat points in \mathbb{P}^n by Davis and Geramita (see [6]). They also showed that this bound is attained if and only if points P_1, \dots, P_s lie on a line in \mathbb{P}^n .

A set of fat points $Z = m_1P_1 + \cdots + m_sP_s$ in \mathbb{P}^n is said to be in general position if no $j + 2$ of the points P_1, \dots, P_s are on any j -plane for $j < n$. A set of fat points $Z = m_1P_1 + \cdots + m_sP_s$ of \mathbb{P}^n is said to be non-degenerate if all points P_1, \dots, P_s do not lie on a hyperplane of \mathbb{P}^n . In 1991, Catalisano (see [3], [4]) extended Segre's result to fat points in general position in \mathbb{P}^2 , and later Catalisano, Trung and Valla (see [5]) extended the result to fat points in general position in \mathbb{P}^n , they proved:

$$\text{reg}(Z) \leq \max \left\{ m_1 + m_2 - 1, \left\lceil \frac{m_1 + \cdots + m_s + n - 2}{n} \right\rceil \right\}.$$

In 1996, N.V. Trung gave the following conjecture: Let $Z = m_1P_1 + \cdots + m_sP_s$ be arbitrary fat points in \mathbb{P}^n . Then

$$\text{reg}(Z) \leq \max \left\{ T_j \mid j = 1, \dots, n \right\},$$

where

$$T_j = \max \left\{ \left\lceil \frac{\sum_{i=1}^q m_{i_l} + j - 2}{j} \right\rceil \mid P_{i_1}, \dots, P_{i_q} \text{ lie on a } j\text{-plane} \right\}.$$

This upper bound nowadays is called the Segre's upper bound.

The Segre's upper bound is proved right in projective spaces with $n = 2$, $n = 3$ (see [12], [13]), for the case of double points $Z = 2P_1 + \cdots + 2P_s$ in \mathbb{P}^n with $n = 4$ (see [14]) by Thien; also for case $n = 2, n = 3$, independently by Fatabbi and Lorenzini (see [7], [8]).

In 2012, Benedetti, Fatabbi and Lorenzini proved the Segre's bound for any set of $n + 2$ non-degenerate fat points $Z = m_1P_1 + \cdots + m_{n+2}P_{n+2}$ of \mathbb{P}^n (see [1]), and independently Thien also proved the Segre's bound for a set of $s + 2$ fat points which is not on a $(s - 1)$ -space in \mathbb{P}^n , $s \leq n$ (see [15]).

Recently, Ballico, Dumitrescu and Postinghen proved the Segre's upper bound for the case $n + 3$ non-degenerate fat points $Z = m_1P_1 + \cdots + m_{n+3}P_{n+3}$ in \mathbb{P}^n (see [2]) and Sinh proved the Segre's upper bound for the regularity index of $2n + 1$ double points $Z = 2P_1 + \cdots + 2P_{2n+1}$ that do not exist $n + 1$ points lying on a $(n - 2)$ -plane in \mathbb{P}^n (see [11]). Up to now, there have not been any other result of Trung's conjecture published yet.

In this article, we prove the Segre's upper bound in the case $2n + 2$ non-degenerate double points $Z = 2P_1 + \cdots + 2P_{2n+2}$ that do not exist $n + 1$ points lying on a $(n - 2)$ -plane in \mathbb{P}^n .

2. Preliminaries

We will use the following lemmas which have been proved. The first lemma allows us to compute the regularity index by induction.

Lemma 2.1. [5, Lemma 1]. *Let P_1, \dots, P_r, P be distinct points in \mathbb{P}^n , and let \wp be the defining ideal of P . If m_1, \dots, m_r and a are positive integers, $J = \wp_1^{m_1} \cap \cdots \cap \wp_r^{m_r}$, and $I = J \cap \wp^a$, then*

$$\operatorname{reg}(R/I) = \max \left\{ a - 1, \operatorname{reg}(R/J), \operatorname{reg}(R/(J + \wp^a)) \right\}.$$

To compute $\operatorname{reg}(R/(J + \wp^a))$, we need the following lemma.

Lemma 2.2. [5, Lemma 3]. *Let P_1, \dots, P_r be distinct points in \mathbb{P}^n and a, m_1, \dots, m_r positive integers. Put $J = \wp_1^{m_1} \cap \cdots \cap \wp_r^{m_r}$ and $\wp = (x_1, \dots, x_n)$. Then*

$$\operatorname{reg}(R/(J + \wp^a)) \leq b$$

if and only if $x_0^{b-i}M \in J + \wp^{i+1}$ for every monomial M of degree i in x_1, \dots, x_n , $i = 0, \dots, a - 1$.

To find such a number b , we will find t hyperplanes L_1, \dots, L_t avoiding P such that $L_1 \cdots L_t M \in J$. For $j = 1, \dots, t$, since we can write $L_j = x_0 + G_j$ for some linear form $G_j \in \wp$, we get $x_0^t M \in J + \wp^{t+1}$. Therefore, if we put

$$\delta = \max \left\{ t + i \mid M \text{ is a monomial of degree } i, 0 \leq i \leq a - 1 \right\}$$

then

$$\operatorname{reg}(R/(J + \wp^a)) \leq \delta.$$

The hyperplanes L_1, \dots, L_t will be constructed by the help of the following lemma.

Lemma 2.3. [5, Lemma 4]. *Let P_1, \dots, P_r, P be distinct points in general position in \mathbb{P}^n , let $m_1 \geq \dots \geq m_r$ be positive integers, and let $J = \wp_1^{m_1} \cap \dots \cap \wp_r^{m_r}$. If t is an integer such that $nt \geq \sum_{i=1}^r m_i$ and $t \geq m_1$, we can find t hyperplanes, say L_1, \dots, L_t avoiding P such that for every $P_l, l = 1, \dots, r$, there exist m_l hyperplanes of $\{L_1, \dots, L_t\}$ passing through P_l .*

The two following lemmas are used to prove main results by induction.

Lemma 2.4. [11, Proposition 2.1]. *Let $X = \{P_1, \dots, P_{2n+1}\}$ be a set of $2n+1$ distinct points that do not exist $n+1$ points of X lying on a $(n-2)$ -plane in \mathbb{P}^n . Let \wp_i be the homogeneous prime ideal corresponding $P_i, i = 1, \dots, 2n+1$. Let*

$$Z = 2P_1 + \dots + 2P_{2n+1}.$$

Put

$$T_j = \max\left\{\left\lfloor \frac{1}{j}(2q+j-2) \right\rfloor \mid P_{i_1}, \dots, P_{i_q} \text{ lie on a } j\text{-plane}\right\},$$

$$T_Z = \max\{T_j \mid j = 1, \dots, n\}.$$

Then, there exists a point $P_{i_0} \in X$ such that

$$\operatorname{reg}(R/(J + \wp_{i_0}^2)) \leq T_Z,$$

where

$$J = \bigcap_{k \neq i_0} \wp_k^2.$$

Lemma 2.5. [11, Proposition 2.2]. *Let $X = \{P_1, \dots, P_{2n+1}\}$ be a set of $2n+1$ distinct points which do not exist $n+1$ points of X lying on a $(n-2)$ -plane in \mathbb{P}^n . Let $Y = \{P_{i_1}, \dots, P_{i_s}\}, 2 \leq s \leq 2n$, be a subset of X . Let \wp_i be the homogeneous prime ideal corresponding $P_i, i = 1, \dots, 2n+1$. Let*

$$Z = 2P_1 + \dots + 2P_{2n+1}.$$

Put

$$T_j = \max\left\{\left\lfloor \frac{1}{j}(2q+j-2) \right\rfloor \mid P_{i_1}, \dots, P_{i_q} \text{ lie on a } j\text{-plane}\right\},$$

$$T_Z = \max\{T_j \mid j = 1, \dots, n\}.$$

Then, there exists a point $P_{i_0} \in Y$ such that

$$\operatorname{reg}(R/(J + \wp_{i_0}^2)) \leq T_Z,$$

where

$$J = \bigcap_{P_k \in Y \setminus \{P_{i_0}\}} \wp_k^2.$$

3. Segre's upper bound for the regularity index of $2n + 2$ non-degenerate double points in \mathbb{P}^n

From now on, we consider a hyperplane and its identical defining linear form. These following propositions are important for proving of Segre' upper bound.

Proposition 3.1. *Let $X = \{P_1, \dots, P_{2n+2}\}$ be a non-degenerate set of $2n + 2$ distinct points that do not exist $n + 1$ points of X lying on a $(n - 2)$ -plane in \mathbb{P}^n . Let \wp_i be the homogeneous prime ideal corresponding $P_i, i = 1, \dots, 2n + 2$, and*

$$Z = 2P_1 + \dots + 2P_{2n+2}.$$

Put

$$T_j = \max \left\{ \left\lceil \frac{1}{j}(2q + j - 2) \right\rceil \mid P_{i_1}, \dots, P_{i_q} \text{ lie on a } j\text{-plane} \right\},$$

$$T_Z = \max\{T_j \mid j = 1, \dots, n\}.$$

Then, there exists a point $P_{i_0} \in X$ such that

$$\operatorname{reg}(R/(J + \wp_{i_0}^2)) \leq T_Z,$$

where

$$J = \bigcap_{k \neq i_0} \wp_k^2.$$

Proof. We denote $|H|$ by the number points of X lying on a j -plane H . The proposition was proved in projective spaces with $n \leq 4$ (see [7], [8], [12]–[14]). Thus, we will prove the case with $n \geq 5$.

We can see that there are $(n - 1)$ -planes H_1, \dots, H_d in \mathbb{P}^n with d as the least integer such that the two following conditions satisfied:

- (i) $X \subset \bigcup_{i=1}^d H_i$,
- (ii) $|H_i \cap (X) \setminus \bigcup_{j=1}^{i-1} H_j| = \max\{|H \cap (X \setminus \bigcup_{j=1}^{i-1} H_j)| \mid H \text{ is an } (n - 1)\text{-plane}\}.$

Since X non-degenerate and $n + 1$ points do not lie on a $(n - 2)$ -plane, $2 \leq d \leq 3$. We consider the following cases:

Case 1. $d = 3$. Since a hyperplane always passes through at least n points of X and $d = 3$, we have the two following cases:

- (i) $|H_1| = n$, $|H_2| = n$, $|H_3| = 2$.
- (ii) $|H_1 = n + 1| = |H_2 \setminus H_1| = n$, $|H_3| = 1$.

Case 1.1. $|H_1| = n$, $|H_2| = n$, $|H_3| = 2$. Since $|H_1| = n$, there do not exist $n + 1$ points of X lying on a hyperplane. Therefore, X is general position. By Lemma 2.3 and Lemma 2.2 we have

$$\text{reg}(R/(J + \wp_{i_0}^2)) \leq T_Z.$$

Case 1.2. $|H_1| = n + 1$, $|H_2| = n$, $|H_3| = 1$. We may assume that $P_1 \in H_3$. Choose $P_1 = P_{i_0} = (1, 0, \dots, 0)$, then $\wp_{i_0} = (x_1, \dots, x_n)$. Clearly, H_1, H_2 avoiding P_{i_0} . We have $H_1 H_1 H_2 H_2 \in J$ for every monomial $M = x_1^{c_1} \cdots x_n^{c_n}$, $c_1 + \cdots + c_n = i$, $i = 0, 1$. By Lemma 2.2 we have

$$\text{reg}(R/(J + \wp_{i_0}^2)) \leq 4 + i \leq 5 \leq T_Z.$$

Case 2. $d = 2$. We have $X \subset H_1 \cup H_2$. Therefore, $|H_1| \geq n + 1$ and $H_1 \geq |H_2|$. We call q the number points of X lying on $H_2 \setminus H_1$, we have $1 \leq q \leq n + 1$, without loss of generality, we assume $P_1, \dots, P_q \in H_2 \setminus H_1$. Put $Y = \{P_1, \dots, P_q\}$. Since $n + 1$ points of X do not lie on a $(n - 2)$ -plane, Y does not lie on a $(q - 3)$ -plane. We consider the following cases:

Case 2.1. Y lies on a $(q - 1)$ -plane and Y does not lie on a $(q - 2)$ -plane. Choose $P_q = P_{i_0} = (1, 0, \dots, 0)$, $P_1 = (0, \underbrace{1}_2, \dots, 0), \dots, P_{q-1} = (0, \dots, \underbrace{1}_q, \dots, 0)$,

then $\wp_{i_0} = (x_1, \dots, x_n)$. Since we always have a $(q - 2)$ -plane, say K , passing through P_1, \dots, P_{q-1} and avoiding P_{i_0} ; therefore, we always have a hyperplane, say L , containing K and avoiding P_{i_0} . We have $H_1 H_1 L L \in J$. Thus $H_1 H_1 L L M \in J$ for every monomial $M = x_1^{c_1} \cdots x_n^{c_n}$, $c_1 + \cdots + c_n = i$, $i = 0, 1$. By Lemma 2.2 we have

$$\text{reg}(R/(J + \wp_{i_0}^2)) \leq 4 + i \leq 5 \leq T_Z.$$

Case 2.2. Y lies on a $(q - 2)$ -plane α , $q \geq 3$. We consider the following cases of Y :

Case 2.2.1. There are $q - 1$ points of Y lying on a $(q - 3)$ -plane. Assume that P_1, \dots, P_{q-1} lying on a $(q - 3)$ -plane, say K and $P_q \notin K$. Choose $P_q = P_{i_0} = (1, 0, \dots, 0)$, then $\wp_{i_0} = (x_1, \dots, x_n)$. Since $q \leq n + 1$, we have $q - 3 \leq n - 2$ and $P_{i_0} \notin K$, we always have a hyperplane L containing K and avoiding

P_{i_0} . We have $H_1H_1LL \in J$, thus $H_1H_1LLM \in J$ for every monomial $M = x_1^{c_1} \cdots x_n^{c_n}$, $c_1 + \cdots + c_n = i$, $i = 0, 1$. By Lemma 2.2 we have

$$\operatorname{reg}(R/(J + \wp_{i_0}^2)) \leq 4 + i \leq 5 \leq T_Z.$$

Case 2.2.2. There are not $q - 1$ points of Y lying on a $(q - 3)$ -plane. We consider the three following cases of q :

Case 2.2.2.1. $q \geq 5$. Since any $(q - 3)$ -planes only pass through $q - 2$ points of Y . Choose $P_q = P_{i_0} = (1, 0, \dots, 0)$, $P_1 = (0, \underbrace{1}_2, 0, \dots, 0), \dots, P_{q-2} = (0, \dots, 0, \underbrace{1}_{q-1}, 0, \dots, 0)$. Put $m_l = 2 - i + c_l$, $l = 1, \dots, q - 2$, $m_{q-1} = 2$ and

$$t = \max \left\{ 2, \left[\left(\sum_{i=1}^{q-1} m_l + (q - 2) - 1 \right) / (q - 2) \right] \right\}.$$

We have

$$\begin{aligned} t + i &= \max \left\{ 2, \left[\left(\sum_{i=1}^{q-1} m_l + q - 3 \right) / (q - 2) \right] \right\} + i \leq \\ &\leq \max \left\{ 2 + i, \left[\left(\sum_{i=1}^{q-1} m_l + (q - 2)i + q - 3 \right) / (q - 2) \right] \right\} \leq \\ &\leq \max \left\{ 2 + i, \left[(3q - 4) / (q - 2) \right] \right\} \leq 3. \end{aligned}$$

Therefore,

$$t \leq 3 - i.$$

By Lemma 2.2, we can find t $(q - 3)$ -planes, say G_1, \dots, G_t avoiding P_{i_0} such that for every point P_l , $l = 1, \dots, q - 1$, there are m_l $(q - 3)$ -planes of G_1, \dots, G_t passing through P_l . With $j = 1, \dots, t$ we find a hyperplane L_j containing G_j and avoiding P_{i_0} . Therefore

$$L_1 \cdots L_t \in \wp_1^{m_1} \cap \cdots \cap \wp_{q-2}^{m_{q-2}} \cap \wp_{q-1}^2.$$

Moreover, since $H_1H_1 \in \wp_{q+1}^2 \cap \cdots \cap \wp_{2n+2}^2$ and $M \in \wp_1^{i-c_1} \cap \cdots \cap \wp_{q-2}^{i-c_{q-2}}$, then

$$H_1H_1L_1 \cdots L_tM \in J.$$

By Lemma 2.2 we have

$$\operatorname{reg}(R/(J + \wp_{i_0}^2)) \leq 2 + (3 - i) + i \leq T_Z.$$

Case 2.2.2.2. $q = 4$. We have $P_1, P_2, P_3, P_4 \notin H_1$. Choose $P_1 = P_{i_0} = (1, 0, \dots, 0)$, $P_3 = (0, \underbrace{1}_2, 0, \dots, 0)$, $P_4 = (0, 0, \underbrace{1}_3, 0, \dots, 0)$, \dots , $P_{n+1} = (0, \dots, 0, \underbrace{1}_n, 0)$, $P_{n+2} = (0, \dots, 0, \underbrace{1}_{n+1})$, therefore $\wp_{i_0} = (x_1, \dots, x_n)$. We

call l_1 a line passing through P_2, P_3 ; l_2 a line passing through P_3, P_4 ; l_3 a line passing through P_2, P_4 . We consider the two following cases of i :

a) $i = 0$. With $j = 1, 2, 3$, since $P_{i_0} \notin l_j$, then we always have a hyperplane L_j containing l_j and avoiding P_{i_0} . We have $H_1 H_1 L_1 L_2 L_3 \in J$, thus $H_1 H_1 L_1 L_2 L_3 M \in J$. By Lemma 2.2 we have

$$\text{reg}(R/(J + \wp_{i_0}^2)) \leq 5 \leq T_Z.$$

b) $i = 1$. Since $c_1 + \dots + c_n = 1$, then there exists $j \in \{1, \dots, n\}$ such that $c_j = 1, c_k = 0, k \in \{1, \dots, n\} \setminus \{j\}$.

◦ If $j \in \{1, 2\}$, assume that $c_1 = 1$ then

$$M \in \wp_4 \cap \wp_5 \cap \dots \cap \wp_{n+2}.$$

We have a $(n-2)$ -plane, say K_1 passing through P_{n+3}, \dots, P_{2n-1} and l_1 , a $(n-2)$ -plane, say K_2 passing through P_{2n}, P_{2n+1} and l_1 , a $(n-2)$ -plane, say K_3 passing through P_4, P_{2n+2} avoiding P_{i_0} . With $i = 1, 2, 3$, we always have hyperplanes L_i containing K_i and avoiding P_{i_0} . We have

$$H_1 L_1 L_2 L_3 \in \wp_2^2 \cap \wp_3^2 \cap \wp_4 \cap \wp_5 \cap \dots \cap \wp_{n+2} \cap \wp_{n+3}^2 \cap \dots \cap \wp_{2n+2}^2.$$

Therefore

$$H_1 L_1 L_2 L_3 M \in J.$$

By Lemma 2.2 we have

$$\text{reg}(R/(J + \wp_{i_0}^2)) \leq 4 + i \leq T_Z.$$

◦ If $j \in \{3, \dots, n\}$, assume that $c_3 = 1$ then

$$M \in \wp_3 \cap \wp_4 \cap \wp_6 \cap \dots \cap \wp_{n+2}.$$

We call l_1 a line passing through P_2, P_3 and l_2 a line passing through P_2, P_4 . With $i = 1, 2$, since $P_{i_0} \notin l_i$, then we always have hyperplanes L_i containing l_i and avoiding P_{i_0} . We have

$$L_1 L_2 \in \wp_2^2 \cap \wp_3 \cap \wp_4$$

Since $H_1 H_1 \in \wp_5^2 \cap \dots \cap \wp_{2n+2}^2$ then

$$H_1 H_1 L_1 L_2 M \in J.$$

By Lemma 2.2 we have

$$\text{reg}(R/(J + \wp_{i_0}^2)) \leq 4 + i \leq T_Z.$$

Case 2.2.2.3. $q = 3$. We have $P_1, P_2, P_3 \notin H_1$. We call l a line passing through P_1, P_2, P_3 and $W = \{P_4, \dots, P_{2n+2}\}$ are the points of X lying on $H_1 \cap X$, then there are $(n-2)$ -planes Q_1, \dots, Q_r in \mathbb{P}^n such that the two following conditions satisfied:

- (i) $W \subset \bigcup_{i=1}^r Q_i$,
- (ii) $|Q_i \cap (W \setminus \bigcup_{j=1}^{i-1} Q_j)| = \max\{|Q \cap (W \setminus \bigcup_{j=1}^{i-1} Q_j)| \mid Q \text{ is a } (n-2)\text{-plane}\}$.

Since $n+1$ of X do not lie on a $(n-2)$ -plane, then we consider the two following cases of Q_1 :

a) $|Q_1| = n$. We have $r = 2$ and $|Q_2| = n-1$. Put $U = \{P_4, \dots, P_{n+2}\}$ to be $n-1$ points lying on Q_2 v $T = \{P_1, \dots, P_{n+2}\}$. We consider the two following cases of T :

a.1) T does not lie on a $(n-1)$ -plane. Since P_1, P_2, P_3 lie on a line l , then we always have a hyperplane containing l and passing through $n-2$ points of U . Assume that L to be a hyperplane containing l and passing through points P_4, \dots, P_{n+1} . Clearly, the hyperplane L avoiding P_{n+2} (if not, then T lies on a $(n-1)$ -plane). Choose $P_{n+2} = P_{i_0} = (1, 0, \dots, 0)$, then $\wp_{i_0} = (x_1, \dots, x_n)$. Since $P_{i_0} \notin Q_1$, therefore we always have a hyperplane L_1 containing Q_1 and avoiding P_{i_0} . We have $LLL_1 L_1 \in J$ then $LLL_1 L_1 M \in J$ for every monomial $M = x_1^{c_1} \cdots x_n^{c_n}, c_1 + \dots + c_n = i, i = 0, 1$. By Lemma 2.2 we have

$$\text{reg}(R/(J + \wp_{i_0}^2)) \leq 4 + i \leq 5 \leq T_Z.$$

a.2) T lies on a $(n-1)$ -plane, say H . Assume that $|Q_1 \cap H \cap X| = s$. When hyperplane H passing through $n+2+s$ points of X . Consider $n-s$ points lying on $Q_1 \setminus H$, say $P_{i_1}, \dots, P_{i_{n-s}} \in Q_1 \setminus H$.

a.2.1) Case $P_{i_1}, \dots, P_{i_{n-s}}$ lie on a $(n-s-1)$ -plane and they do not lie on a $(n-s-2)$ -plane. Choose $P_{i_1} = P_{i_0} = (1, 0, \dots, 0)$, then $\wp_{i_0} = (x_1, \dots, x_n)$. Since we always have a $(n-s-2)$ -plane, say β passing through $P_{i_2}, \dots, P_{i_{n-s-1}}$. Moreover, since $n-s-2 \leq n-2$ then we always have a hyperplane L containing β and avoiding P_{i_0} . We have $HHLL \in J$ then $HHLLM \in J$ for every monomial $M = x_1^{c_1} \cdots x_n^{c_n}, c_1 + \dots + c_n = i, i = 0, 1$. By Lemma 2.2 we have

$$\text{reg}(R/(J + \wp_{i_0}^2)) \leq 4 + i \leq 5 \leq T_Z.$$

a.2.2) Case $P_{i_1}, \dots, P_{i_{n-s}}$ lie on a $(n-s-2)$ -plane. Since P_1, P_2, P_3 lie on a line, then $P_1, P_2, P_3, P_{i_1}, \dots, P_{i_{n-s}}$ lie on a $(n-s)$ -plane. So, $n-1 \leq n-s \leq n$ or $0 \leq s \leq 1$.

• If $\{P_{i_1}, \dots, P_{i_{n-s}}\}$ has $n-s-1$ points lying on a $(n-s-3)$ -plane, say γ . Assume that $P_{i_1} \notin \gamma$, then choose $P_{i_1} = P_{i_0} = (1, 0, \dots, 0)$, then $\wp_{i_0} = (x_1, \dots, x_n)$. Since $P_{i_0} \notin \gamma$ therefore we always have a hyperplane L containing γ and avoiding P_{i_0} . We have $LLHH \in J$ then $LLHHM \in J$ for every

monomial $M = x_1^{c_1} \cdots x_n^{c_n}, c_1 + \cdots + c_n = i, i = 0, 1$. By Lemma 2.2 we have

$$\operatorname{reg}(R/(J + \wp_{i_0}^2)) \leq 4 + i \leq 5 \leq T_Z.$$

• If $\{P_{i_1}, \dots, P_{i_{n-s}}\}$ without $n-s-1$ points lying on a $(n-s-3)$ -plane, then any $(n-s-3)$ -plane only pass through $n-s-2$ points of $\{P_{i_1}, \dots, P_{i_{n-s}}\}$. Choose $P_{i_1} = P_{i_0} = (1, 0, \dots, 0)$, $P_{i_2} = (0, \underbrace{1}_2, 0, \dots, 0), \dots, P_{i_{n-s-1}} = (0, \dots, 0, \underbrace{1}_{n-s-1}, 0, \dots, 0)$ then $\wp_{i_0} = (x_1, \dots, x_n)$. Put $m_l = 2 - i + c_l, l = 2, \dots, n-s-1, m_{n-s} = 2$ and

$$t = \max \left\{ 2, \left[\left(\sum_{i=1}^{n-s-1} m_l + (n-s-2) - 1 \right) / (n-s-2) \right] \right\}.$$

We have

$$\begin{aligned} t + i &= \max \{ 2, \left[\left(\sum_{i=1}^{n-s-1} m_l + n-s-3 \right) / (n-s-2) \right] \} + i \leq \\ &\leq \max \{ 2 + i, \left[\left(\sum_{i=1}^{n-s-1} m_l + (n-s-2)i + n-s-3 \right) / (n-s-2) \right] \} \leq \\ &\leq \max \{ 2 + i, \left[(3(n-s-2) + 2) / (n-s-2) \right] \}. \end{aligned}$$

✓ $s = 0$ or $n \geq 6$, we have

$$t \leq 3 - i.$$

By Lemma 2.3 we can find t $(q-3)$ -planes, say G_1, \dots, G_t avoiding P_{i_0} such that for every point $P_l, l = 1, \dots, q-1$, there are m_l $(q-3)$ -planes of G_1, \dots, G_t passing through. With $j = 1, \dots, t$ we find a hyperplane L_j containing G_j and avoiding P_{i_0} . Therefore

$$L_1 \cdots L_t \in \wp_{i_2}^{m_2} \cap \cdots \cap \wp_{i_{n-s-1}}^{m_{n-s-1}} \cap \wp_{i_{n-s}}^2.$$

So, $HHL_1 \cdots L_t M \in J$ for every monomial $M = x_1^{c_1} \cdots x_n^{c_n}, c_1 + \cdots + c_n = i, i = 0, 1$. By Lemma 2.2 we have

$$\operatorname{reg}(R/(J + \wp_{i_0}^2)) \leq 4 + i \leq 5 \leq T_Z.$$

✓ $s = 1$ and $n = 5$. Then hyperplane H pass through eight points of X and there are four points $P_{i_1}, P_{i_2}, P_{i_3}, P_{i_4}$ lying on a 2-plane, say $\gamma_1 \setminus H$. According to Case 2.2.2.2 we have proved it.

b) If $|Q_1| = n-1$, then $W = \{P_4, \dots, P_{2n+2}\}$ lie on the general position in H_1 . We call H a hyperplane containing l and passing through $n-3+u$ points of $W \cap H_1$. We have $u \geq 1$.

• If $u = 1$, then consider $n+1$ points of $H_1 \setminus H$. Without loss of generality, assume that $P_{n+2}, \dots, P_{2n+2} \in H_1 \setminus H$. Put $V = \{P_{n+2}, \dots, P_{2n+2}\}$. Since there do not exist n points of V lying on a $(n-2)$ -plane. Choose $P_{n+2} = P_{i_0} =$

$= (1, 0, \dots, 0)$, $P_{n+3} = (0, \underbrace{1}_2, 0, \dots, 0), \dots, P_{2n+1} = (0, \dots, 0, \underbrace{1}_n, 0)$ then $\wp_{i_0} = (x_1, \dots, x_n)$. Put $m_l = 2 - i + c_l$, $l = n + 3, \dots, 2n + 1$, $m_{2n+2} = 2$ and

$$t = \max \left\{ 2, \left[\left(\sum_{i=n+3}^{2n+2} m_l + (n-1) - 1 \right) / (n-1) \right] \right\}.$$

We have

$$\begin{aligned} t + i &= \max \left\{ 2, \left[\left(\sum_{i=n+3}^{2n+2} m_l + n - 2 \right) / (n-1) \right] \right\} + i \leq \\ &\leq \max \left\{ 2 + i, \left[\left(\sum_{i=n+3}^{2n+2} m_l + (n-1)i + n - 2 \right) / (n-1) \right] \right\} \leq \\ &\leq \max \left\{ 2 + i, \left[(3n-1) / (n-1) \right] \right\} \leq 3. \end{aligned}$$

Therefore

$$t \leq 3 - i.$$

By Lemma 2.3 we can find t $(n-2)$ -planes G_1, \dots, G_t avoiding P_{i_0} such that for every $P_l, l = n + 3, \dots, 2n + 2$, there are m_l $(n-2)$ -planes of G_1, \dots, G_t passing through. With $j = 1, \dots, t$ we find a hyperplane L_j containing G_j and avoiding P_{i_0} . Therefore

$$L_1 \cdots L_t \in \wp_{n+3}^{m_{n+3}} \cap \cdots \cap \wp_{2n+1}^{m_{2n+1}} \cap \wp_{2n+2}^2.$$

Moreover, since $HH \in \wp_1^2 \cap \cdots \cap \wp_{n+1}^2$ and $M \in \wp_{n+3}^{i-c_1} \cap \cdots \cap \wp_{2n+1}^{i-c_{n-1}}$ then

$$H_1 H_1 L_1 \cdots L_t M \in J.$$

By Lemma 2.2 we have

$$\text{reg}(R/(J + \wp_{i_0}^2)) \leq 2 + (3 - i) + i \leq T_Z.$$

• If $u \geq 2$, then there are $n+2-u$ points, assume that $P_{i_1}, \dots, P_{n+2-u} \in H_1 \setminus H$. Since $u \geq 2$ then $n+2-u \leq n$. Moreover, since $P_{i_1}, \dots, P_{n+2-u}$ lie on the general position in H_1 , then we have a $(n-u)$ -plane, say π , passing through $n+1-u$ points $P_{i_2}, \dots, P_{n+2-u}$ and avoiding P_{i_1} . Choose $P_{i_1} = P_{i_0} = (1, 0, \dots, 0)$, then $\wp_{i_0} = (x_1, \dots, x_n)$. Since $P_{i_0} \notin \pi$, we always have a hyperplane, say L , containing π and avoiding P_{i_0} . We have $HHLL \in J$, therefore $HHLLM \in J$ for every monomial $M = x_1^{c_1} \cdots x_n^{c_n}, c_1 + \cdots + c_n = i, i = 0, 1$. By Lemma 2.2 we have

$$\text{reg}(R/(J + \wp_{i_0}^2)) \leq 4 + i \leq 5 \leq T_Z.$$

The proof of proposition 3.1 is completed. ■

From Lemma 2.4, Lemma 2.5 and Proposition 3.1, we get the following remark.

Remark 3.1. Let $X = \{P_1, \dots, P_{2n+2}\}$ be a non-degenerate set of $2n + 2$ distinct points that do not exist $n + 1$ points of X lying on a $(n - 2)$ -plane in \mathbb{P}^n . Let $Y = \{P_{i_1}, \dots, P_{i_s}\}, 2 \leq s \leq 2n + 1$, be a subset of X . Let \wp_i be the homogeneous prime ideal corresponding $P_i, i = 1, \dots, 2n + 1$, and

$$Z = 2P_1 + \dots + 2P_{2n+2}.$$

Put

$$T_j = \max \left\{ \left\lfloor \frac{1}{j} (2q + j - 2) \right\rfloor \mid P_{i_1}, \dots, P_{i_q} \text{ lie on a } j\text{-plane} \right\},$$

$$T_Z = \max \{T_j \mid j = 1, \dots, n\}.$$

Then, there exists a point $P_{i_0} \in Y$ such that

$$\text{reg}(R/(J + \wp_{i_0}^2)) \leq T_Z,$$

where

$$J = \bigcap_{P_k \in Y \setminus \{P_{i_0}\}} \wp_k^2.$$

The theorem below is the main result of this paper.

Theorem 3.2. *Let $X = \{P_1, \dots, P_{2n+2}\}$ be a non-degenerate set of $2n + 2$ distinct points that do not exist $n + 1$ points of X lying on a $(n - 2)$ -plane in \mathbb{P}^n . Let*

$$Z = 2P_1 + \dots + 2P_{2n+2}.$$

Then

$$\text{reg}(Z) \leq \max \{T_j \mid j = 1, \dots, n\} = T_Z,$$

where

$$T_j = \left\{ \left\lfloor \frac{2q + j - 2}{j} \right\rfloor \mid P_{i_1}, \dots, P_{i_q} \text{ lie on a } j\text{-plane} \right\}.$$

Proof. Firstly, we have the following claim:

Let $X = \{P_1, \dots, P_{2n+2}\}$ in \mathbb{P}^n , $Y = \{P_{i_1}, \dots, P_{i_s}\}$ be a subset of X , $1 \leq s \leq 2n + 1$. Then

$$\text{reg}(R/J_s) \leq T_Z,$$

where

$$J_s = \bigcap_{P_i \in Y} \wp_i^2.$$

We will prove this claim by induction on number points of Y .

If $s = 1$. Let \wp_1 be the defining homogeneous prime ideal of P_1 . Put $J_1 = \wp_1^2$, $A = R/J_1$. Then,

$$\text{reg}(R/J_1) = 1 \leq T_Z.$$

Assume that the claim is right for all subsets Y of X , whose number points are smaller or equal $s - 1$. Let $Y = \{P_{i_1}, \dots, P_{i_s}\}$. By Remark 3.1, there exists a point $P_{i_0} \in Y$ such that

$$(1) \quad \text{reg}(R/(J_{s-1} + \wp_{i_0}^2)) \leq T_Z,$$

where $J_{s-1} = \bigcap_{P_i \in Y \setminus \{P_{i_0}\}} \wp_i^2$. Note that, J_{s-1} is the intersection of ideals containing $s - 1$ double points of Y . By conjecture of induction, we have

$$(2) \quad \text{reg}(R/J_{s-1}) \leq T_Z.$$

By Lemma 2.1 we have

$$(3) \quad \text{reg}(R/J_s) = \left\{ 1, \text{reg}(R/(J_{s-1})), \text{reg}(R/(J_{s-1} + \wp_{i_0}^2)) \right\}.$$

From (1), (2) and (3) we have

$$\text{reg}(R/J_s) \leq T_Z.$$

The proof of the above claim is completed.

Now, we prove Theorem 3.2. Let $X = \{P_1, \dots, P_{2n+2}\}$ in \mathbb{P}^n , by Proposition 3.1, there exists a point $P_{i_0} \in X$ such that

$$(4) \quad \text{reg}(R/(J + \wp_{i_0}^2)) \leq T_Z.$$

where $J = \bigcap_{P_i \in X \setminus \{P_{i_0}\}} \wp_i^2$. Note that, J is the intersection of ideals containing $2n + 1$ double points of X . Therefore, by the above claim with $s = 2n + 1$, we have

$$(5) \quad \text{reg}(R/J) \leq T_Z.$$

By Lemma 2.1 we have

$$(6) \quad \text{reg } R/I = \left\{ 1, \text{reg}(R/J), \text{reg}(R/(J + \wp_{i_0}^2)) \right\}$$

where $I = J \cap \wp_{i_0}^2$.

From (4), (5) and (6) we have

$$\text{reg}(Z) \leq T_Z.$$

The proof of Theorem 3.2 is completed. ■

References

- [1] **Benedetti, B., G. Fatabbi and A. Lorenzini**, Segre's bound and the case of $n + 2$ fat points of \mathbb{P}^n , *Comm. Algebra*, **40**, (2012), 395–5473.
- [2] **Ballico, E., O. Dumitrescu and E. Postinghel**, On Segre's bound for fat points in \mathbb{P}^n , *J. Pure and Appl. Algebra*, **220**, (2016), 2307–2323.
- [3] **Catalisano, M.V.**, Linear systems of plane curves through fixed fat points of \mathbb{P}^2 , *J. Algebra*, **142(1)** (1991), 81–100.
- [4] **Catalisano, M.V.**, Fat points on a conic, *Comm. Algebra*, **19** (1991), 2153–2168.
- [5] **Catalisano, M.V., N.V. Trung and G. Valla**, A sharp bound for the regularity index of fat points in general position, *Proc. Amer. Math. Soc.*, **118** (1993), 717–724.
- [6] **Davis, E.D. and A.V. Geramita**, The Hilbert function of a special class of 1-dimension Cohen–Macaulay grade algebras, The Curves Seminar at Queen's, *Queen's Paper in Pure and Appl. Math.*, **67** (1984), 1–29.
- [7] **Fatabbi, G.**, Regularity index of fat points in the projective plane, *J. Algebra*, **170** (1994), 916–928.
- [8] **Fatabbi, G. and A. Lorenzini**, On the sharp bound for the regularity index of any set of fat points, *J. Pure Appl. Algebra*, **161** (2001), 91–111.
- [9] **Fulton, W.**, *Algebraic Curves*, Math. Lect. Note Series, Benjamin, New York, 1969.
- [10] **Serge, B.**, Alcune question su insiemi finiti di punti in geometria algebrica, *Atti. Convergnno. Intern. di Torino*, 1961, 15–33.
- [11] **Sinh, T.N.**, Segre's upper bound for the regularity index of $2n + 1$ double points in \mathbb{P}^n , (to appear in *Journal of Science, Hue University*).
- [12] **Thien, P.V.**, On Serge bound for the regularity index of fat points in \mathbb{P}^2 , *Acta Math. Vietnamica* **24** (1999), 75–81.
- [13] **Thien, P.V.**, Serge bound for the regularity index of fat points in \mathbb{P}^3 , *J. Pure and Appl. Algebra*, **151** (2000), 197–214.
- [14] **Thien, P.V.**, Sharp upper bound for the regularity of zero-schemes of double points in \mathbb{P}^4 , *Comm. Algebra*, **30** (2002), 5825–5847.
- [15] **Thien, P.V.**, Regularity index of $s + 2$ fat points not on a $(s - 1)$ -space, *Comm. Algebra*, **40** (2012), 3704–3715.
- [16] **Thien, P.V. and T.N. Sinh**, On the regularity index of s fat points not on a $(r - 1)$ -space, $s \leq r + 3$, *Comm. Algebra*, **45** (2017), 4123–4138.

T.N. Sinh and P.V. Thien

Department of Mathematics, College of Education

Hue University

34 Le Loi, Hue City, Vietnam

trannamsinh80@gmail.com, tphanvannl@yahoo.com