

A NEW CLASS OF UNIQUE RANGE SETS FOR MEROMORPHIC FUNCTIONS

Vu Hoai An and Pham Ngoc Hoa

(Hai Duong, Vietnam)

Communicated by Bui Minh Phong

(Received February 24, 2018; accepted May 30, 2018)

Abstract. In this paper, we give a new class of unique range sets for meromorphic functions. Note that this class different from Yi's [6], Frank-Reinders's [3] and Fujimoto's [4].

1. Introduction

In this paper, by a meromorphic function we mean a meromorphic function in the complex plane \mathbb{C} . We assume that the reader is familiar with the notations in the Nevanlinna theory (see [4], [5] and [8]). Let f be a non-constant meromorphic function on \mathbb{C} . For every $a \in \mathbb{C}$, define the function $\nu_f^a : \mathbb{C} \rightarrow \mathbb{N}$ by

$$\nu_f^a(z) = \begin{cases} 0 & \text{if } f(z) \neq a \\ m & \text{if } f(z) = a \text{ with multiplicity } m, \end{cases}$$

and set $\nu_f^\infty = \nu_{\frac{1}{f}}^0$. For $f \in \mathcal{M}(\mathbb{C})$ and $S \subset \mathbb{C} \cup \{\infty\}$, we define

$$E_f(S) = \bigcup_{a \in S} \{(z, \nu_f^a(z)) : z \in \mathbb{C}\}.$$

Key words and phrases: Meromorphic function, Unique range set.

2010 Mathematics Subject Classification: 30D35.

This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 101.01-2012.19.

<https://doi.org/10.71352/ac.47.109>

Two meromorphic functions f, g are said to *share* S , *counting multiplicity*, if $E_f(S) = E_g(S)$. Let a set $S \subset \mathbb{C} \cup \{\infty\}$ and f and g be two non-constant meromorphic (entire) functions. If $E_f(S) = E_g(S)$ implies $f = g$ for any two non-constant meromorphic (entire) functions f, g , then S is called a unique range set for meromorphic(entire) functions or, in brief, $URSM(URSE)$. Gross and Yang [2] showed that the set $S = \{z \in \mathbb{C} \mid z + e^z = 0\}$ is a $URSE$. Since then, $URSE$ and $URSM$ with finitely many elements have been found by Yi [6], Mues and Reinders [1], Frank and Reinders [3], Fujimoto [4]. In fact, examples of unique range sets given by most authors are sets of the form $\{z \in \mathbb{C} \mid z^n + az^m + b = 0\}$ under suitable conditions on the constants a and b and on the positive integers n and m (see [6]). So far, the smallest unique range set for meromorphic functions has 11 elements and was given by Frank and Reinders [3]. They proved the following result.

Theorem A. *The set*

$$\left\{ z \in \mathbb{C} \mid \frac{(n-1)(n-2)}{2} z^n + n(n-2) z^{n-1} + \frac{(n-1)n}{2} z^{n-2} + b = 0 \right\},$$

where $n \geq 11$ and $b \neq 0, 1$, is a unique range set for meromorphic functions.

Fujimoto [4] extended this result to zero sets of more general polynomials $P_F(z)$ satisfying the condition: for any zeros $e_i \neq e_j$ of $P_F(z)$ we have $P_F(e_i) \neq P_F(e_j)$.

In this paper, we give a new class of unique range sets for meromorphic functions. Note that this class is different from Yi's [6], Frank–Reinders's [3] and Fujimoto's [4] (see Theorem 2.1, Theorem 2.2).

2. A new class of unique range sets for meromorphic functions

We assume that the reader is familiar with the notations in the Nevanlinna theory (see [3], [4] and [8]).

We first need the following Lemmas.

Lemma 2.1. (See [8].) *Let f be a non-constant meromorphic function on \mathbb{C} and let a_1, a_2, \dots, a_q be distinct points of $\mathbb{C} \cup \{\infty\}$. Then*

$$(q-2)T(r, f) \leq \sum_{i=1}^q N_1(r, \frac{1}{f - a_i}) + S(r, f),$$

where $S(r, f) = o(T(r, f))$ for all r , except for a set of finite Lebesgue measure.

Lemma 2.2. (See [7].) *Let $d, n \in \mathbb{N}^*$, $d \geq n^2$, and let f_1, \dots, f_{n+1} be entire functions on \mathbb{C} , not identically zero and satisfying the condition $f_1^d + f_2^d + \dots + f_{n+1}^d = 0$. Then there is a decomposition of indices, $\{1, \dots, n+1\} = \cup I_v$, such that*

- i. *Every I_v contains at least 2 indices;*
- ii. *For $j, i \in I_v$; $f_i = c_{ij}f_j$, where c_{ij} is a non-zero constant.*

Now let us describe main result of the paper.

Let $d \in \mathbb{N}^*$, $d \geq 25$ and $a, b, c \in \mathbb{C}$, $a, b, c \neq 0$,

$$(A_1) \quad \text{with } c \neq \frac{b^d}{a^d}, \quad a^{2d} \neq 1, \quad c \neq a^d b^d, \quad c \neq \frac{(-1)^d b^d}{a^{2d}}, \quad c \neq (-1)^d b^d.$$

Then we consider following polynomial

$$(A_2) \quad P(z) = z^d + (az + b)^d + c, \text{ and let } P(z) \text{ has only simple zeros.}$$

We need following lemma.

Set $v_1 = (1, 0)$, $v_2 = (0, e)$ with $e^d = c$, $v_3 = (a, b)$. Define the set

$A := \{\alpha = (\alpha_1, \alpha_2)\}$, where α_1, α_2 are 2 distinct numbers of $\{1, 2, 3\}$. For each element $\alpha \in A$, we associate the matrix

$$A_\alpha = \begin{pmatrix} v_{\alpha_1} \\ v_{\alpha_2} \end{pmatrix}.$$

Main result of the paper is following theorem.

Theorem 2.1. *Let S be the set of zeros of the above polynomial $P(z)$. Assume that the conditions $(A_1), (A_2)$ are satisfied. Then S is a URSM.*

Proof. Write $f = \frac{f_1}{f_2}$ (resp., $g = \frac{g_1}{g_2}$), where f_1, f_2 (resp., g_1, g_2) are entire functions on \mathbb{C} having no common zeros. Set

$$Q(z_1, z_2) = z_1^d + (az_1 + bz_2)^d + e^d z_2^d, \text{ with } e^d = c$$

We consider following linear forms $L_i(z_1, z_2), i = 1, 2, 3$, on \mathbb{C}^2 :

$$L_1(z_1, z_2) = z_1, \quad L_2(z_1, z_2) = ez_2, \quad L_3(z_1, z_2) = az_1 + bz_2.$$

We first prove that if

$$Q(f_1, f_2) = Q(g_1, g_2), \text{ then } g_i = tf_i, i = 1, 2, \text{ where } t \in \mathbb{C}, t \neq 0,$$

and therefore $f = g$. From $Q(f_1, f_2) = Q(g_1, g_2)$ we have

$$(L_1(f_1, f_2))^d + (L_2(f_1, f_2))^d + (L_3(f_1, f_2))^d = (L_1(g_1, g_2))^d + (L_2(g_1, g_2))^d + (L_3(g_1, g_2))^d. \quad (2.1)$$

For simplicity, set $L_i(\tilde{f}) = L_i(f_1, f_2)$, $L_i(\tilde{g}) = L_i(g_1, g_2)$. Then from (2.1) we have

$$(L_1(\tilde{f}))^d + (L_2(\tilde{f}))^d + (L_3(\tilde{f}))^d = (L_1(\tilde{g}))^d + (L_2(\tilde{g}))^d + (L_3(\tilde{g}))^d. \quad (2.2)$$

We shall prove that for each $i = 1, 2, 3$, there exists a non-zero constant c_i such that $L_i(\tilde{f}) = c_i L_i(\tilde{g})$.

By non-constant of the functions f and g we give $L_i(\tilde{f}) \neq 0$, $L_i(\tilde{g}) \neq 0$. Since $d \geq 25$, from Lemma 2.2 it follows that for each $i = 1, 2, 3$, we have one of the following possibilities:

i/ there exists a $i' \in \{1, 2, 3\}$ with $i' \neq i$ such that

$$L_i(\tilde{f}) = b_{ii'} L_{i'}(\tilde{f}), b_{ii'} \neq 0. \quad (2.3)$$

ii/ there exists a $i' \in \{1, 2, 3\}$ such that

$$L_i(\tilde{f}) = c_{ii'} L_{i'}(\tilde{g}), c_{ii'} \neq 0. \quad (2.4)$$

iii/ there exist $i', i'' \in \{1, 2, 3\}$, $i' \neq i''$ such that

$$L_i(\tilde{f}) = c_{ii'} L_{i'}(\tilde{g}) = c_{ii''} L_{i''}(\tilde{g}), c_{ii'}, c_{ii''} \neq 0,$$

and then

$$L_{i'}(\tilde{g}) = c_{i'i''} L_{i''}(\tilde{g}), c_{i'i''} \neq 0. \quad (2.5)$$

If we have (2.3) or (2.5), we get a contradiction to the hypothesis of non-constant of the functions f and g . Thus, we have only possibility (2.4), i. e., for each $i = 1, 2, 3$, there exists an unique $\sigma(i) \in \{1, 2, 3\}$ with σ is a permutation of $\{1, 2, 3\}$ such that

$$L_i(\tilde{f}) = c_{\sigma(i)} L_{\sigma(i)}(\tilde{g}), \text{ this means that, } L_i(f_1, f_2) = c_{\sigma(i)} L_{\sigma(i)}(g_1, g_2), \quad (2.6)$$

where $c_{\sigma(i)}^d = 1$.

Set $\alpha = (1, 2)$, $\beta = (2, 3)$, and $\alpha' = (\sigma(1), \sigma(2))$, $\beta' = (\sigma(2), \sigma(3))$. Then

$$A_\alpha = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}, A_\beta = \begin{pmatrix} v_2 \\ v_3 \end{pmatrix}, \text{ and } \det A_\alpha = e, \det A_\beta = -ae. \quad (2.7)$$

Now we consider the following possibilities for (2.6):

Case 1. $\alpha' = (2, 1)$, $\beta' = (1, 3)$. Then

$$(2.8) \quad A_{\alpha'} = \begin{pmatrix} v_2 \\ v_1 \end{pmatrix}, \quad A_{\beta'} = \begin{pmatrix} v_1 \\ v_3 \end{pmatrix}, \quad \text{and} \quad \det A_{\alpha'} = -e, \quad \det A_{\beta'} = b.$$

From this and (2.6) we give

$$L_1(f_1, f_2) = c_2 L_2(g_1, g_2), \quad L_2(f_1, f_2) = c_1 L_1(g_1, g_2),$$

$$(2.9) \quad L_3(f_1, f_2) = c_3 L_3(g_1, g_2).$$

Then we get by (2.9)

$$(2.10) \quad A_{\alpha} f^t = B A_{\alpha'} g^t,$$

where

$$B = \begin{pmatrix} c_2 & 0 \\ 0 & c_1 \end{pmatrix},$$

and

$$(2.11) \quad A_{\beta} f^t = C A_{\beta'} g^t,$$

where

$$C = \begin{pmatrix} c_1 & 0 \\ 0 & c_3 \end{pmatrix}.$$

From the equations (2.10), (2.11) we get

$$(2.12) \quad f^t = A_{\alpha}^{-1} B A_{\alpha'} g^t, \quad f^t = A_{\beta}^{-1} C A_{\beta'} g^t.$$

By deleting f^t from the equations (2.12) we obtain $A_{\alpha}^{-1} B A_{\alpha'} g^t = A_{\beta}^{-1} C A_{\beta'} g^t$. By non-constant of g we have $A_{\alpha}^{-1} B A_{\alpha'} = A_{\beta}^{-1} C A_{\beta'}$. By $c_i^d = 1, i = 1, 2, 3$, and noting that

$$\det A_{\alpha} \det A_{\alpha}^{-1} = 1, \quad \det A_{\beta} \det A_{\beta}^{-1} = 1,$$

we obtain

$$(\det B)^d = 1, \quad (\det C)^d = 1,$$

$$\left(\frac{\det A_{\alpha}}{\det A_{\alpha'}} \right)^d = \left(\frac{\det A_{\beta}}{\det A_{\beta'}} \right)^d, \quad c = \frac{b^d}{a^d}.$$

a contradiction to the hypothesis $c \neq \frac{b^d}{a^d}$.

Case 2. $\alpha' = (3, 2)$, $\beta' = (2, 1)$. From this and (2.6) we give

$$\begin{aligned} L_1(f_1, f_2) &= c_3 L_3(g_1, g_2), \quad L_2(f_1, f_2) = c_2 L_2(g_1, g_2), \\ (2.13) \quad L_3(f_1, f_2) &= c_1 L_1(g_1, g_2). \end{aligned}$$

By the similar arguments as in **Case 1** we obtain a contradiction to the hypothesis $a^{2d} \neq 1$.

Case 3. $\alpha' = (3, 1)$, $\beta' = (1, 2)$. From this and (2.6) we give

$$\begin{aligned} L_1(f_1, f_2) &= c_3 L_3(g_1, g_2), \quad L_2(f_1, f_2) = c_1 L_1(g_1, g_2), \\ (2.14) \quad L_3(f_1, f_2) &= c_2 L_2(g_1, g_2). \end{aligned}$$

By the similar arguments as in **Case 1** we obtain a contradiction to the hypothesis $c \neq a^d b^d$.

Case 4. $\alpha' = (2, 3)$, $\beta' = (3, 1)$. From this and (2.6) we give

$$\begin{aligned} L_1(f_1, f_2) &= c_2 L_2(g_1, g_2), \quad L_2(f_1, f_2) = c_3 L_3(g_1, g_2), \\ (2.15) \quad L_3(f_1, f_2) &= c_1 L_1(g_1, g_2). \end{aligned}$$

By the similar arguments as in **Case 1** we obtain a contradiction to the hypothesis $c \neq \frac{(-1)^d b^d}{a^{2d}}$.

Case 5. $\alpha' = (1, 3)$, $\beta' = (3, 2)$. From this and (2.6) we give

$$\begin{aligned} L_1(f_1, f_2) &= c_1 L_1(g_1, g_2), \quad L_2(f_1, f_2) = c_3 L_3(g_1, g_2), \\ (2.16) \quad L_3(f_1, f_2) &= c_2 L_2(g_1, g_2). \end{aligned}$$

By the similar arguments as in **Case 1** we obtain a contradiction to the hypothesis $c \neq (-1)^d b^d$.

Case 6. $\alpha' = (1, 2)$, $\beta' = (2, 3)$. From this and (2.6) we give

$$\begin{aligned} L_1(f_1, f_2) &= c_1 L_1(g_1, g_2), \quad L_2(f_1, f_2) = c_2 L_2(g_1, g_2), \\ (2.17) \quad L_3(f_1, f_2) &= c_3 L_3(g_1, g_2). \end{aligned}$$

Since L_1, L_2 are linearly independent, L_1, L_2, L_3 are linearly dependent, there exist non-zero constants t_k such that

$$L_3 = \sum_{k=1}^2 t_k L_k, \quad \text{and} \quad L_3(\tilde{f}) = \sum_{k=1}^2 t_k L_k(\tilde{f}), \quad L_3(\tilde{g}) = \sum_{k=1}^2 t_k L_k(\tilde{g}),$$

$$L_k(\tilde{f}) = c_k L_k(\tilde{g}), k = 1, 2, L_3(\tilde{f}) = c_3 L_3(\tilde{g}).$$

Thus,

$$\sum_{k=1}^2 (c_3 - c_k) t_k L_k(\tilde{g}) = 0.$$

Since f_1, f_2 are linearly independent, it follows that all the c_i are equal each to other, say $c_i = t$. Then we have $g_i = t f_i$ for $i = 1, 2$. Therefore $f = g$.

Now we are going to complete the proof of Theorem 2.1. By $E_f(S) = E_g(S)$ it is easy to see that there exists an entire function h such that $Q(f_1, f_2) = e^h Q(g_1, g_2)$. Set $l = e^{\frac{h}{a}}$ and $G_1 = l g_1, G_2 = l g_2$. Then $Q(f_1, f_2) = Q(G_1, G_2)$. By the similar arguments as above we have $\frac{f_1}{f_2} = \frac{G_1}{G_2}$. Therefore $f = g$. Theorem 2.1 is proved. \blacksquare

A example of new class of unique range sets for meromorphic functions in Theorem 2.1 is following.

Theorem 2.2. *Let $d \in \mathbb{N}^*$, $d \geq 25$ and S be the set of zeros of polynomial $P(z) = z^d + (2z + 5)^d + 1$. Then S is a URSM.*

Proof. By $P(z) = z^d + (2z + 5)^d + 1$ we have $a = 2, b = 5, c = 1$. From this it follows that

$$a, b, c \neq 0, \text{ and } c \neq \frac{b^d}{a^d}, a^{2d} \neq 1, c \neq a^d b^d, c \neq \frac{(-1)^d b^d}{a^{2d}}, c \neq (-1)^d b^d.$$

So the condition (A_1) is satisfied. We shall prove that the condition (A_2) is satisfied. Take l is a any zero of $P'(z) = d(z^{d-1} + 2(2z + 5)^{d-1})$. Then

$$\begin{aligned} l^{d-1} + 2(2l + 5)^{d-1} &= 0, (2 + \frac{5}{l})^{d-1} = -\frac{1}{2}. \text{ Set } 2 + \frac{5}{l} = h. \text{ Then } h^{d-1} = -\frac{1}{2}, \\ l &= \frac{5}{h-2}, (2l + 5)^{d-1} = -\frac{1}{2} l^{d-1}, l^d + (2l + 5)^d + 1 = l^d - \frac{1}{2} l^{d-1} (2l + 5) + 1 \\ (2.18) \quad &= -\frac{5}{2} l^{d-1} + 1 = -\frac{5}{2} \frac{5^{d-1}}{(h-2)^{d-1}} + 1 = -\frac{5^d}{2(h-2)^{d-1}} + 1. \end{aligned}$$

Moreover

$$\begin{aligned} |h|^{d-1} &= \frac{1}{2}, |h| = (\frac{1}{2})^{\frac{1}{d-1}}, 0 < |h-2|^{d-1} \leq (|h| + 2)^{d-1}, \\ 0 < |h-2|^{d-1} &\leq ((\frac{1}{2})^{\frac{1}{d-1}} + 2)^{d-1} = \frac{(2.2^{\frac{1}{d-1}} + 1)^{d-1}}{2}, \\ 0 < 2.|h-2|^{d-1} &\leq (2.2^{\frac{1}{d-1}} + 1)^{d-1}, \end{aligned}$$

$$(2.19) \quad \frac{5^d}{2 \cdot |h-2|^{d-1}} \geq \frac{5^d}{(2 \cdot 2^{\frac{1}{d-1}} + 1)^{d-1}} > 1.$$

Combining (2.18) and (2.19) we get $-\frac{5^d}{2(h-2)^{d-1}} + 1 \neq 0$. Thus $P(l) \neq 0$. So the condition (A_2) is satisfied.

Now applying Theorem 2.1 to the set of zeros of polynomial $P(z) = z^d + (2z+5)^d + 1$ we obtain conclusion of Theorem 2.2. ■

References

- [1] **Mues, E. and M. Reinders**, Meromorphic functions sharing one value and unique range sets, *Kodai Math. J.*, **18** (1995), 515–522.
- [2] **Gross, F. and C.C. Yang**, On preimage and range sets of meromorphic functions, *Proc. Japan Acad. Ser. A Math. Sci.*, **58** (1982), 1–20.
- [3] **Frank, G. and M. Reinders**, A unique range set for meromorphic functions with 11 elements, *Complex Variables Theory Appl.*, **37(1-4)** (1998), 185–193.
- [4] **Fujimoto, H.**, On uniqueness of meromorphic functions sharing finite sets, *Amer. J. Math.*, **122(6)** (2000), 1175–1203.
- [5] **Ha Huy Khoai, Vu Hoai An and Le Quang Ninh**, Uniqueness theorems for holomorphic curves with hypersurfaces of Fermat–Waring type, *Complex Anal. Oper. Theory*, **8** (2014), 591–794.
- [6] **Yi, H.X.**, Unicity theorems for meromorphic or entire functions III, *Bull. Austr. Math. Soc.*, **53** (1996), 71–82.
- [7] **Masuda, K. and J. Noguchi**, A construction of hyperbolic hypersurface of $P^N(\mathbb{C})$, *Math. Ann.*, **304** (1996), 339–362.
- [8] **Hayman, W.K.**, *Meromorphic Functions*, Clarendon, Oxford, 1964.

V.H. An and P.N. Hoa

Hai Duong College

Hai Duong Province

Thang Long Institute of Mathematics and Applied Sciences

Ha Noi City

Vietnam

vuhoaianmai@yahoo.com

hphamngoc577@gmail.com