

ON SYMMETRY OF MAKÓ-PÁLES MEANS

Pál Burai (Debrecen, Hungary)

Justyna Jarczyk (Zielona Góra, Poland)

Communicated by Antal Járai

(Received March 9, 2018; accepted June 25, 2018)

Abstract. Given a nonempty real interval I , a continuous strictly monotonic function $\varphi : I \rightarrow \mathbb{R}$, and a Borel probability measure μ on $[0, 1]$, we characterize all symmetric Makó-Páles means $M_{\varphi, \mu}$ defined on $I \times I$ by

$$M_{\varphi, \mu}(x, y) = \varphi^{-1} \left(\int_0^1 \varphi(tx + (1-t)y) d\mu(t) \right).$$

1. Introduction

In the whole paper I stands for a nonempty interval of reals. Among two variable means on I , that is functions $M : I \times I \rightarrow I$ satisfying the condition

$$\min \{x, y\} \leq M(x, y) \leq \max \{x, y\}, \quad x, y \in I,$$

the Makó-Páles means play an important role. Denote by $\mathcal{CM}(I)$ the class of all continuous strictly monotonic functions mapping I into \mathbb{R} . Let $\varphi \in \mathcal{CM}(I)$

Key words and phrases: Mean, Makó-Páles mean.

2010 Mathematics Subject Classification: Primary 26E60, Secondary 39B22.

The research of the first author has been supported by the Hungarian Scientific Research Fund OTKA K-111651

<https://doi.org/10.71352/ac.47.173>

and μ be a Borel probability measure on $[0, 1]$. In 2008 Makó and Páles [2] introduced the mean $M_{\varphi, \mu}$ on I by the formula

$$M_{\varphi, \mu}(x, y) = \varphi^{-1} \left(\int_0^1 \varphi(tx + (1-t)y) d\mu(t) \right).$$

The class of Makó–Páles means is pretty large. Taking any $p \in (0, 1)$ and $\mu = (1-p)\delta_0 + p\delta_1$, where δ_t denotes the Dirac measure on $[0, 1]$ concentrated at a given fixed $t \in [0, 1]$, one can see that $M_{\varphi, \mu}$ is a weighted quasi-arithmetic mean

$$A_p^\varphi(x, y) = \varphi^{-1} (p\varphi(x) + (1-p)\varphi(y)), \quad x, y \in I.$$

If μ is the Lebesgue measure on $[0, 1]$, then for all $x, y \in I$, substituting $u = tx + (1-t)y$ and using the change-of-variable theorem, one can see that $M_{\varphi, \mu} = L_\varphi$, where the formula

$$L_\varphi(x, y) = \begin{cases} \varphi^{-1} \left(\frac{1}{x-y} \int_x^y \varphi(u) du \right), & \text{if } x, y \in I, \quad x \neq y, \\ x, & \text{if } x, y \in I, \quad x = y, \end{cases}$$

defines the Lagrangean mean. Therefore the class of Makó–Páles means contains two important classes of means. The investigation presented here is, in some sense, a continuation of that given in [1].

On one hand, observe that any Lagrangean mean is symmetric. That is to say, $L_\varphi(x, y) = L_\varphi(y, x)$ for all $x, y \in I$. On the other hand, the mean A_p^φ is symmetric, i.e.

$$A_p^\varphi(x, y) = A_p^\varphi(y, x), \quad x, y \in I,$$

if and only if $p = 1/2$. Thus the following natural problem arises: Describe all the Makó–Páles means that are symmetric, that is, satisfy the condition

$$(1.1) \quad M_{\varphi, \mu}(x, y) = M_{\varphi, \mu}(y, x).$$

The result below provide a complete answer to that question.

Theorem. *Let μ be a Borel probability measure on $[0, 1]$ and $\varphi \in \mathcal{CM}(I)$ be a continuous strictly monotonic function. Then the mean $M_{\varphi, \mu}$ is symmetric if and only if at least one of the conditions below holds:*

(i) *the measure μ is symmetric with respect to $1/2$, that is*

$$(1.2) \quad \mu(1-A) = \mu(A)$$

for all Borel subsets A of $[0, 1]$;

(ii) φ is a polynomial of degree at most $n \in \mathbb{N}$ and

$$(1.3) \quad \int_0^1 t^k d\mu(t) = \int_0^1 (1-t)^k d\mu(t) \quad \text{for all } k = 1, \dots, n.$$

Condition (1.1) means that

$$\int_0^1 \varphi(tx + (1-t)y) d\mu(t) = \int_0^1 \varphi((1-t)x + ty) d\mu(t), \quad x, y \in I,$$

or equivalently, by putting

$$(1.4) \quad \mu^*(A) = \mu(1-A)$$

for all Borel sets $A \subset [0, 1]$ and using the change-of-variable theorem,

$$\int_0^1 \varphi(tx + (1-t)y) d\mu(t) = \int_0^1 \varphi(tx + (1-t)y) d\mu^*(t), \quad x, y \in I.$$

Consequently, the original condition (1.1) is equivalent to the requirement

$$(1.5) \quad M_{\varphi, \mu} = M_{\varphi, \mu^*},$$

which is a particular case of the general equality problem

$$(1.6) \quad M_{\varphi, \mu} = M_{\psi, \nu}$$

solved in [2]. So one can expect that the description of the symmetric Makó-Páles means can be derived from the paper [2]. A thorough reading of it shows that this is almost true. Applying [2, Corollary 3, Theorems 7-9 and 12-15], which solve equation (1.6), to (1.5) being a particular case of (1.6), the reader can obtain the assertion of the Theorem. However, the results of [2] require some regularity of φ and ψ .

In the next section we give a short immediate proof of the Theorem, making use of the main result of the paper [3] by Páles (used also in the proof of [2, Theorem 8]).

2. Proof of the Theorem

In what follows, with any Borel probability measure μ on $[0, 1]$ we associate its moments $\hat{\mu}_k$, $k \in \mathbb{N} \cup \{0\}$, given by

$$\hat{\mu}_k = \int_0^1 t^k d\mu(t).$$

Then, by the change-of-variable theorem, condition (1.3) can be reformulated as

$$(2.1) \quad \hat{\mu}_k = \hat{\mu}_k^* \quad \text{for all } k = 1, \dots, n,$$

where the measure μ^* is defined by (1.4).

Proof of the Theorem. Assume that the mean $M_{\varphi, \mu}$ is symmetric. If $\mu = \mu^*$, that is equality (1.2) holds for all Borel sets $A \subset [0, 1]$, then μ is symmetric with respect to $1/2$, i.e. (i) is satisfied. So assume that $\mu \neq \mu^*$. By well-known results of measure theory and Weierstrass approximation theorem there exists an $n \in \mathbb{N} \cup \{0\}$ such that

$$\hat{\mu}_{n+1} \neq \hat{\mu}_{n+1}^*.$$

Let n be the smallest such a number. Now, rewriting (1.5) in the form

$$\int_0^1 \varphi(tx + (1-t)y) d(\mu - \mu^*)(t) = 0, \quad x, y \in I,$$

and applying [3, Theorem 1] by Páles for the continuous function φ and signed Borel measure $\mu - \mu^*$, we see that φ is a polynomial of degree at most n and condition (2.1), i.e. (1.3), holds. However, since φ is strict monotonic, it is not constant and consequently, n cannot be zero. Having $n \geq 1$ we come to (ii).

Now assume (i). Then $\mu = \mu^*$, hence

$$\begin{aligned} & \int_0^1 \varphi(tx + (1-t)y) d\mu(t) = \\ &= \int_0^1 \varphi((1-s)x + sy) d\mu^*(s) = \int_0^1 \varphi(sy + (1-s)x) d\mu(s) \end{aligned}$$

for all $x, y \in I$, which implies the symmetry of $M_{\varphi, \mu}$.

Finally assume condition (ii). Fix any $x, y \in I$. Then also $I \ni t \mapsto \varphi(tx + (1 - t)y)$ is a polynomial of degree at most n , say

$$\varphi(tx + (1 - t)y) = \sum_{k=0}^n a_k t^k, \quad t \in [0, 1],$$

with some $a_0, \dots, a_n \in \mathbb{R}$. Then, by (1.3),

$$\begin{aligned} \int_0^1 \varphi(tx + (1 - t)y) d\mu(t) &= \int_0^1 \sum_{k=0}^n a_k t^k d\mu(t) = \int_0^1 \sum_{k=0}^n a_k (1 - t)^k d\mu(t) = \\ &= \int_0^1 \varphi((1 - t)x + ty) d\mu(t) \end{aligned}$$

which completes the proof. ■

Acknowledgement. The authors would like to thank the anonymous reviewer for his/her valuable suggestion to improve the statement of the Theorem.

References

- [1] **Burai, P. and J. Jarczyk J.**, Conditional homogeneity and translativity of Makó–Páles means, *Annales Univ. Sci. Budapest., Sect. Comp.*, **40** (2013), 159–172. MR 3129165
- [2] **Makó, Z. and Zs. Páles**, On the equality of generalized quasi-arithmetic means, *Publ. Math. Debrecen*, **72** (2008), 407–440. MR 2406700 (2009b:39029)
- [3] **Páles, Zs.**, On functional equations characterizing polynomials, *Acta Sci. Math. (Szeged)*, **74** (2008), 581–592. MR 2487933 (2009j:39026)

P. Burai

University of Debrecen
Faculty of Informatics
Debrecen
Hungary
burai.pal@inf.unideb.hu

J. Jarczyk

University of Zielona Góra
Faculty of Mathematics
Computer Science and Econometrics
Zielona Góra
Poland
j.jarczyk@wmie.uz.zgora.pl

