

SUMMABILITY IN MIXED-NORM HARDY SPACES

Ferenc Weisz (Budapest, Hungary)

Communicated by Ferenc Schipp

(Received February 20, 2018; accepted March 12, 2018)

Abstract. The mixed norm Hardy spaces $H_{\vec{p}}(\mathbb{R}^d)$ is investigated, where $\vec{p} = (p_1, \dots, p_d) \in (0, \infty]^d$. A general summability method, the so called θ -summability is considered for multi-dimensional Fourier transforms. Under some conditions on θ , it is proved that the maximal operator of the θ -means is bounded from $H_{\vec{p}}(\mathbb{R}^d)$ to $L_{\vec{p}}(\mathbb{R}^d)$. This implies some norm and almost everywhere convergence results for the θ -means, amongst others the generalization of the well known Lebesgue's theorem.

1. Introduction

It is due to Lebesgue [16] that the Fejér means [6] of the trigonometric Fourier transforms of a function $f \in L_p(\mathbb{R})$ ($1 \leq p < \infty$) converge almost everywhere to the function. In this paper we generalize this result to mixed norm Lebesgue spaces and other summability methods as well. A general method of summation, the so called θ -summation method, which is generated by a single function θ and which includes all well known summations, is studied intensively in the literature (see e.g. Butzer and Nessel [2], Trigub and Belinsky [24], Gát [7, 8, 9], Goginava [10, 11, 12], Persson, Tephnadze and Wall [18], Simon [19, 20] and Feichtinger and Weisz [4, 5, 27, 28, 29]). The means generated

Key words and phrases: Mixed norm Hardy spaces, atomic decomposition, θ -summability, maximal operator.

2010 Mathematics Subject Classification: Primary 42B08, Secondary 42A38, 42A24, 42B25, 42B30.

This research was supported by the Hungarian Scientific Research Funds (OTKA) No K115804.

<https://doi.org/10.71352/ac.48.233>

by the θ -summation are defined for multi-dimensional functions by

$$\sigma_T^\theta f(x) := \int_{\mathbb{R}^d} \theta\left(\frac{|u|}{T}\right) \widehat{f}(u) e^{2\pi i x \cdot u} du,$$

where $|\cdot|$ denotes the Euclidean norm and $\widehat{f}$ the Fourier transform of f . The choice $\theta(u) = \max(1 - |u|, 0)$ yields the Fejér summation.

Stein, Taibleson and Weiss [22] proved for the Bochner-Riesz summability that the maximal operator σ_*^θ of the θ -means is bounded from the Hardy space $H_p(\mathbb{R}^d)$ to $L_p(\mathbb{R}^d)$ if $p > p_0$ (see also Grafakos [13] and Lu [17]). Later we generalized this result to other summability methods in [4, 5, 27, 29].

In this paper, we generalize these results to mixed norm Lebesgue and Hardy spaces, $L_{\vec{p}}(\mathbb{R}^d)$ and $H_{\vec{p}}(\mathbb{R}^d)$, where $\vec{p} = (p_1, \dots, p_d) \in (0, \infty]^d$. We give the atomic decomposition of this Hardy space. If $\vec{p}$ is the vector $(p, \dots, p)$, then we get back the classical Lebesgue and Hardy spaces. Under some conditions on θ , we will prove that the maximal operator σ_*^θ is bounded from $H_{\vec{p}}(\mathbb{R}^d)$ to $L_{\vec{p}}(\mathbb{R}^d)$ when each $p_i > p_0$. As a consequence, we prove some norm and almost everywhere convergence results for the θ -means. In this way, the well known Lebesgue's theorem is generalized. As special cases of the θ -summation, we consider the Riesz, Bochner-Riesz, Weierstrass, Picard and Bessel summations.

2. Mixed norm Lebesgue spaces

The $L_p(\mathbb{R}^d)$ space is equipped with the quasi-norm

$$\|f\|_p := \left(\int_{\mathbb{R}^d} |f(x)|^p dx \right)^{1/p} \quad (0 < p < \infty),$$

with the usual modification for $p = \infty$. Here we integrate with respect to the Lebesgue measure λ . The Lebesgue measure of a set H will be denoted also by $|H|$. Benedek and Panzone [1] generalized this definition as follows. Let $\vec{p} = (p_1, \dots, p_d) \in (0, \infty]^d$. The *mixed-norm Lebesgue space* $L_{\vec{p}}(\mathbb{R}^d)$ is defined to be the set of all measurable functions f such that

$$\begin{aligned} \|f\|_{L_{\vec{p}}(\mathbb{R}^d)} &:= \\ &:= \left(\int_{\mathbb{R}} \dots \left(\int_{\mathbb{R}} \left(\int_{\mathbb{R}} |f(x_1, \dots, x_d)|^{p_1} dx_1 \right)^{p_2/p_1} dx_2 \right)^{p_3/p_2} \dots dx_d \right)^{1/p_d} < \infty, \end{aligned}$$

with the usual modifications if $p_i = \infty$ for some $i = 1, \dots, d$. If $\vec{p} = (p, \dots, p)$, then we get back the space $L_p(\mathbb{R}^d)$. Let

$$p_- := \min\{p_1, \dots, p_d\} \quad \text{and} \quad \underline{p} = \min\{p_-, 1\}.$$

It is known that

$$(2.1) \quad \| |f|^s \|_{L_{\vec{p}}(\mathbb{R}^d)} = \|f\|_{L_{s\vec{p}}(\mathbb{R}^d)}^s.$$

Given a locally integrable function f , the *Hardy-Littlewood maximal operator* M is defined by

$$Mf(x) := \sup_{x \in B} \frac{1}{|B|} \int_B |f(y)| dy \quad (x \in \mathbb{R}^d),$$

where the supremum is taken over all balls B of $\mathbb{R}^d$ containing x . It is known that M is bounded on $L_p(\mathbb{R}^d)$ if $1 < p < \infty$. This is extended to the mixed norm spaces in Huang at al. [14].

Lemma 1. *If $p_- > 1$ and $f \in L_{\vec{p}}(\mathbb{R}^d)$, then*

$$(2.2) \quad \|Mf\|_{L_{\vec{p}}(\mathbb{R}^d)} \leq C \|f\|_{L_{\vec{p}}(\mathbb{R}^d)}.$$

The vector-valued extension of inequality (2.2) holds also. In the classical case see Fefferman–Stein [3], for the mixed norm spaces see Huang at al. [14].

Lemma 2. *If $p_- > 1$ and $1 < r < \infty$, then*

$$\left\| \left(\sum_{j=1}^{\infty} (Mf_j)^r \right)^{1/r} \right\|_{L_{\vec{p}}(\mathbb{R}^d)} \leq C \left\| \left(\sum_{j=1}^{\infty} |f_j|^r \right)^{1/r} \right\|_{L_{\vec{p}}(\mathbb{R}^d)}.$$

We will write $A \lesssim B$ if there exists a constant C such that $A \leq CB$.

3. Mixed norm Hardy spaces

Now we introduce the *mixed norm Hardy spaces* and give the atomic decompositions. Denote by $S(\mathbb{R}^d)$ the space of all Schwartz functions and by $S'(\mathbb{R}^d)$ the space of all tempered distributions. For $N \in \mathbb{N}$, let

$$\mathcal{F}_N(\mathbb{R}^d) := \left\{ \psi \in S(\mathbb{R}^d) : \sup_{\|\alpha\|_1 \leq N} \sup_{x \in \mathbb{R}^d} (1 + |x|)^N |\partial^\alpha \psi(x)| \leq 1 \right\},$$

where $\|\alpha\|_1 = |\alpha_1| + \cdots + |\alpha_d|$. For $t \in (0, \infty)$ and $\xi \in \mathbb{R}^d$, let

$$\psi_t(\xi) := t^{-d}\psi(\xi/t).$$

For any $d(1/p_- - 1) + 1 < N < \infty$, the *non-tangential grand maximal function* of $f \in S'(\mathbb{R}^d)$ is defined by

$$f_{\square}(x) := \sup_{\psi \in \mathcal{F}_N(\mathbb{R}^d)} \sup_{0 < t < \infty, |y-x| < t} |f * \psi_t(y)|.$$

Let $d(1/p_- - 1) + 1 < N < \infty$ be a positive integer. The mixed norm Hardy spaces $H_{\vec{p}}(\mathbb{R}^d)$ are consisting of all tempered distributions $f \in S'(\mathbb{R}^d)$ such that

$$\|f\|_{H_{\vec{p}}(\mathbb{R}^d)} := \|f_{\square}\|_{L_{\vec{p}}(\mathbb{R}^d)} < \infty.$$

It is known that different integers N give the same space with equivalent norms. Moreover, all $f \in H_{\vec{p}}(\mathbb{R}^d)$ are bounded distributions, i.e. $f * \phi \in L_{\infty}(\mathbb{R}^d)$ for all $\phi \in S(\mathbb{R}^d)$. Similarly to the classical case, one can show (see Huang et al. [14]) that

$$H_{\vec{p}}(\mathbb{R}^d) \sim L_{\vec{p}}(\mathbb{R}^d)$$

whenever $p_- > 1$. If each $p_i = p$, then we get back the classical Hardy spaces $H_p(\mathbb{R}^d)$ investigated in Fefferman, Stein and Weiss [3, 23, 21], Lu [17], Uchiyama [25].

The atomic decomposition is a useful characterization of the Hardy spaces by the help of which some boundedness results, duality theorems, inequalities and interpolation results can be proved. A measurable function a is called a $\vec{p}$ -atom if there exists a ball B such that

- (a) $\text{supp } a \subset B$,
- (b) $\|a\|_{L_{\infty}(\mathbb{R}^d)} \leq \frac{1}{\|\chi_B\|_{L_{\vec{p}}(\mathbb{R}^d)}}$,
- (c) $\int_{\mathbb{R}^d} a(x)x^{\alpha}dx = 0$ for all multi-indices α with $|\alpha| \leq s$,

where $d(1/p_- - 1) < s < \infty$ is an integer. In the classical case, i.e., if each $p_i = p$, the atomic decomposition theorem can be formulated as follows (see e.g. Latter [15], Lu [17]). Assume that $0 < p \leq 1$. A tempered distribution f is in $H_p(\mathbb{R}^d)$ if and only if there exist a sequence $\{a_i\}_{i \in \mathbb{N}}$ of p -atoms with support $\{B_i\}_{i \in \mathbb{N}}$ and a sequence $\{\lambda_i\}_{i \in \mathbb{N}}$ of positive numbers such that $f = \sum_{i \in \mathbb{N}} \lambda_i a_i$ in $S'(\mathbb{R}^d)$. Moreover,

$$\|f\|_{H_p(\mathbb{R}^d)} \sim \inf \left(\sum_{i \in \mathbb{N}} \lambda_i^p \right)^{1/p}.$$

It is easy to see that the right hand sides of the previous and next equations are the same. Thus

$$\|f\|_{H_p(\mathbb{R}^d)} \sim \inf \left\| \left(\sum_{i \in \mathbb{N}} \left(\frac{\lambda_i \chi_{B_i}}{\|\chi_{B_i}\|_{L_p(\mathbb{R}^d)}} \right)^p \right)^{1/p} \right\|_{L_p(\mathbb{R}^d)}.$$

And this form can be generalized to all $0 < p < \infty$ as follows. The next theorem is due to Huang et al. [14].

Theorem 1. *A tempered distribution $f \in S'(\mathbb{R}^d)$ is in $H_{\vec{p}}(\mathbb{R}^d)$ if and only if there exist a sequence $\{a_i\}_{i \in \mathbb{N}}$ of $\vec{p}$ -atoms with support $\{B_i\}_{i \in \mathbb{N}}$ and a sequence $\{\lambda_i\}_{i \in \mathbb{N}}$ of positive numbers such that*

$$f = \sum_{i \in \mathbb{N}} \lambda_i a_i \quad \text{in} \quad S'(\mathbb{R}^d).$$

Moreover,

$$\|f\|_{H_{\vec{p}}(\mathbb{R}^d)} \sim \inf \left\| \left(\sum_{i \in \mathbb{N}} \left(\frac{\lambda_i \chi_{B_i}}{\|\chi_{B_i}\|_{L_{\vec{p}}(\mathbb{R}^d)}} \right)^p \right)^{1/p} \right\|_{L_{\vec{p}}(\mathbb{R}^d)},$$

where the infimum is taken over all decompositions of f as above.

4. θ -summability of Fourier transforms

The Fourier transform of a function $f \in L^1(\mathbb{R}^d)$ is defined by

$$\widehat{f}(x) := \int_{\mathbb{R}^d} f(t) e^{-2\pi i x \cdot t} dt \quad (x \in \mathbb{R}^d),$$

where $i = \sqrt{-1}$ and $x \cdot t := \sum_{k=1}^d x_k t_k$. Suppose first that $f \in L_p(\mathbb{R}^d)$ for some $1 \leq p \leq 2$. The Fourier inversion formula

$$f(x) = \int_{\mathbb{R}^d} \widehat{f}(t) e^{2\pi i x \cdot t} dt \quad (x \in \mathbb{R}^d)$$

holds if $\widehat{f} \in L^1(\mathbb{R}^d)$. This motivates the following definition of θ -summability, which is a general summation generated by a single function $\theta : [0, \infty) \rightarrow \mathbb{R}$.

This summation was considered in a great number of papers and books, see e.g. Butzer and Nessel [2], Grafakos [13], Trigub and Belinsky [24] and Feichtinger and Weisz [5, 27, 28, 29] and the references therein. Let $\theta_0(x) := \theta(|x|)$ and suppose that

$$(4.1) \quad \theta \in C_0[0, \infty), \quad \theta(0) = 1, \quad \theta_0 \in L_1(\mathbb{R}^d), \quad \widehat{\theta}_0 \in L^1(\mathbb{R}^d),$$

where $C_0[0, \infty)$ denotes the spaces of continuous functions vanishing at infinity and $|\cdot|$ denotes the Euclidean norm. For $T > 0$, the T th θ -mean of the function $f \in L_p(\mathbb{R}^d)$ ($1 \leq p \leq 2$) is given by

$$\sigma_T^\theta f(x) := \int_{\mathbb{R}^d} \theta\left(\frac{|u|}{T}\right) \widehat{f}(u) e^{2\pi i x \cdot u} du \quad (x \in \mathbb{R}^d, T > 0).$$

This integral is well defined because $\theta_0 \in L^p(\mathbb{R}^d)$ and $\widehat{f} \in L_{p'}(\mathbb{R}^d)$, where $1/p + 1/p' = 1$.

For an integrable function f , it is known that we can rewrite $\sigma_T^\theta f$ as

$$\sigma_T^\theta f(x) = \int_{\mathbb{R}^d} f(x-t) K_T^\theta(t) dt = f * K_T^\theta(x) \quad (x \in \mathbb{R}^d, T > 0),$$

where the T th θ -kernel is given by

$$K_T^\theta(x) := \int_{\mathbb{R}^d} \theta\left(\frac{|t|}{T}\right) e^{2\pi i x \cdot t} dt = T^d \widehat{\theta}_0(Tx) \quad (x \in \mathbb{R}^d, T > 0).$$

We can extend the θ -means to all $f \in L_{\widehat{p}}(\mathbb{R}^d)$ with $p_- \geq 1$ and to all $f \in H_{\widehat{p}}(\mathbb{R}^d)$ with $p_- > 0$ by

$$\sigma_T^\theta f := f * K_T^\theta \quad (T > 0).$$

The maximal θ -operator is introduced by

$$\sigma_*^\theta f := \sup_{T>0} |\sigma_T^\theta f|.$$

For a ball B with center c and radius ρ , let τB denotes the ball with the same center and with radius $\tau\rho$ ($\tau > 0$). The following theorem can be proved as in [26].

Theorem 2. *Suppose that (4.1) is satisfied, $\widehat{\theta}_0$ is $(N+1)$ -times differentiable for some $N \in \mathbb{N}$ and there exists $d+N < \beta \leq d+N+1$ such that*

$$(4.2) \quad \left| \partial_1^{i_1} \dots \partial_d^{i_d} \widehat{\theta}_0(x) \right| \leq C|x|^{-\beta} \quad (x \neq 0)$$

whenever $i_1 + \dots + i_d = N$ or $i_1 + \dots + i_d = N + 1$. Then

$$(4.3) \quad |\sigma_*^\theta a(x)| \leq C \|\chi_B\|_{L_{\vec{p}}(\mathbb{R}^d)}^{-1} |M\chi_B(x)|^{\beta/d}$$

for all $\vec{p}$ -atoms a and all $x \notin 2B$, where the ball B is the support of the atom. If $\beta = d + N + 1$, then it is enough to suppose that (4.2) holds whenever $i_1 + \dots + i_d = N + 1$.

5. Boundedness in $H_{\vec{p}}(\mathbb{R}^d)$

In the proof of the boundedness of σ_*^θ , we will use the next lemma.

Lemma 3. *Let (4.1) be satisfied. If $\lim_{k \rightarrow \infty} f_k = f$ in the $H_{\vec{p}}(\mathbb{R}^d)$ -norm, then $\lim_{k \rightarrow \infty} \sigma_t^\theta f_k = \sigma_t^\theta f$ in $S'(\mathbb{R}^d)$ for all $t > 0$.*

Proof. The proof is similar to that of Theorem 7 in [26], so we outline the differences, only. We have to show that $\sigma_t^\theta f$ is a tempered distribution for each $f \in H_{\vec{p}}(\mathbb{R}^d)$ and $t > 0$. To this end, the main point is to show that $f * \check{h}_k$ is uniformly bounded in k if $\lim_{k \rightarrow \infty} h_k = h$ in $S(\mathbb{R}^d)$, where $\check{h}(x) := h(-x)$. We may suppose that $h \in \mathcal{F}_N(\mathbb{R}^d)$, and hence that $h_k \in \mathcal{F}_N(\mathbb{R}^d)$ for large k 's. Then for such a k ,

$$\left| (f * \check{h}_k)(x) \right| \leq f_\square(y) \quad \text{for every } y \text{ with } |x - y| \leq 1.$$

Thus, with the same x and y ,

$$\begin{aligned} & \left| f * \check{h}_k(x) \right| \leq \\ & \leq \left(\int_{-1/2}^{1/2} \dots \left(\int_{-1/2}^{1/2} \left(\int_{-1/2}^{1/2} |f_\square(y_1, \dots, y_d)|^{p_1} dy_1 \right)^{p_2/p_1} dy_2 \right)^{p_3/p_2} \dots dy_d \right)^{1/p_d} \leq \\ & \leq \|f\|_{H_{\vec{p}}(\mathbb{R}^d)}, \end{aligned}$$

which shows the uniform boundedness of $f * \check{h}_k$. The proof can be finished as Theorem 7 in [26]. ■

Theorem 3. *If (4.1) and (4.2) are satisfied and $p_- > d/\beta$, then*

$$\|\sigma_*^\theta f\|_{L_{\vec{p}}(\mathbb{R}^d)} \lesssim \|f\|_{H_{\vec{p}}(\mathbb{R}^d)} \quad (f \in H_{\vec{p}}(\mathbb{R}^d)).$$

Proof. By the atomic decomposition theorem, $f \in H_{\vec{p}}(\mathbb{R}^d)$ can be written as

$$f = \sum_{i \in \mathbb{N}} \lambda_i a_i,$$

where λ_i is positive and a_i is a $\vec{p}$ -atom with support B_i . It is known (see e.g. Weisz [28]) that the series converge in the $H_1(\mathbb{R}^d)$ -norm as well as in the $L_1(\mathbb{R}^d)$ -norm if $f \in H_{\vec{p}}(\mathbb{R}^d) \cap H_1(\mathbb{R}^d)$. It is easy to see that σ_t^θ is bounded on the $L_1(\mathbb{R}^d)$ space, hence

$$\sigma_t^\theta(f) = \sum_{i \in \mathbb{N}} \lambda_i \sigma_t^\theta(a_i) \quad (t > 0)$$

and so

$$\sigma_*^\theta(f) \leq \sum_{i \in \mathbb{N}} \lambda_i \sigma_*^\theta(a_i).$$

Then

$$\begin{aligned} \|\sigma_*^\theta f\|_{L_{\vec{p}}(\mathbb{R}^d)} &\lesssim \left\| \sum_{i \in \mathbb{N}} \lambda_i \sigma_*^\theta(a_i) \chi_{2B_i} \right\|_{L_{\vec{p}}(\mathbb{R}^d)} + \left\| \sum_{i \in \mathbb{N}} \lambda_i \sigma_*^\theta(a_i) \chi_{(2B_i)^c} \right\|_{L_{\vec{p}}(\mathbb{R}^d)} = \\ &=: A_1 + A_2. \end{aligned}$$

Using (2.1) and the fact that $\underline{p} \leq 1$, we can see

$$A_1 \leq \left\| \sum_{i \in \mathbb{N}} \lambda_i^{\underline{p}} \sigma_*^\theta(a_i)^{\underline{p}} \chi_{2B_i} \right\|_{L_{\vec{p}/\underline{p}}(\mathbb{R}^d)}^{1/\underline{p}}.$$

Let $(\vec{p})' = (p'_1, \dots, p'_d)$ denote the conjugate index vector, where $\frac{1}{p_i} + \frac{1}{p'_i} = 1$ for every $i = 1, \dots, d$. By Theorem 1 of Benedek and Panzone [1], there exists $g \in L_{(\vec{p}/\underline{p})'}(\mathbb{R}^d)$ with $\|g\|_{L_{(\vec{p}/\underline{p})'}(\mathbb{R}^d)} \leq 1$ such that

$$\left\| \sum_{i \in \mathbb{N}} \lambda_i^{\underline{p}} \sigma_*^\theta(a_i)^{\underline{p}} \chi_{2B_i} \right\|_{L_{\vec{p}/\underline{p}}(\mathbb{R}^d)} = \int_{\mathbb{R}^d} \sum_{i \in \mathbb{N}} \lambda_i^{\underline{p}} \sigma_*^\theta(a_i)^{\underline{p}} \chi_{2B_i} g.$$

Choosing a real number r such that $p_i/\underline{p} < r < \infty$ for all $i = 1, \dots, d$ and applying Hölder's inequality, we deduce

$$\begin{aligned} A_1^{\underline{p}} &\leq \int_{\mathbb{R}^d} \sum_{i \in \mathbb{N}} \lambda_i^{\underline{p}} \sigma_*^\theta(a_i)^{\underline{p}} \chi_{2B_i} g \leq \\ &\leq \sum_{i \in \mathbb{N}} \lambda_i^{\underline{p}} \|\sigma_*^\theta(a_i)^{\underline{p}} \chi_{2B_i}\|_{L_r(\mathbb{R}^d)} \|\chi_{2B_i} g\|_{L_{r'}(\mathbb{R}^d)} \lesssim \\ &\lesssim \sum_{i \in \mathbb{N}} \lambda_i^{\underline{p}} \|\sigma_*^\theta(a_i)^{\underline{p}}\|_{L_\infty(\mathbb{R}^d)} \lambda(2B_i)^{1/r} \|\chi_{2B_i} g\|_{L_{r'}(\mathbb{R}^d)}. \end{aligned}$$

Observe that σ_*^θ is bounded from $L_\infty(\mathbb{R}^d)$ to $L_\infty(\mathbb{R}^d)$. By the definition of the $\vec{p}$ -atom,

$$\begin{aligned} A_1^p &\lesssim \sum_{i \in \mathbb{N}} \lambda_i^p \|\chi_{B_i}\|_{L_{\vec{p}}(\mathbb{R}^d)}^{-p} \lambda(2B_i) \left(\frac{1}{\lambda(2B_i)} \int_{2B_i} g^{r'} \right)^{1/r'} \leq \\ &\leq \int_{\mathbb{R}^d} \sum_{i \in \mathbb{N}} \lambda_i^p \|\chi_{B_i}\|_{L_{\vec{p}}(\mathbb{R}^d)}^{-p} \chi_{2B_i} \left(M(g^{r'}) \right)^{1/r'} d\lambda. \end{aligned}$$

Again by Hölder's inequality,

$$A_1^p \lesssim \left\| \sum_{i \in \mathbb{N}} \lambda_i^p \|\chi_{B_i}\|_{L_{\vec{p}}(\mathbb{R}^d)}^{-p} \chi_{2B_i} \right\|_{L_{\vec{p}/\underline{p}}(\mathbb{R}^d)} \left\| \left(M(g^{r'}) \right)^{1/r'} \right\|_{L_{(\vec{p}/\underline{p})'(\mathbb{R}^d)}}.$$

Since $p_i/\underline{p} < r < \infty$ imply $(p_i/\underline{p})' > r'$ ($i = 1, \dots, d$), we get by (2.1) and (2.2) that

$$\begin{aligned} A_1 &\lesssim \left\| \sum_{i \in \mathbb{N}} \lambda_i^p \|\chi_{B_i}\|_{L_{\vec{p}}(\mathbb{R}^d)}^{-p} \chi_{2B_i} \right\|_{L_{\vec{p}/\underline{p}}(\mathbb{R}^d)}^{1/\underline{p}} \left\| M(g^{r'}) \right\|_{L_{((\vec{p}/\underline{p})')/r'}(\mathbb{R}^d)}^{1/p r'} \lesssim \\ &\lesssim \left\| \sum_{i \in \mathbb{N}} \lambda_i^p \|\chi_{B_i}\|_{L_{\vec{p}}(\mathbb{R}^d)}^{-p} \chi_{2B_i} \right\|_{L_{\vec{p}/\underline{p}}(\mathbb{R}^d)}^{1/\underline{p}} \|g\|_{L_{(\vec{p}/\underline{p})'(\mathbb{R}^d)}^{1/p}} \lesssim \\ &\lesssim \left\| \left(\sum_{i \in \mathbb{N}} \left(\frac{\lambda_i \chi_{2B_i}}{\|\chi_{2B_i}\|_{L_{\vec{p}}(\mathbb{R}^d)}} \right)^p \right)^{1/p} \right\|_{L_{\vec{p}}(\mathbb{R}^d)} \lesssim \|f\|_{H_{\vec{p}}(\mathbb{R}^d)}. \end{aligned}$$

On the other hand, using (4.3) and Lemma 2, we establish that

$$\begin{aligned} A_2 &\lesssim \left\| \sum_{i \in \mathbb{N}} \lambda_i \|\chi_{B_i}\|_{L_{\vec{p}}(\mathbb{R}^d)}^{-1} |M\chi_{B_i}|^{\beta/d} \chi_{(2B_i)^c} \right\|_{L_{\vec{p}}(\mathbb{R}^d)} \leq \\ &\leq \left\| \left(\sum_{i \in \mathbb{N}} \left(\lambda_i^{d/\beta} \|\chi_{B_i}\|_{L_{\vec{p}}(\mathbb{R}^d)}^{-d/\beta} |M\chi_{B_i}| \right)^{\beta/d} \right)^{d/\beta} \right\|_{L_{\beta\vec{p}/d}(\mathbb{R}^d)}^{\beta/d} \leq \\ &\leq \left\| \left(\sum_{i \in \mathbb{N}} \lambda_i \|\chi_{B_i}\|_{L_{\vec{p}}(\mathbb{R}^d)}^{-1} \chi_{B_i} \right)^{d/\beta} \right\|_{L_{\beta\vec{p}/d}(\mathbb{R}^d)}^{\beta/d} \lesssim \\ &\lesssim \left\| \sum_{i \in \mathbb{N}} \frac{\lambda_i \chi_{B_i}}{\|\chi_{B_i}\|_{L_{\vec{p}}(\mathbb{R}^d)}} \right\|_{L_{\vec{p}}(\mathbb{R}^d)} \lesssim \|f\|_{H_{\vec{p}}(\mathbb{R}^d)}, \end{aligned}$$

which proves the theorem for $f \in H_{\vec{p}}(\mathbb{R}^d) \cap H_1(\mathbb{R}^d)$. Note that $\beta/d > 1$ and $p_- > d/\beta$. Using lemma 3, the proof can be finished by a standard density argument as in [26]. $\blacksquare$

Note that if each $p_i = p$, then we get back the classical result (see Weisz [27, 28]). The classical result was proved in a special case, for the Bochner-Riesz means in Stein, Taibleson and Weiss [22], Grafakos [13] and Lu [17]. For the same case [22] contains a counterexample which shows that the theorem is not true for $p \leq n/\beta$.

Using Theorem 3 and a usual density argument, we obtain the next convergence results. We do not give the details here, because they can be found in similar cases in [26].

Corollary 1. *Suppose that (4.1) and (4.2) are satisfied and $p_- > d/\beta$. If $f \in H_{\vec{p}}(\mathbb{R}^d)$, then $\sigma_T^\theta f$ converges almost everywhere as well as in the $L_{\vec{p}}(\mathbb{R}^d)$ -norm as $T \rightarrow \infty$.*

For functions from the Hardy spaces, the limit of $\sigma_T^\theta f$ will be exactly the function.

Corollary 2. *Suppose that (4.1) and (4.2) are satisfied and $p_- > d/\beta$. If $f \in H_{\vec{p}}(\mathbb{R}^d)$ and there exists an interval $I \subset \mathbb{R}^d$ such that the restriction $f|_I \in L_{\vec{r}}(I)$ with $r_- \geq 1$, then*

$$\lim_{T \rightarrow \infty} \sigma_T^\theta f(x) = f(x) \quad \text{for a.e. } x \in I \text{ as well as in the } L_{\vec{p}}(I)\text{-norm.}$$

The next consequence follows from the fact that $L_{\vec{p}}(\mathbb{R}^d)$ is equivalent to $H_{\vec{p}}(\mathbb{R}^d)$ if $p_- > 1$.

Corollary 3. *Suppose that (4.1) and (4.2) are satisfied and $p_- > d/\beta$. If $p_- > 1$ and $f \in L_{\vec{p}}(\mathbb{R}^d)$, then*

$$\lim_{T \rightarrow \infty} \sigma_T^\theta f(x) = f(x) \quad \text{for a.e. } x \in \mathbb{R}^d \text{ as well as in the } L_{\vec{p}}(\mathbb{R}^d)\text{-norm.}$$

6. Some summability methods

As special cases, we consider some summability methods. The details of the necessary computations are left to the reader.

6.1. Riesz summation

The function

$$\theta_0(t) = \begin{cases} (1 - |t|^\gamma)^\alpha, & \text{if } |t| > 1; \\ 0, & \text{if } |t| \leq 1 \end{cases} \quad (t \in \mathbb{R}^d)$$

defines the *Riesz summation* if $0 < \alpha < \infty$ and γ is a positive integer. It is called *Bochner-Riesz summation* if $\gamma = 2$. The next lemma can be found in Stein and Weiss [23] (see also Lu [17, p. 132] and Weisz [29]).

Lemma 4. *Condition (4.1) is satisfied if $\alpha > \frac{d-1}{2}$ and*

$$\left| \partial_1^{i_1} \dots \partial_d^{i_d} \widehat{\theta}_0(x) \right| \leq C|x|^{-d/2-\alpha-1/2} \quad (x \neq 0)$$

for all $i_1, \dots, i_d \in \mathbb{N}$.

The following result follows from Theorem 3.

Corollary 4. *If*

$$\alpha > \frac{d-1}{2}, \quad \frac{d}{d/2 + \alpha + 1/2} < p_- < \infty,$$

then

$$\|\sigma_*^\theta f\|_{L_{\vec{p}}(\mathbb{R}^d)} \lesssim \|f\|_{H_{\vec{p}}(\mathbb{R}^d)} \quad (f \in H_{\vec{p}}(\mathbb{R}^d)).$$

Moreover, the corresponding Corollaries 1–3 hold as well.

6.2. Weierstrass summation

The *Weierstrass summation* is defined by

$$(6.1) \quad \theta_0(t) = e^{-|t|^2/2} \quad (t \in \mathbb{R}^d)$$

or by

$$(6.2) \quad \theta_0(t) = e^{-|t|} \quad (t \in \mathbb{R}^d),$$

or, in the one-dimensional case, by

$$(6.3) \quad \theta_0(t) = e^{-|t|^\gamma} \quad (t \in \mathbb{R}, 1 \leq \gamma < \infty).$$

It is called **Abel summation** if $\gamma = 1$. It is known that in the first case $\widehat{\theta}_0(x) = e^{-|x|^2/2}$ and in the second one $\widehat{\theta}_0(x) = c_d/(1 + |x|^2)^{(d+1)/2}$ for some $c_d \in \mathbb{R}$ (see Stein and Weiss [23, p. 6.]). The following lemma is easy to verify.

Lemma 5. *Let θ_0 be as in (6.1) or in (6.2) or in (6.3). Then condition (4.1) is satisfied and for any $N \in \mathbb{N}$,*

$$\left| \partial_1^{i_1} \dots \partial_d^{i_d} \widehat{\theta}_0(x) \right| \leq C|x|^{-d-N-1} \quad (x \neq 0),$$

where $i_1 + \dots + i_d = N + 1$.

The following result is an easy consequence of Theorem 3.

Corollary 5. *Let θ_0 be as in (6.1) or in (6.2) or in (6.3). Then*

$$\|\sigma_*^\theta f\|_{L_{\vec{p}}(\mathbb{R}^d)} \lesssim \|f\|_{H_{\vec{p}}(\mathbb{R}^d)} \quad (f \in H_{\vec{p}}(\mathbb{R}^d)).$$

Moreover, the corresponding Corollaries 1–3 and hold as well.

6.3. Picard–Bessel summation

Now let

$$(6.4) \quad \theta_0(t) = \frac{1}{(1 + |t|^2)^{(d+1)/2}} \quad (t \in \mathbb{R}^d).$$

Here $\widehat{\theta}_0(x) = c_d e^{-|x|}$ for some $c_d \in \mathbb{R}^d$.

Corollary 6. *Let θ_0 be as in (6.4). Then Lemma 5 and Corollary 5 hold.*

References

- [1] **Benedek, A. and R. Panzone**, The spaces L^p , with mixed norm, *Duke Math. J.*, **28** (1961), 301–324.
- [2] **Butzer, P.L. and R.J. Nessel**, *Fourier Analysis and Approximation*, Birkhäuser Verlag, Basel, 1971.
- [3] **Fefferman, C. and E.M. Stein**, H^p spaces of several variables, *Acta Math.*, **129** (1972), 137–194.
- [4] **Feichtinger, H.G. and F. Weisz**, The Segal algebra $\mathbf{S}_0(\mathbb{R}^d)$ and norm summability of Fourier series and Fourier transforms, *Monatsh. Math.*, **148** (2006), 333–349.
- [5] **Feichtinger, H.G. and F. Weisz**, Wiener amalgams and pointwise summability of Fourier transforms and Fourier series, *Math. Proc. Cambridge Philos. Soc.*, **140** (2006), 509–536.
- [6] **Fejér, L.**, Untersuchungen über Fouriersche Reihen, *Math. Ann.*, **58** (1904), 51–69.
- [7] **Gát, G.**, Pointwise convergence of cone-like restricted two-dimensional $(C,1)$ means of trigonometric Fourier series, *J. Approx. Theory.*, **149** (2007), 74–102.

- [8] **Gát, G.**, Almost everywhere convergence of sequences of Cesàro and Riesz means of integrable functions with respect to the multidimensional Walsh system, *Acta Math. Sin., Engl. Ser.*, **30(2)** (2014), 311–322.
- [9] **Gát, G., U. Goginava and K. Nagy**, On the Marcinkiewicz-Fejér means of double Fourier series with respect to Walsh-Kaczmarz system, *Studia Sci. Math. Hungar.*, **46** (2009), 399–421.
- [10] **Goginava, U.**, Marcinkiewicz-Fejér means of d -dimensional Walsh-Fourier series, *J. Math. Anal. Appl.*, **307** (2005), 206–218.
- [11] **Goginava, U.**, Almost everywhere convergence of (C, α) -means of cubical partial sums of d -dimensional Walsh-Fourier series, *J. Approx. Theory*, **141** (2006), 8–28.
- [12] **Goginava, U.**, The maximal operator of the Marcinkiewicz-Fejér means of d -dimensional Walsh-Fourier series, *East J. Approx.*, **12** (2006), 295–302.
- [13] **Grafakos, L.**, *Classical and Modern Fourier Analysis*. Pearson Education, New Jersey, 2004.
- [14] **Huang, L., J. Liu, D. Yang and W. Yuan**, Atomic and Littlewood-Paley characterizations of anisotropic mixed-norm Hardy spaces and their applications, <http://arxiv.org/pdf/1801.06251.pdf>, 2018.
- [15] **Latter, R.H.**, A characterization of $H^p(\mathbb{R}^n)$ in terms of atoms, *Studia Math.*, **62** (1978), 92–101.
- [16] **Lebesgue, H.**, Recherches sur la convergence des séries de Fourier, *Math. Ann.*, **61** (1905), 251–280.
- [17] **Lu, S.**, *Four Lectures on Real H^p Spaces*, World Scientific, Singapore, 1995.
- [18] **Persson, L.E., G. Tephnadze and P. Wall**, Maximal operators of Vilenkin-Nörlund means, *J. Fourier Anal. Appl.*, **21(1)** (2015), 76–94.
- [19] **Simon, P.**, Cesàro summability with respect to two-parameter Walsh systems, *Monatsh. Math.*, **131** (2000), 321–334.
- [20] **Simon, P.**, (C, α) summability of Walsh-Kaczmarz-Fourier series, *J. Approx. Theory*, **127** (2004), 39–60.
- [21] **Stein, E.M.**, *Harmonic Analysis: Real-variable Methods, Orthogonality and Oscillatory Integrals*, Princeton Univ. Press, Princeton, N.J., 1993.
- [22] **Stein, E.M., M.H. Taibleson and G. Weiss**, Weak type estimates for maximal operators on certain H^p classes, *Rend. Circ. Mat. Palermo, Suppl.*, **1** (1981), 81–97.
- [23] **Stein, E.M. and G. Weiss**, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton Univ. Press, Princeton, N.J., 1971.
- [24] **Trigub, R.M. and E.S. Belinsky**, *Fourier Analysis and Approximation of Functions*, Kluwer Academic Publishers, Dordrecht, Boston, London, 2004.

- [25] **Uchiyama, A.**, *Hardy Spaces on the Euclidean Space*, Springer Monographs in Mathematics. Springer, Berlin, 2001.
- [26] **Weisz, F.**, Summability of Fourier transforms in variable Hardy and Hardy-Lorentz spaces, (preprint).
- [27] **Weisz, F.**, *Summability of Multi-dimensional Fourier Series and Hardy Spaces*, Mathematics and Its Applications. Kluwer Academic Publishers, Dordrecht, Boston, London, 2002.
- [28] **Weisz, F.**, Summability of multi-dimensional trigonometric Fourier series, *Surv. Approx. Theory*, 7:1–179, 2012.
- [29] **Weisz, F.**, *Convergence and Summability of Fourier Transforms and Hardy Spaces*, Applied and Numerical Harmonic Analysis. Springer, Birkhäuser, Basel, 2017.

F. Weisz

Department of Numerical Analysis

Eötvös L. University

Pázmány P. sétány 1/C.

H-1117 Budapest

Hungary

`weisz@inf.elte.hu`