

# DIGIT PATTERNS IN REAL NUMBERS CREATED FROM PERMUTATIONS

**Jean-Marie De Koninck** (Québec, Canada)

**Imre Kátai** (Budapest, Hungary)

*Dedicated to the memory of Dr. László Dringó*

Communicated by Karl-Heinz Indlekofer

(Received January 24, 2020; accepted June 17, 2020)

**Abstract.** Given a positive integer  $k$ , we construct a binary number  $0.a_1a_2a_3\dots$  having the property that any sequence  $a_{m+1}\dots a_{m+k}$  of  $k$  consecutive digits from its binary expansion appears with a frequency directly related to the various permutations of the set  $\{1, 2, \dots, k+1\}$ .

## 1. Introduction

Given a positive integer  $k$ , let  $\Pi_k$  be the set of the permutations of the set  $\{1, 2, \dots, k+1\}$ . Various interesting aspects of this set  $\Pi_k$  can be studied; see for instance the book of Pemmaraju [1]. Here, we use this set to construct real numbers with an interesting property, as follows. Given  $\pi \in \Pi_k$ , let  $j_1, j_2, \dots, j_{k+1}$  be defined by  $\pi(i) = j_i$ . Further set, for each  $h = 1, 2, \dots, k$ ,

$$\rho(j_h, j_{h+1}) = \begin{cases} 1 & \text{if } j_{h+1} > j_h, \\ 0 & \text{if } j_{h+1} < j_h. \end{cases}$$

Moreover, given  $(\delta_1, \delta_2, \dots, \delta_k) \in \{0, 1\}^k$ , set

$$D(\delta_1, \delta_2, \dots, \delta_k) := \#\{\pi \in \Pi_k : \rho(\pi(i), \pi(i+1)) = \delta_i \text{ for } i = 1, 2, \dots, k\}$$

---

*Key words and phrases:* Permutations, digit patterns.

*2010 Mathematics Subject Classification:* 05A05, 20B05.

<https://doi.org/10.71352/ac.50.093>

and

$$\kappa(\delta_1, \delta_2, \dots, \delta_k) := \frac{D(\delta_1, \delta_2, \dots, \delta_k)}{(k+1)!}.$$

As we will see in Section 4,

$$\kappa(\delta_1, \delta_2, \dots, \delta_k) \geq \frac{1}{(k+1)!} \quad (k = 1, 2, \dots).$$

To illustrate the function  $\kappa(\delta_1, \delta_2, \dots, \delta_k)$ , if we choose the case  $k = 4$ , we obtain the following table.

$(\delta_1, \delta_2, \delta_3, \delta_4)$	$D(\delta_1, \delta_2, \delta_3, \delta_4)$	$\kappa(\delta_1, \delta_2, \delta_3, \delta_4)$
(0,0,0,0)	1	1/120
(0,0,0,1)	4	1/30
(0,0,1,0)	9	3/40
(0,0,1,1)	6	1/20
(0,1,0,0)	9	3/40
(0,1,0,1)	16	2/15
(0,1,1,0)	11	11/120
(0,1,1,1)	4	1/30
(1,0,0,0)	4	1/30
(1,0,0,1)	11	11/120
(1,0,1,0)	16	2/15
(1,0,1,1)	9	3/40
(1,1,0,0)	6	1/20
(1,1,0,1)	9	3/40
(1,1,1,0)	4	1/30
(1,1,1,1)	1	1/120

Our purpose in this short paper is to construct some binary number

$$\alpha = 0.a_1a_2a_3\dots,$$

that is, where each digit  $a_i \in \{0, 1\}$ , and such that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \#\{m \leq N : a_{m+1} \dots a_{m+k} = \delta_1 \dots \delta_k\} = \kappa(\delta_1, \dots, \delta_k).$$

To construct  $\alpha$ , we proceed as follows. First we set

$$\mathcal{F}_N = [e^N, e^{N+1}) \quad \text{and} \quad \mathcal{L}_N = [\log N, N] \quad (N = 1, 2, \dots).$$

Let  $p(n) = p_N(n)$  stand for the smallest prime divisor of  $n$  which is located in the interval  $\mathcal{L}_N$ . Observe that the number of those  $n \in \mathcal{F}_N$  which do not contain any prime divisors in  $\mathcal{L}_N$  is bounded by

$$ce^N \prod_{\substack{p \in \mathcal{L}_N \\ p \in \wp}} \left(1 - \frac{1}{p}\right) \leq ce^N \frac{\log \log N}{\log N}.$$

To each number  $n \in \mathcal{F}_N$ , we associate the number

$$\epsilon_n = \begin{cases} 1 & \text{if } p(n+1) > p(n) \text{ and } n+1 \in \mathcal{F}_N, \\ 0 & \text{otherwise} \end{cases}$$

for some absolute constant  $c > 0$ , where  $\wp$  stands for the set of all primes. Thus,  $\epsilon_n = 0$  if  $p(n+1) < p(n)$  or if  $n < e^{N+1} < n+1$  or if either  $p(n)$  or  $p(n+1)$  does not exist. Then, to each  $N \in \mathbb{N}$ , we associate the number

$$\xi_N = \text{Concat}(\epsilon_n : n \in \mathcal{F}_N),$$

and we then define

$$(1.1) \quad \alpha = 0.\xi_2\xi_3\xi_4\dots$$

## 2. The distribution function of $(\{2^n\alpha\})_{n \geq 1}$

With  $\alpha$  as in (1.1), let  $0 < u < 1$  written as

$$(2.1) \quad u = \frac{t_1}{2} + \frac{t_2}{2^2} + \frac{t_3}{2^3} + \dots$$

Here, we may assume that  $t_n = 0$  for infinitely many  $n \in \mathbb{N}$ . We can prove that

$$(2.2) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \#\{n \leq N : \{2^n\alpha\} \leq u\} = F(u) \quad \text{exists.}$$

To see this, we proceed as follows. Let  $r_1 < r_2 < \dots$  be a sequence of integers such that  $t_{r_j} = 0$  for some  $j \in \mathbb{N}$  and then set  $u_j := \sum_{\nu=1}^{r_j-1} \frac{t_\nu}{2^\nu}$  and further define

$$\tilde{u}_j := \frac{1}{2^{r_j}} + u_j. \quad \text{It is clear that}$$

$$u_j \leq u < \tilde{u}_j \quad (j \in \mathbb{N}).$$

We then introduce the two functions

$$\begin{aligned} F_1(u) &= \liminf_{N \rightarrow \infty} \frac{1}{N} \#\{n \leq N : \{2^n \alpha\} < u\}, \\ F_2(u) &= \limsup_{N \rightarrow \infty} \frac{1}{N} \#\{n \leq N : \{2^n \alpha\} < u\}. \end{aligned}$$

With these definitions, we easily see that

$$(2.3) \quad F(u_j) = \sum_{\frac{a_1}{2} + \frac{a_2}{2^2} + \dots + \frac{a_{r_j}}{2^{r_j}} \leq u_j} \kappa(a_1, \dots, a_{r_j}) \leq F_1(u),$$

$$(2.4) \quad F(\widetilde{u}_j) = \sum_{\frac{a_1}{2} + \frac{a_2}{2^2} + \dots + \frac{a_{r_j}}{2^{r_j}} \leq \widetilde{u}_j} \kappa(a_1, \dots, a_{r_j}) \geq F_2(u).$$

Moreover,

$$(2.5) \quad F(\widetilde{u}_j) - F(u_j) = \kappa(t_1, \dots, t_{r_{j-1}}, 1).$$

Also, observe that it will follow from Theorem 4.1 below that

$$(2.6) \quad \lim_{m \rightarrow \infty} \max_{\delta_1, \dots, \delta_m \in \{0,1\}^m} \kappa(\delta_1, \dots, \delta_m) = 0.$$

It then follows from (2.3), (2.4), (2.5) and (2.6) that  $F_1(u) = F_2(u)$  and therefore that  $F(u)$  exists, as claimed.

We can even prove that  $F(u)$  is a continuous function. To show this, we first fix  $u$  and choose two sequences of numbers  $(u_M)_{M \geq 1}$  and  $(v_M)_{M \geq 1}$  such that  $u_M < u < v_M$  for each  $M \geq 1$ , and such that  $u_M \rightarrow u$  and  $v_M \rightarrow u$  as  $M \rightarrow \infty$ . Then, let  $s$  be an integer such that  $\lfloor u \cdot 2^M \rfloor = s$  and choose  $u_M = \frac{s}{2^M}$  and  $v_M = \frac{s+1}{2^M}$ . We then have

$$\begin{aligned} F(u_M) &= \sum_{\frac{a_1}{2} + \dots + \frac{a_M}{2^M} \leq s/2^M} \kappa(a_1, \dots, a_M), \\ F(v_M) &= \sum_{\frac{a_1}{2} + \dots + \frac{a_M}{2^M} \leq (s+1)/2^M} \kappa(a_1, \dots, a_M), \end{aligned}$$

with

$$\frac{s+1}{2^M} = \frac{b_1}{2} + \dots + \frac{b_M}{2^M}.$$

Since  $F(v_M) - F(u_M) = \kappa(b_1, \dots, b_M) \rightarrow 0$  as  $M \rightarrow \infty$ , we therefore have that  $\lim_{M \rightarrow \infty} F(u_M) = F(u)$  and  $\lim_{M \rightarrow \infty} F(v_M) = F(u)$ . Thus we proved that  $F(u)$  is continuous at the points  $u \in \mathbb{R} \setminus \mathbb{Q}$  and also continuous from

the right at those points  $u \in \mathbb{Q}$  of the form  $u = s/(2^R)$ , where  $s$  is an odd integer. Moreover, assuming that  $K$  is an integer larger than  $R$  and setting  $u_K = u - 1/2^K$ , we have  $F(u) - F(u_K) = \kappa(t_1, t_2, \dots, t_{R-1}, 0, 1, 1, \dots, 1)$  which tends to 0 as  $K$  tends to infinity, thereby establishing that  $F(u)$  is continuous from the left, as well.

### 3. Main theorem

**Theorem 3.1.** *Given an integer  $k \geq 2$  and an arbitrary  $k$ -tuple  $(\delta_1, \dots, \delta_k) \in \{0, 1\}^k$ , we have*

$$\lim_{M \rightarrow \infty} \frac{1}{M} \# \left\{ m \leq M : \{2^m \alpha\} \in \left[ \frac{\delta_1}{2} + \dots + \frac{\delta_k}{2^k}, \frac{\delta_1}{2} + \dots + \frac{\delta_k}{2^k} + \frac{1}{2^k} \right) \right\} = \\ = \kappa(\delta_1, \dots, \delta_k).$$

**Proof.** Let  $p_1, p_2, \dots, p_{k+1}$  be distinct primes located in the interval  $\mathcal{L}_N$ . Let us count those  $n \in \mathcal{F}_N$  for which  $p(n+j) = p_j$ . Also, let  $\{i_1, \dots, i_{k+1}\}$  be that particular permutation of the set  $\{1, 2, \dots, k+1\}$  for which  $p_{i_1} < p_{i_2} < \dots < p_{i_{k+1}}$ . Then, set

$$Q_U := \prod_{\substack{\log N < p < U \\ p \in \wp}} p.$$

Since  $n+j \equiv 0 \pmod{p_j}$  for  $j = 1, 2, \dots, k+1$ , it follows that  $n = m \cdot p_1 p_2 \cdots p_{k+1} + r$  with  $(r, p_1 p_2 \cdots p_{k+1}) = 1$ . Moreover,  $(n+i_1, Q_{p_{i_1}}) = 1$ ,  $(n+i_2, Q_{p_{i_2}}) = 1, \dots, (n+i_{k+1}, Q_{p_{i_{k+1}}}) = 1$ . Using standard asymptotic sieve techniques, we can write these conditions in the form

$$\prod_{\ell=1}^{k+1} (n+i_\ell, Q_{p_{i_\ell}}) = 1, \prod_{\ell=2}^{k+1} \left( n+i_\ell, \frac{Q_{p_{i_2}}}{Q_{p_{i_1}}} \right) = 1, \dots, \left( n+i_{k+1}, \frac{Q_{p_{i_{k+1}}}}{Q_{p_{i_k}}} \right) = 1, \\ n \equiv r \pmod{p_1 p_2 \cdots p_{k+1}}.$$

Thus the number of such numbers  $n \in \mathcal{F}_N$  is, as  $N \rightarrow \infty$ ,

$$(1+o(1)) \frac{\#\mathcal{F}_N}{p_1 p_2 \cdots p_{k+1}} \cdot \prod_{\substack{\log N < q < p_{i_1} \\ q \in \wp}} \left( 1 - \frac{k+1}{q} \right) \cdot \\ \cdot \prod_{\substack{p_{i_1} < q < p_{i_2} \\ q \in \wp}} \left( 1 - \frac{k}{q} \right) \cdots \prod_{\substack{p_{i_k} < q < p_{i_{k+1}} \\ q \in \wp}} \left( 1 - \frac{1}{q} \right) = \\ = (1+o(1)) \#\mathcal{F}_N \cdot \log \log \log N \cdot \prod_{i=1}^{k+1} \frac{1}{p_i \log \log p_i}.$$

The important observation here is that this asymptotic behavior does not depend on the particular permutation of the primes  $p_1, p_2, \dots, p_{k+1}$  we choose. We may therefore conclude that, as  $N \rightarrow \infty$ ,

$$\begin{aligned} \frac{1}{\#\mathcal{F}_N} \# \left\{ m \in \mathcal{F}_N : \{2^m \alpha\} \in \left[ \frac{\delta_1}{2} + \dots + \frac{\delta_k}{2^k}, \frac{\delta_1}{2} + \dots + \frac{\delta_k}{2^k} + \frac{1}{2^k} \right) \right\} = \\ = (1 + o(1)) \kappa(\delta_1, \dots, \delta_k). \end{aligned}$$

Now, we need to count those  $\{2^m \alpha\}$  ( $m = 1, 2, \dots, \lfloor x \rfloor$ ) not only for the particular values  $x = e^N$ , but also for the more general values  $x \in (e^N, e^{N+1})$ .

So, let  $\varepsilon > 0$  be an arbitrarily small number and set  $x = e^{N+\theta}$  with  $0 < \theta < 1$ . We now examine two separate cases. If  $\theta < \varepsilon$ , then

$$\#\{n : e^N \leq n < x\} < e^N(e^\varepsilon - 1) < 2\varepsilon e^N.$$

On the other hand, if  $\theta > \varepsilon$ , setting  $S := [e^N, e^{N+\theta})$ , we may then repeat the above argument for the interval  $S$  instead of  $\mathcal{F}_N$  and obtain the same result. Therefore, in both cases, the proof is complete.  $\blacksquare$

#### 4. The size of $\kappa(\delta_1, \dots, \delta_k)$

**Theorem 4.1.** *Let  $k \geq 2$  and let  $a_1, a_2, \dots, a_k \in \{0, 1\}$  be given. Then,*

$$\frac{1}{(k+1)!} \leq \kappa(a_1, a_2, \dots, a_k) < \frac{1}{2^{\lfloor k/2 \rfloor}}.$$

**Proof.** First, we prove the first inequality, namely

$$(4.1) \quad \kappa(a_1, a_2, \dots, a_k) \geq \frac{1}{(k+1)!}.$$

To do so, we let  $j_1, \dots, j_r$  be the indices of those  $a_i$ 's for which  $a_{j_\nu} = 0$  ( $\nu = 1, \dots, r$ ) and let  $t_1, \dots, t_s$  be the indices of those  $a_i$ 's for which  $a_{t_\mu} = 1$  ( $\mu = 1, \dots, s$ ). The case where one of the two sets  $\{j_1, \dots, j_r\}$  or  $\{t_1, \dots, t_s\}$  is empty is much more simple. So, let  $S = \{1, \dots, r\}$  and  $M = \{r+1, \dots, k+1\}$ . Now, let  $\{u(1), \dots, u(k+1)\}$  be a permutation of  $\{1, 2, \dots, k+1\}$  satisfying

1.  $\{u(j_1 + 1), u(j_2 + 1), \dots, u(j_r + 1)\}$  is a permutation of  $S$  satisfying the condition

$$\text{If } j_{\ell+1} = j_\ell + 1 \text{ for some } \ell \in \{1, \dots, r\}, \text{ then } u(j_\ell + 1) > u(j_{\ell+1} + 1).$$

Observe that such a permutation clearly exists.

2.  $\{u(t_1 + 1), u(t_2 + 1), \dots, u(t_{k-r} + 1)\}$  is a permutation of  $M$  satisfying the condition

If  $t_{\nu+1} = t_\nu + 1$  for some  $\nu \in \{1, \dots, s\}$ , then  $u(t_\nu + 1) < u(t_{\nu+1} + 1)$ .

Such a permutation also clearly exists.

For such a permutation  $\{u(1), \dots, u(k+1)\}$ , we have that  $\rho(u(\ell), u(\ell+1)) = a_\ell$  for  $\ell = 1, \dots, k$ . The special case  $S = \emptyset$  is very simple, because in this case,  $u(j) = j$  for each  $j = 1, \dots, k+1$ . On the other hand if  $M = \emptyset$ , then

$$u(1) = k+1, u(2) = k, \dots, u(k+1) = 1.$$

This completes the proof of (4.1).

We will now prove the second inequality in Theorem 4.1, namely

$$(4.2) \quad \kappa(a_1, a_2, \dots, a_k) < \frac{1}{2^{\lfloor k/2 \rfloor}}.$$

Assume that  $\{j_1, j_2, \dots, j_{k+1}\}$  is a permutation of  $\{1, 2, \dots, k+1\}$  satisfying  $\rho(j_\ell, j_{\ell+1}) = a_\ell$  for  $\ell = 1, \dots, k$ . Assume first that  $k+1$  is even and consider the pairs

$$(j_1, j_2), \quad (j_3, j_4), \quad \dots, \quad (j_k, j_{k+1}).$$

If  $a_1 = 1$ , then  $j_1 < j_2$ ; if  $a_1 = 0$ , then  $j_2 > j_1$ ; if  $a_3 = 1$ , then  $j_3 < j_4$ ; if  $a_3 = 0$ , then  $j_3 > j_4$ , and so on, up to if  $a_k = 1$ , then  $j_k < j_{k+1}$ ; if  $a_k = 0$ , then  $j_{k+1} < j_k$ .

To sum up, this means that the number of associated permutations is no larger than  $\frac{(k+1)!}{2^{(k+1)/2}}$ .

The case where  $k+1$  is odd can be treated in a similar manner, since we then have that  $k$  is even, in which case we consider the  $k/2 + 1$  numbers

$$(j_1, j_2), \quad (j_3, j_4), \quad \dots, \quad (j_{k-1}, j_k), \quad j_{k+1},$$

which allows us in the end to conclude that the number of associated permutations is no larger than  $\frac{(k+1)!}{2^{k/2}}$ .

In both cases, we have proved (4.2) and the proof of Theorem 4.1 is complete. ■

## References

- [1] **Pemmaraju, S.**, *Computational Discrete Mathematics*, Cambridge University Press, 2003, xiv+480 pages.

### **J.-M. De Koninck**

Dép. math. et statistique  
Université Laval  
Québec  
Québec G1V 0A6  
Canada  
`jmdk@mat.ulaval.ca`

### **I. Káta**

Department of Computer Algebra  
Eötvös Loránd University  
H-1117 Budapest  
Pázmány Péter sétány 1/C  
Hungary  
`katai@inf.elte.hu`