

ON THE EQUATION $F(n^2 + 1) = bF(n^2 - 1) + c$

Imre Káta, Bui Minh Mai Khanh and Bui Minh Phong

(Budapest, Hungary)

Communicated by Gábor Farkas

(Received 16 March 2025; accepted 18 June 2025)

Abstract. Under a suitable conjecture concerning prime numbers, we provide a complete solution to the equation

$$F(n^2 + 1) = bF(n^2 - 1) + c \quad \text{for every } n \in \mathbb{N}, \quad n > 1,$$

where $F : \mathbb{N} \rightarrow \mathbb{C}$ is a completely multiplicative function and $b, c \in \mathbb{C}$, $c \neq 0$.

1. Introduction

Let \mathcal{P} , \mathbb{N} , \mathbb{Z} and \mathbb{C} denote the set of primes, positive integers, integers and complex numbers, respectively. Let $\mathbb{N}_1 := \mathbb{N} \setminus \{1\}$. We denote by \mathcal{M} (\mathcal{M}^*) the set of all complex-valued multiplicative (completely multiplicative) functions, respectively. For each $a \in \mathbb{Z}$ and $p \in \mathcal{P}$ let $\left(\frac{a}{p}\right)$ be the Legendre symbol. Let $P(m)$ be the largest prime divisor of $m \in \mathbb{N}$ and let p_k denote the k -th prime number. For each $k \in \mathbb{N}$ we denote by \mathcal{F}_k the set of all arithmetical functions $F : \mathbb{N} \rightarrow \mathbb{C}$ such that $F(n^2 + 1) = k$ for every $n \in \mathbb{N}$.

Let $\mathbb{E}(n) = 1$ and $\mathbb{I}(n) = n$ for every $n \in \mathbb{N}$.

The problem of characterising the identity function as a multiplicative arithmetical function satisfying certain equations has been studied by several authors. C. Spiro [4] proved that if $f \in \mathcal{M}$ satisfies the relations

$$f(p + q) = f(p) + f(q) \quad \text{for every } p, q \in \mathcal{P}$$

and $f(p_0) \neq 0$ for some $p_0 \in \mathcal{P}$, then f is the identity function.

Key words and phrases: Completely multiplicative function, the identity function, Dirichlet character.

2010 Mathematics Subject Classification: Primary 11A07, 11A25, 11K6, Secondary 11N25, 11N64.

<https://doi.org/10.71352/ac.58.180625>

First published online: 24 June 2025

In [1], we gave all solutions of the equation

$$F(n^2 + m^2 + k) = H(n) + H(m) + K \quad \text{for every } n \in \mathbb{N},$$

where $k \in \mathbb{N}$ is the sum of two fixed squares, $K \in \mathbb{C}$ and F, H are completely multiplicative functions.

The equation

$$G(n) = F(n^2 - 1) + D$$

was completely solved in [2], where $G, F \in \mathcal{M}^*$ and $D \in \mathbb{C}$.

Now we turn to the equation

$$F(Q(n)) = bF(Q(n) - \ell) + c,$$

where $\ell \in \mathbb{N}$, $b, c \in \mathbb{C}$ and $Q(x) \in \mathbb{Z}[x]$. The special case

$$(Q(n), \ell, b, c) = \{n^3, 1, 1, D\}$$

was solved in [3].

In this note we consider the case $Q(n) = n^2 + 1$, $\ell = 1$ and $b, c \in \mathbb{C}$, $c \neq 0$. Unfortunately, we can solve the equation only under the following conjecture:

Conjecture 1. *For every prime $p > p_{93} = 487$, there exists a positive integer m such that $p \mid m^2 - 1$ and*

$$P\left(\frac{(m^2 - 1)(m^2 + 1)}{p}\right) < p.$$

We note that, with the help of Maple, we have verified that Conjecture 1 holds for every p_k with $93 < k \leq 10^6$. In the range $k \in \{1, \dots, 93\}$, we could not verify Conjecture 1 for the primes p_k , where

$$\begin{aligned} k &\in \mathcal{F} := \\ &:= \{1, 2, 3, 4, 5, 6, 11, 13, 14, 15, 16, 18, 24, 25, 28, 29, 30, 33, 39, 56, 67, 74, 93\}. \end{aligned}$$

Define the sets

$$\mathcal{K} := \left\{ p \in \mathcal{P} \mid \left(\frac{-1}{p}\right) = \left(\frac{-7}{p}\right) = -1 \right\} \quad \text{and} \quad \mathcal{H} = \mathcal{P} \setminus \mathcal{K}.$$

It is easy to show that

$$\mathcal{K} = \left\{ p \in \mathcal{P} \mid p \equiv t \pmod{28}, \text{ where } t \in \{3, 19, 27\} \right\}.$$

In fact, for the proof of our main result, we require the following weaker conjecture.

Conjecture 2. *For every prime $p > p_{39} = 167$ with $p \in \mathcal{K}$, there exists a positive integer ℓ such that $p \mid \ell^2 - 1$ and*

$$P\left(\frac{(\ell^2 - 1)(\ell^2 + 1)}{p}\right) < p.$$

We have verified Conjecture 2 for every $p = p_k$ with $39 < k \leq 10^6$, and we have checked that in the range $k \in \{1, \dots, 39\}$, Conjecture 2 holds for all $p = p_k$ except:

$$p \in \{p_2 = 3, p_{11} = 31, p_{15} = 47, p_{39} = 167\}.$$

In this paper we prove the following result.

Theorem 1. *Assume that $b, c \in \mathbb{C}$, $c \neq 0$ and the function $F \in \mathcal{M}^*$ satisfies the relation*

$$(1.1) \quad F(n^2 + 1) = bF(n^2 - 1) + c \quad \text{for every } n \in \mathbb{N}_1.$$

Then the following assertions hold:

(a) *If $b = 0$, then $c = 1$ and $F \in \mathcal{F}_1$, furthermore*

$$F(p) = 1 \quad \text{for every } p \in \mathcal{P}, p \equiv 1 \pmod{4}.$$

(b) *If $bc \neq 0$ and Conjecture 2 holds, then*

$$(F, b, c) \in \{(\mathbb{E}, b, -b + 1), (\mathbb{I}, 1, 2)\}.$$

2. Lemmas

We first observe from the assumption $F \in \mathcal{M}^*$ and (1.1) that

$$(2.1) \quad E_n := F(n^2 + 1) - bF(n - 1)F(n + 1) - c = 0 \quad \text{for every } n \in \mathbb{N}_1.$$

Since

$$(2.2) \quad (n^2 + n + 1)^2 + 1 = (n^2 + 1)((n + 1)^2 + 1) \quad \text{for every } n \in \mathbb{N},$$

it follows from (2.1) and (2.2) that

$$bF(n^2 + n)F(n^2 + n + 2) + c = \left(bF(n - 1)F(n + 1) + c\right)\left(bF(n)F(n + 2) + c\right).$$

Therefore, we have the identity

$$(2.3) \quad \begin{aligned} L_n := & bF(n)F(n + 1)F(n^2 + n + 2) - \\ & - b^2F(n - 1)F(n)F(n + 1)F(n + 2) - \\ & - bcF(n)F(n + 2) - bcF(n - 1)F(n + 1) + c - c^2 = 0, \end{aligned}$$

which holds for every $n \in \mathbb{N}_1$.

From now on, let $F(2) := x$ and $F(3) := y$.

In the proof of our results, we will use the relation (2.1) for the values

$$n \in \{2, 3, 4, 5, 6, 7, 8, 17, 21, 31, 41, 57\}.$$

This yields the following system of equations:

$$\begin{aligned}
 (2.4) \quad & E_2 = F(5) - by - c = 0, \\
 & E_3 = xF(5) - bx^3 - c = 0, \\
 & E_4 = F(17) - bxF(5) - c = 0, \\
 & E_5 = xF(13) - byF(5) - c = 0, \\
 & E_6 = F(37) - bF(5)F(7) - c = 0, \\
 & E_7 = xF(5)^2 - bx^4y - c = 0, \\
 & E_8 = F(5)F(13) - by^2F(7) - c = 0, \\
 & E_{17} = xF(13)F(17) - bx^3F(5)F(11) - c = 0, \\
 & E_{21} = xF(5)F(29) - bx^5y^2 - c = 0, \\
 & E_{31} = xF(13)F(37) - bx^6yF(5) - c = 0, \\
 & E_{41} = xF(29)^2 - bx^4yF(5)F(7) - c = 0, \\
 & E_{57} = xF(5)^3F(13) - bx^4F(7)F(29) - c = 0.
 \end{aligned}$$

Lemma 1. *If $F \in \mathcal{M}^*$ and $c \in \mathbb{C}$ satisfy*

$$(2.5) \quad F(n^2 + 1) = c \quad \text{for every } n \in \mathbb{N},$$

then $c \in \{0, 1\}$.

If $c = 0$, then $F \in \mathcal{F}_0$.

If $c = 1$, then $F(2) = 1$ and

$$(2.6) \quad F(p) = 1 \quad \text{for every } p \in \mathcal{P}, \quad p \equiv 1 \pmod{4}.$$

Proof. From (2.2) and (2.5), we have

$$\begin{aligned}
 c &= F\left((n^2 + n + 1)^2 + 1\right) = F\left((n^2 + 1)((n + 1)^2 + 1)\right) = \\
 &= F(n^2 + 1)F((n + 1)^2 + 1) = c^2,
 \end{aligned}$$

which implies $c \in \{0, 1\}$.

If $c = 0$, then $F(n^2 + 1) = 0$ for every $n \in \mathbb{N}$ and so $F \in \mathcal{F}_0$.

If $c = 1$, then $F(n^2 + 1) = 1$ for every $n \in \mathbb{N}$ and so $F \in \mathcal{F}_1$.

We now prove (2.6). It is easy to check that $F(2) = 1, F(5) = F(13) = F(17) = 1$.

Assume that $F(q) = 1$ for every $q \in \mathcal{P}, q \equiv 1 \pmod{4}, q < P$ and let $P \in \mathcal{P}, P \equiv 1 \pmod{4}, P > 17$. Since $P \equiv 1 \pmod{4}$, there is $m, Q \in \mathbb{N}$ with $m < P$, such that

$$m^2 + 1 = PQ.$$

Clearly, $Q < P$ since $PQ = m^2 + 1 \leq (P - 1)^2 + 1$. For any $q \in \mathcal{P}, q|Q$, we have $m^2 + 1 \equiv 0 \pmod{q}$ and

$$(-1)^{\frac{q-1}{2}} = \left(\frac{-1}{q}\right) = \left(\frac{m^2}{q}\right) = 1,$$

so $q \equiv 1 \pmod{4}$, and by our assumption $F(q) = 1$. Therefore $F(Q) = 1$, and thus

$$1 = F(m^2 + 1) = F(PQ) = F(P)F(Q) = F(P).$$

This proves (2.6) and completes the proof of Lemma 1. ■

Lemma 2. *If $F \in \mathcal{M}^*$ satisfies (1.1) and $bc \neq 0$, then*

$$F(2)F(3)F(5) \neq 0.$$

Proof. (i) The proof of $x = F(2) \neq 0$.

From (2.4), we have

$$E_3 = xF(5) - bx^3 - c = 0,$$

which, together with $c \neq 0$, implies $x = F(2) \neq 0$.

(ii) The proof of $y = F(3) \neq 0$.

Assume that $y = 0$. Then from E_2 and E_4 in (2.4), we obtain:

$$(2.7) \quad F(5) = F(17) = c.$$

Consequently,

$$\begin{aligned} c^3 &= F(5)F(17)^2 = F(1445) = F(38^2 + 1) = bF(38^2 - 1) + c = \\ &= bF(3)F(13)F(37) + c = c, \end{aligned}$$

which gives

$$(2.8) \quad c^2 = 1.$$

Thus, we have

$$(2.9) \quad x = xc^2 = F(2)F(5)^2 = F(50) = bF(48) + c = bx^4y + c = c.$$

On the other hand, from (1.1) and (2.8)–(2.9), we have

$$c^2 = F(2)F(5) = F(10) = F(3^2 + 1) = bF(3^2 - 1) + c = bF(2)^3 + c = bc^3 + c.$$

This with (2.8) implies $c = b + 1$, and hence

$$1 = c^2 = (b + 1)^2 = b(b + 2) + 1 \Rightarrow b(b + 2) = 0.$$

Since $b \neq 0$, it follows that

$$(2.10) \quad b = -2 \quad \text{and} \quad c = b + 1 = -1.$$

Now, from (2.7) and (2.10), we have

$$(2.11) \quad x = F(2) = -1, F(5) = -1 \quad \text{and} \quad F(17) = -1.$$

Now from (2.4), we compute:

$$\begin{aligned} 0 = E_5 &= xF(13) - bx^3y - c = -F(13) + 1, \\ 0 = E_{17} &= xF(5)F(29) - bx^5y^2 - c = F(29) + 1, \\ 0 = E_{57} &= xF(5)^2F(13) - bx^4F(7)F(29) - c = \\ &= -F(13) + 2F(7)F(29) + 1. \end{aligned}$$

From this, we get

$$F(13) = 1, \quad F(29) = -1 \quad \text{and} \quad F(7) = 0.$$

Now consider:

$$\begin{aligned} 0 = E_6 &= F(37) - bF(5)F(7) - c = F(37) + 1 \quad \implies \quad F(37) = -1, \\ 0 = E_{31} &= xF(13)F(37) - bx^6yF(5) - c = -F(37) + 1 \quad \implies \quad F(37) = 1, \end{aligned}$$

which are impossible. Therefore, our assumption $y = 0$ leads to a contradiction, so $y \neq 0$.

(iii) The proof of $F(5) \neq 0$.

Assume that $xy = F(2)F(3) \neq 0$, but $F(5) = 0$. Then from (2.4), we get:

$$\begin{aligned} (2.12) \quad E_2 &= -by - c = 0, \\ E_3 &= -bx^3 - c = 0, \\ E_7 &= -bx^4y - c = 0, \\ E_8 &= -by^2F(7) - c = 0, \\ E_{41} &= xF(29)^2 - c = 0, \\ E_{57} &= -bx^4F(7)F(29) - c = 0. \end{aligned}$$

From E_3 and E_2 , we get:

$$c = -bx^3, \quad y = -\frac{c}{b} = x^3.$$

Using E_7 :

$$E_7 = -bx^3(x^4 - 1) = 0 \implies x^4 = 1.$$

Also

$$E_8 = -by^2F(7) - c = -bx^3(x^3F(7) - 1) = 0 \implies F(7) = \frac{1}{x^3} = x.$$

Thus, we have

$$E_{57} = -bx^4F(7)F(29) - c = -bx^3(x^2F(29) - 1) = 0,$$

which implies

$$F(29) = x^2$$

and

$$E_{41} = xF(29)^2 - c = x^3(x^2 + b) = 0.$$

Hence

$$b = -x^2.$$

Therefore, we have

$$x^4 = 1, \quad b = -x^2, \quad c = x^5, \quad y = x^3, \quad F(7) = x, \quad F(29) = x^2.$$

Now compute:

$$\begin{aligned} L_2 &= bx^6yF(5) - b^2x^3yF(5)F(7) - bcF(5)F(7) - bcx^3y + \\ &\quad + c - c^2 = x^5(x^8 - x^5 + 1) = x(-x + 2) = 0, \end{aligned}$$

which leads to $x = F(2) = 2$. This shows that $1 = x^4 = F(2)^4 = 2^4 = 16$, which is impossible.

Thus, Lemma 2 is proved. ■

Lemma 3. *If $F \in \mathcal{M}^*$ satisfies (1.1) and $bc \neq 0$, then*

$$\begin{aligned} &\left(b, c, x, y, F(5), F(7), F(11), F(13), F(17) \right) \in \\ &\in \left\{ (b, -b + 1, 1, 1, 1, 1, 1, 1), (1, 2, 2, 3, 5, 7, 11, 13, 17) \right\}. \end{aligned}$$

Proof. Assume that $bc \neq 0$. Then Lemma 2 implies that $bcF(2)F(3)F(5) \neq 0$.

By applying (2.4) concerning $E(2)$, $E(4)$, $E(5)$, $E(8)$, $E(17)$, $E(21)$, $E(6)$, we obtain the following expressions:

$$\begin{aligned} (2.13) \quad F(5) &= bF(3) + c = by + c, \\ F(17) &= bF(3)F(5) + c = b^2y^2 + bcy + c, \\ F(13) &= \frac{bF(2)^3F(3) + c}{F(2)} = \frac{bx^3y + c}{x}, \\ F(7) &= \frac{F(5)F(13) - c}{bF(3)^2} = \frac{b^2x^3y^2 + bcx^3y + bcy + c^2 - cx}{bxy^2}, \end{aligned}$$

$$\begin{aligned}
F(29) &= \frac{bF(2)^5F(3)^2 + c}{F(2)F(5)} = \frac{bx^5y^2 + c}{x(by + c)}, \\
F(11) &= \frac{F(2)F(13)F(17) - c}{bF(2)^3F(5)} = \\
(2.13) \quad &= \frac{b^3x^3y^3 + b^2cx^3y^2 + bcx^3y + b^2cy^2 + bc^2y + c^2 - c}{bx^3(by + c)}, \\
F(37) &= bF(5)F(7) + c = \\
&= \frac{b^3x^3y^3 + 2b^2cx^3y^2 + bc^2x^3y + b^2cy^2 + 2bc^2y - bcxy + cxy^2 + c^3 - c^2x}{xy^2}.
\end{aligned}$$

From (1.1), we also have

$$(2.14) \quad E(n) = F(n^2 + 1) - bF(n^2 - 1) - c = 0 \quad \text{for every } n \in \mathbb{N}_1.$$

Thus we infer from (2.13) and (2.14) that

$$\begin{aligned}
E(3) &= F(10) - bF(2)F(4) - c = -bx^3 + bxy + cx - c = 0, \\
E(12) &= F(5)F(29) - bF(11)F(13) - c = \\
&= -\frac{1}{x^4(by + c)} \left(b^4x^6y^4 + b^3cx^6y^3 - b^2x^8y^3 - bcx^8y^2 + b^2cx^6y^2 + \right. \\
&\quad + 2b^3cx^3y^3 + 2b^2c^2x^3y^2 + 2bc^2x^3y + bcx^4y + b^2c^2y^2 - \\
&\quad \left. - 2bcx^3y + c^2x^4 + bc^3y - c^2x^3 + c^3 - c^2 \right) = 0.
\end{aligned}$$

By applying the above relations with $n \in \{2, 3, 8\}$, we infer from (2.3) that

$$\begin{aligned}
L(2) &= bF(2)F(3)F(8) - b^2F(1)F(2)F(3)F(4) - bcF(2)F(4) - bcF(3) - \\
&\quad - c^2 + c = bF(2)^4F(3) - b^2F(2)^3F(3) - bcF(2)^3 - bcF(3) - c^2 + c = \\
&= bx^4y - b^2x^3y - bcx^3 - bcy - c^2 + c = 0, \\
L(3) &= bF(3)F(4)F(14) - b^2F(2)F(3)F(4)F(5) - bcF(3)F(5) - bcF(2)F(4) - \\
&\quad - c^2 + c = bF(3)F(2)^3F(7) - b^2F(2)^3F(3)F(5) - bcF(3)F(5) - \\
&\quad - bcF(2)^3 - c^2 + c = -\frac{1}{y} \left(b^3x^3y^3 - b^2x^5y^2 + b^2cx^3y^2 - bcx^5y + b^2cy^3 + \right. \\
&\quad \left. + bcx^3y + bc^2y^2 - bcx^2y - c^2x^2 + cx^3 + c^2y - cy \right) = 0, \\
L(8) &= bF(8)F(9)F(74) - b^2F(7)F(8)F(9)F(10) - bcF(8)F(10) - \\
&\quad - bcF(7)F(9) - c^2 + c = bF(2)^4F(3)^2F(37) - b^2F(7)F(2)^4F(3)^2F(5) - \\
&\quad - bcF(2)^4F(5) - bcF(7)F(3)^2 - c^2 + c = \\
&= \frac{-c}{x} \left(b^2x^5y - bx^5y^2 + b^2x^3y^2 + bcx^5 + bcx^3y + bcy + c^2 - x \right) = 0.
\end{aligned}$$

Define:

$$\begin{aligned} W_1 &= -E(3), \\ W_2 &= -L(2), \\ W_3 &= -xL(8)/c, \\ W_4 &= -yL(3), \\ W_5 &= -x^4(by + c)E(12). \end{aligned}$$

Then we have the following system of equations

$$\begin{aligned} W_1 &= bx^3 - bxy - cx + c = 0, \\ W_2 &= b^2x^3y - bx^4y + bcx^3 + bcy + c^2 - c = 0, \\ W_3 &= b^2x^5y - bx^5y^2 + b^2x^3y^2 + bcx^5 + bcx^3y + bcy + c^2 - x = 0, \\ W_4 &= b^3x^3y^3 - b^2x^5y^2 + b^2cx^3y^2 - bcx^5y + b^2cy^3 + bcx^3y + bc^2y^2 - bcx^2y - \\ &\quad - c^2x^2 + cx^3 + c^2y - cy = 0, \\ W_5 &= b^4x^6y^4 + b^3cx^6y^3 - b^2x^8y^3 - bcx^8y^2 + b^2cx^6y^2 + 2b^3cx^3y^3 + 2b^2c^2x^3y^2 + \\ &\quad + 2b^2c^2x^3y^2 + 2bc^2x^3y + bcx^4y + b^2c^2y^2 - 2bcx^3y + c^2x^4 + bc^3y - \\ &\quad - c^2x^3 + c^3 - c^2 = 0. \end{aligned}$$

These yield the following system of five polynomial equations in variables b, c, x, y . Using a computer algebra system (Maple), we find that the system has the following solutions:

$$(b, c, x, y) \in \{(b, 0, 0, y), (0, 1, 1, y), (b, -b + 1, 1, 1), (1, 2, 2, 3), (-1, 1, 1, 1), (-2, -1, -1, 0)\}.$$

From these, the only tuples consistent with $bcF(2)F(3)F(5) = bcxy(by + c) \neq 0$ are

$$(b, c, x, y) \in \{(b, -b + 1, 1, 1), (1, 2, 2, 3)\}.$$

Substituting these into the earlier formulas gives the corresponding function values:

$$\begin{aligned} &\left(b, c, x, y, F(5), F(7), F(11), F(13), F(17)\right) \in \\ &\in \left\{(b, -b + 1, 1, 1, 1, 1, 1, 1, 1), (1, 2, 2, 3, 5, 7, 11, 13, 17)\right\}, \end{aligned}$$

completing the proof of Lemma 3. ■

3. Proof of Theorem 1

Proof of Theorem 1 (a). If $b = 0$ and $c \neq 0$, then $F(n^2 + 1) = c$ for every $n \in \mathbb{N}$, and so Theorem 1 (a) follows from Lemma 1. ■

Proof of Theorem 1 (b). Assume that $F \in \mathcal{M}^*$ satisfies (1.1) and $bc \neq 0$. Then, by using Lemma 2 and Lemma 3, we have $bcF(2)F(3)F(5) \neq 0$. Thus, by using Lemma 3, we will prove our theorem by considering two separate cases.

Case 1: $(b, c, x, y, F(5), F(7), F(11), F(13), F(17)) = (b, -b+1, 1, 1, 1, 1, 1, 1)$

Case 2: $(b, c, x, y, F(5), F(7), F(11), F(13), F(17)) = (1, 2, 2, 3, 5, 7, 11, 13, 17)$.

Proof of Case 1. In this case, we can assume that $F(n) = 1$ for every $n \leq P$, where $P > 17$. We aim to prove that $F(P) = 1$.

It is clear from our assumptions that $F(P) = 1$ if $P \notin \mathcal{P}$. So we now assume $P \in \mathcal{P}$.

(a) We first prove that $F(P) = 1$ if $P \in \mathcal{H}$.

(a1). Let $P \in \mathcal{H}$ and $\left(\frac{-1}{P}\right) = 1$.

Then $P \equiv 1 \pmod{4}$ and similarly as in the proof of Lemma 1, there is $m, Q \in \mathbb{N}, m < P$ such that

$$m^2 + 1 = PQ.$$

Clearly, $Q < P$. From our assumptions, $F(Q) = 1$ and

$$F(m-1) = 1, \quad F(m+1) = F(2)F\left(\frac{m+1}{2}\right) = 1.$$

Thus we have

$$F(m^2 + 1) = F(PQ) = F(P)F(Q) = F(P),$$

and from equation (1.1)

$$F(m^2 + 1) = bF(m-1)F(m+1) + 1 - b = b + 1 - b = 1.$$

Therefore, we proved that $F(P) = 1$ if $\left(\frac{-1}{P}\right) = 1$.

(a2). Now suppose $P \in \mathcal{H}$ and $\left(\frac{-7}{P}\right) = 1$.

Then there exist $m, Q \in \mathbb{N}, m < P$ such that

$$m^2 + m + 2 = PQ,$$

since $\left(\frac{-7}{P}\right) = 1$ and

$$m^2 + m + 2 \equiv 0 \pmod{P} \iff (2m+1)^2 \equiv -7 \pmod{P}.$$

Clearly, $m < P$, $Q < P$ and $m \neq P-1, m \neq P-2$. So we have

$$F(Q) = 1, \quad F(m-1) = 1, \quad F(m) = 1, \quad F(m+1) = 1, \quad F(m+2) = 1.$$

Now apply equation (2.3), we have

$$\begin{aligned} L_m &:= bF(m)F(m+1)F(m^2+m+2) - \\ &\quad - b^2F(m-1)F(m)F(m+1)F(m+2) - \\ &\quad - bcF(m)F(m+2) - bcF(m-1)F(m+1) + c - c^2 = 0. \end{aligned}$$

Substituting all values as 1:

$$L_m = bF(P) - b^2 - 2bc + c - c^2 = 0.$$

Recall $c = 1 - b$, we have

$$L_m = bF(P) - b^2 - 2b(1-b) + (1-b) - (1-b)^2 = bF(P) - b = 0,$$

which implies $F(P) = 1$.

Therefore, the theorem is proved for all $p \in \mathcal{H}$.

(b) Now we prove that $F(P) = 1$ if $P \in \mathcal{K}$.

To apply Conjecture 2, by using $P > p_2 = 3$, we first verify the special cases:

$$F(P) = 1 \quad \text{if } P \in \{p_{11} = 31, p_{15} = 47, p_{39} = 167\}.$$

(b1) If $P = p_{11} = 31$, then from E_9 and E_{32} , we compute:

$$F(41) = \frac{bF(2)^4F(5) + c}{F(2)} = b + c = 1$$

and

$$F(31) = \frac{F(5)^2F(41) - c}{bF(3)F(11)} = \frac{1 - c}{b} = 1.$$

(b2) If $P = p_{15} = 47$, then $F(n) = 1$ for every $n < 47$. Thus we infer from E_{27} and E_{46} that

$$F(73) = \frac{bF(2)^3F(13)F(7) + c}{F(2)F(5)} = b + c = 1$$

and

$$F(47) = \frac{F(29)F(73) - c}{bF(3)^2F(5)} = \frac{1 - c}{b} = 1.$$

Consequently

$$F(P) = 1 \quad \text{for } P = 47.$$

(b3) If $P = p_{39} = 167$, then $F(n) = 1$ for every $n < 167$. Thus we infer from E_{107} , E_{8351} that

$$E(107) = F(2)F(5)^2F(229) - bF(2)^3F(53)F(3)^3 - c = F(229) - 1 = 0$$

and

$$\begin{aligned} E(8351) &= F(53)F(2)F(13)^2F(17)F(229) - \\ &\quad - bF(2)F(5)^2F(167)F(29)F(2)^5F(3)^2 - c = \\ &= F(229) - bF(167) - c = 0. \end{aligned}$$

Consequently

$$F(229) = 1 \quad \text{and} \quad F(167) = F(P) = 1.$$

Now we use Conjecture 2. Since $F(n) = 1$ for every $n \leq p_{39} = 167$, we infer from Conjecture 2 that there exists a positive integer ℓ such that

$$P \mid \ell^2 - 1 \quad \text{and} \quad P \left(\frac{(\ell^2 + 1)(\ell^2 - 1)}{P} \right) < P.$$

Thus, it follows from our assumptions that

$$F(\ell^2 + 1) = 1, F(\ell^2 - 1) = F\left(\frac{\ell^2 - 1}{P}\right)F(P) = F(P),$$

which with (1.1) imply that

$$1 = F(\ell^2 + 1) = bF(\ell^2 - 1) + c = bF(P) + c = bF(P) + 1 - b.$$

Therefore $F(P) = 1$ and we proved that $(F, b, c) = (\mathbb{E}, b, -b + 1)$.

Proof of Case 2. Assume that

$$(b, c, x, y, F(5), F(7), F(11), F(13), F(17)) = (1, 2, 2, 3, 5, 7, 11, 13, 17).$$

Similar way as in the proof of Case 1, we can deduce that $F(n) = n$ for every $n \in \mathbb{N}$. Thus, we have $(F, b, c) = (\mathbb{I}, 1, 2)$.

This completes the proof of Theorem 1. ■

4. Remark

Conjecture 3. Let H be a complex valued function defined on the set of Gaussian integers satisfying the equations

$$H(\alpha\beta) = H(\alpha)H(\beta) \quad \text{for every } \alpha, \beta \in \mathbb{Z}[i]$$

$$H(\alpha^2 + 1) = F(\alpha^2 - 1) + 2 \quad \text{for every } \alpha \in \mathbb{Z}[i].$$

Then either $H(\alpha) = \alpha$ or $H(\alpha) = \bar{\alpha}$ for every $\alpha \in \mathbb{Z}[i]$, where $\bar{\alpha}$ is the conjugate of α .

This conjecture is weaker than the assertion (b) in Theorem 1.

References

- [1] **Kátaı, I., B.M.M. Khanh and B.M. Phong**, On the equation $F(n^2 + m^2 + k) = H(n) + H(m) + K$, *Thang Long Journal of Science: Mathematics and Mathematical Sciences*, **1**(4) (2023), 135–151. <https://science.thanglong.edu.vn/index.php/volc/article/view/54>
- [2] **Kátaı, I., B.M.M. Khanh and B.M. Phong**, On the equation $G(n) = F(n^2 - 1) + D$, *Ann. Univ. Sci. Budapest., Sect. Comp.*, **53** (2022), 159–173. <https://doi.org/10.71352/ac.53.159>
- [3] **Kátaı, I., B.M.M. Khanh and B.M. Phong**, On the equation $F(n^3) = F(n^3 - 1) + D$ and some conjectures, *Publ. Math. Debrecen*, **103** (2023), 257–267. <https://doi.org/10.5486/PMD.2023.9656>
- [4] **Spiro, C.**, Additive uniqueness sets for arithmetic functions, *J. Number Theory*, **42** (1992), 232–246. [https://doi.org/10.1016/0022-314X\(92\)90022-H](https://doi.org/10.1016/0022-314X(92)90022-H)

Imre Kátaı, Bui Minh Mai Khanh and Bui Minh Phong

ELTE Eötvös Loránd University

Faculty of Informatics

Department of Computer Algebra

H-1117 Budapest

Pázmány Péter sétány 1/C

Hungary

katai@inf.elte.hu

mbuiminh@yahoo.com

bui@inf.elte.hu

